

Table 6.28A Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2010 and 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	10,929	10,416	19,847	19,080	15,877	16,795	23,011	21,934
Cigarettes	9,231	8,716	16,252	15,437	13,092	13,611	19,761	19,054
Smokeless Tobacco	1,994	1,850	2,784	2,598	1,601	1,434	2,499	2,360
Cigars	2,901 ^a	2,410	4,793	4,268	2,750 ^a	3,399	2,830	2,787
Pipe Tobacco	316	316	569	662	501	554	769	606

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.28B Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2010 and 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	64.7	65.7	47.7 ^a	44.9	21.8	22.4	18.8 ^a	17.7
Cigarettes	54.6	54.9	39.0 ^a	36.3	18.0	18.1	16.2	15.3
Smokeless Tobacco	11.8	11.7	6.7	6.1	2.2	1.9	2.0	1.9
Cigars	17.2	15.2	11.5 ^a	10.0	3.8 ^a	4.5	2.3	2.2
Pipe Tobacco	1.9	2.0	1.4	1.6	0.7	0.7	0.6	0.5

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.29A Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2010 and 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	317	298	819	765	359	390	1,118	1,037
Cigarettes	285	249	660	615	277	307	818	768
Smokeless Tobacco	64	88	201	157	66	65	239	207
Cigars	121	127	288	298	105	136	275	297
Pipe Tobacco	26	29	51	82	21	17	57	55

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.29B Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2010 and 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	77.6	78.7	54.5	52.0	25.5	26.4	5.3	4.8
Cigarettes	69.7	65.8	43.9	41.8	19.7	20.8	3.9	3.5
Smokeless Tobacco	15.7	23.1	13.4	10.7	4.7	4.4	1.1	1.0
Cigars	29.5	33.4	19.2	20.3	7.4	9.2	1.3	1.4
Pipe Tobacco	6.3	7.6	3.4 ^a	5.6	1.5	1.1	0.3	0.3

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.30A Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2010 and 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	3,455 ^b	3,058	5,171	5,107	2,323	2,311	2,985	3,056
Cigarettes	2,950 ^a	2,646	4,296	4,301	1,897	1,933	2,559	2,607
Smokeless Tobacco	758 ^a	605	828	758	220	164	377	341
Cigars	1,158	1,017	1,445	1,331	574	663	681	714
Pipe Tobacco	148	148	258	249	96	128	116	110

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.30B Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2010 and 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	74.8	73.4	56.3 ^a	53.8	32.6	32.3	22.7	22.7
Cigarettes	63.9	63.5	46.8	45.3	26.6	27.0	19.5	19.3
Smokeless Tobacco	16.4	14.5	9.0	8.0	3.1	2.3	2.9	2.5
Cigars	25.1	24.4	15.7	14.0	8.1	9.3	5.2	5.3
Pipe Tobacco	3.2	3.5	2.8	2.6	1.4	1.8	0.9	0.8

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.31A Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2010 and 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	7,156	7,059	13,858	13,209	13,196	14,094	18,908	17,841
Cigarettes	5,996	5,821	11,296	10,521	10,918	11,371	16,384	15,679
Smokeless Tobacco	1,172	1,158	1,756	1,683	1,315	1,205	1,884	1,812
Cigars	1,623	1,267	3,060	2,639	2,071 ^a	2,600	1,874	1,777
Pipe Tobacco	142	140	259	331	384	409	596	441

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.31B Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2010 and 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	60.3	62.4	44.8	41.9	20.5	21.2	21.5	20.0
Cigarettes	50.5	51.4	36.5 ^a	33.4	17.0	17.1	18.6	17.6
Smokeless Tobacco	9.9	10.2	5.7	5.3	2.0	1.8	2.1	2.0
Cigars	13.7	11.2	9.9	8.4	3.2	3.9	2.1	2.0
Pipe Tobacco	1.2	1.2	0.8	1.0	0.6	0.6	0.7	0.5

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.