

Table 6.47A Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	1,978	1,947	271	235	510	495	1,198	1,217
From More Than One Doctor	219	204	31	29	62	46	125	128
Wrote Fake Prescription	24	24	6	6	4	11	14	7
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	60	50	19	13	22	13	19	24
From Friend or Relative for Free	6,189	5,822	594	559	2,059 ^b	1,796	3,536	3,466
Bought from Friend or Relative	1,108 ^a	1,306	118	113	520	513	469 ^b	679
Took from Friend or Relative without Asking	567	473	168 ^b	123	163	148	236	202
Bought from Drug Dealer or Other Stranger	541 ^a	423	49	46	255	222	237	156
Bought on the Internet	49	27	1	0	1	3	46	23
Some Other Way	473	463	75	88	127	128	271	247

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.47B Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	17.7	18.1	20.3	19.3	13.7	14.7	19.5	19.8
From More Than One Doctor	2.0	1.9	2.4	2.4	1.7	1.4	2.0	2.1
Wrote Fake Prescription	0.2	0.2	0.4	0.5	0.1	0.3	0.2	0.1
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.5	0.5	1.4	1.1	0.6	0.4	0.3	0.4
From Friend or Relative for Free	55.2	54.2	44.6	46.1	55.3	53.2	57.5	56.4
Bought from Friend or Relative	9.9 ^b	12.2	8.9	9.4	14.0	15.2	7.6 ^b	11.0
Took from Friend or Relative without Asking	5.1	4.4	12.6 ^a	10.2	4.4	4.4	3.8	3.3
Bought from Drug Dealer or Other Stranger	4.8	3.9	3.7	3.8	6.8	6.6	3.9	2.5
Bought on the Internet	0.4	0.3	0.1	0.0	0.0	0.1	0.8	0.4
Some Other Way	4.2	4.3	5.6	7.2	3.4	3.8	4.4	4.0

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.48A Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	4,408	4,243	303	282	1,324 ^a	1,165	2,781	2,795
From More Than One Doctor	237 ^a	160	26	31	67	70	143 ^b	58
Wrote Fake Prescription	12	17	0	1	1	3	11	14
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	11	18	3	5	8	11	*	2
From Friend or Relative for Free	340	286	56	41	150	123	134	122
Bought from Friend or Relative	237	202	32	29	121	117	84	56
Took from Friend or Relative without Asking	91	92	43	43	26	38	22	11
Bought from Drug Dealer or Other Stranger	103	96	15	17	61	51	27	28
Bought on the Internet	9	9	1	0	3	1	6	8
Some Other Way	63	77	15	23	29	18	19	36

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.48B Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	80.0	81.6	61.3	59.5	73.9	73.0	86.2	89.3
From More Than One Doctor	4.3	3.1	5.3	6.6	3.8	4.4	4.4 ^a	1.9
Wrote Fake Prescription	0.2	0.3	0.1	0.2	0.1	0.2	0.3	0.5
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2	0.3	0.5	1.1	0.5	0.7	*	0.1
From Friend or Relative for Free	6.2	5.5	11.4	8.7	8.4	7.7	4.1	3.9
Bought from Friend or Relative	4.3	3.9	6.4	6.0	6.8	7.3	2.6	1.8
Took from Friend or Relative without Asking	1.6	1.8	8.7	9.1	1.4	2.4	0.7	0.3
Bought from Drug Dealer or Other Stranger	1.9	1.9	3.1	3.7	3.4	3.2	0.8	0.9
Bought on the Internet	0.2	0.2	0.1	0.0	0.2	0.0	0.2	0.2
Some Other Way	1.1	1.5	3.1	4.9	1.6	1.1	0.6	1.2

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.49A Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	1,266	1,146	131	107	273	284	862	755
From More Than One Doctor	238	194	20	24	66	46	152	123
Wrote Fake Prescription	19	27	6	4	3	4	10	19
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	41	49	17 ^a	6	11	15	13	29
From Friend or Relative for Free	2,750	2,565	240	234	905	812	1,605	1,519
Bought from Friend or Relative	1,163	1,127	98	79	471	421	594	628
Took from Friend or Relative without Asking	481	498	108 ^b	71	162	179	212	247
Bought from Drug Dealer or Other Stranger	573	514	64	47	255	264	253	204
Bought on the Internet	49	40	0	1	8	15	40	24
Some Other Way	232	254	44	37	79	77	108	140

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.49B Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	27.9	26.7	26.0	23.3	19.3	22.4	32.9	29.4
From More Than One Doctor	5.2	4.5	3.9	5.3	4.7	3.6	5.8	4.8
Wrote Fake Prescription	0.4	0.6	1.1	0.8	0.2	0.3	0.4	0.7
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.9	1.1	3.3 ^a	1.2	0.8	1.2	0.5	1.1
From Friend or Relative for Free	60.5	59.7	47.7	50.8	63.6	64.0	61.2	59.2
Bought from Friend or Relative	25.6	26.3	19.4	17.1	33.2	33.2	22.7	24.5
Took from Friend or Relative without Asking	10.6	11.6	21.5 ^a	15.5	11.4	14.2	8.1	9.6
Bought from Drug Dealer or Other Stranger	12.6	12.0	12.8	10.2	18.0	20.8	9.7	8.0
Bought on the Internet	1.1	0.9	0.0	0.2	0.6	1.2	1.5	0.9
Some Other Way	5.2	6.1	9.2	8.3	5.7	6.2	4.2	5.6

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.50A Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	1,946	1,814	126	111	606	534	1,214	1,169
From More Than One Doctor	263 ^a	166	25	25	80	70	158 ^a	72
Wrote Fake Prescription	32	37	4 ^a	0	9	4	19	33
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	28	26	6 ^a	2	12	13	10	11
From Friend or Relative for Free	355	328	54 ^b	29	138	129	163	170
Bought from Friend or Relative	311	294	41	27	135	142	135	125
Took from Friend or Relative without Asking	102	98	28	23	37	45	37	30
Bought from Drug Dealer or Other Stranger	211	203	26	20	104	100	81	83
Bought on the Internet	13	6	3	0	6	4	4	1
Some Other Way	75	79	16	16	27	25	32	39

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.50B Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	78.0	77.5	61.3	55.1	74.1	71.4	82.6	83.9
From More Than One Doctor	10.7 ^a	7.1	12.4	12.4	9.8	9.4	11.0 ^a	5.2
Wrote Fake Prescription	1.3	1.6	1.9 ^a	0.1	1.2	0.5	1.3	2.4
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.1	1.1	3.0 ^a	0.8	1.5	1.8	0.7	0.8
From Friend or Relative for Free	14.5	14.1	26.7 ^b	14.7	17.0	17.5	11.3	12.2
Bought from Friend or Relative	12.7	12.6	20.4 ^a	13.5	16.7	19.2	9.4	9.0
Took from Friend or Relative without Asking	4.2	4.2	13.8	11.5	4.5	6.1	2.6	2.2
Bought from Drug Dealer or Other Stranger	8.6	8.7	13.0	9.8	12.8	13.5	5.6	6.0
Bought on the Internet	0.5	0.2	1.4	0.1	0.8	0.6	0.3	0.1
Some Other Way	3.1	3.5	8.4	8.0	3.4	3.4	2.2	2.8

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.51A Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	460 ^b	661	28	31	113	116	318 ^b	514
From More Than One Doctor	56	51	5	5	17	15	34	31
Wrote Fake Prescription	3	4	2	3	1	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	13	14	4	1	8	9	*	4
From Friend or Relative for Free	2,952	2,901	188	192	882	830	1,881	1,879
Bought from Friend or Relative	669	669	70	61	331	293	268	314
Took from Friend or Relative without Asking	199	165	43	33	71	71	85	61
Bought from Drug Dealer or Other Stranger	300	296	41	39	156	159	103	98
Bought on the Internet	71	17	1	1	2	2	67	*
Some Other Way	190	120	15	18	61	42	114	60

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.51B Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	9.4 ^b	13.5	7.1	8.0	6.9	7.6	11.1 ^b	17.3
From More Than One Doctor	1.1	1.0	1.3	1.3	1.0	1.0	1.2	1.0
Wrote Fake Prescription	0.1	0.1	0.5	0.7	0.1	0.1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3	0.3	1.1	0.4	0.5	0.6	*	0.1
From Friend or Relative for Free	60.1	59.2	47.3	50.0	53.7	53.9	65.5	63.2
Bought from Friend or Relative	13.6	13.7	17.5	16.0	20.1	19.0	9.3	10.6
Took from Friend or Relative without Asking	4.0	3.4	10.7	8.6	4.3	4.6	3.0	2.0
Bought from Drug Dealer or Other Stranger	6.1	6.0	10.4	10.3	9.5	10.3	3.6	3.3
Bought on the Internet	1.4	0.4	0.4	0.2	0.1	0.2	2.3	*
Some Other Way	3.9	2.4	3.8	4.6	3.7	2.7	4.0	2.0

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.52A Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	1,977	2,141	77	82	509	524	1,392	1,534
From More Than One Doctor	115 ^a	69	9	8	25	35	82 ^b	26
Wrote Fake Prescription	5	2	2	*	1	2	2	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	9	6	1	*	5	1	3	5
From Friend or Relative for Free	233	221	25	22	109	85	99	115
Bought from Friend or Relative	126	93	17	14	74 ^b	41	34	38
Took from Friend or Relative without Asking	27	25	16	8	9	13	2	4
Bought from Drug Dealer or Other Stranger	45	62	6	12	29	32	10	18
Bought on the Internet	5	7	*	*	3	1	2	5
Some Other Way	28	21	6	8	7	4	15	10

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.52B Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	77.0 ^a	80.9	48.0	53.1	66.2	71.1	84.9	87.4
From More Than One Doctor	4.5 ^a	2.6	5.4	5.0	3.2	4.7	5.0 ^b	1.5
Wrote Fake Prescription	0.2	0.1	1.4	*	0.1	0.2	0.1	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.4	0.2	0.9	*	0.6	0.1	0.2	0.3
From Friend or Relative for Free	9.1	8.4	15.9	14.0	14.1	11.5	6.0	6.5
Bought from Friend or Relative	4.9	3.5	10.9	9.2	9.7 ^b	5.6	2.1	2.2
Took from Friend or Relative without Asking	1.1	0.9	10.2	5.4	1.1	1.7	0.1	0.2
Bought from Drug Dealer or Other Stranger	1.7	2.3	3.6	7.8	3.7	4.4	0.6	1.0
Bought on the Internet	0.2	0.3	*	*	0.4	0.2	0.1	0.3
Some Other Way	1.1	0.8	3.7	4.9	0.9	0.5	0.9	0.6

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.53A Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	234 ^a	334	12	10	59	63	162 ^a	261
From More Than One Doctor	33	14	*	3	7	8	24 ^a	4
Wrote Fake Prescription	1	3	1	1	*	2	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	5	21	*	1	2	4	*	*
From Friend or Relative for Free	1,059	1,040	55	45	289	255	714	740
Bought from Friend or Relative	361	376	22	29	149	130	190	217
Took from Friend or Relative without Asking	93	91	21 ^a	9	35	43	37	39
Bought from Drug Dealer or Other Stranger	177	146	19	17	80	84	77	45
Bought on the Internet	*	12	2	0	1	4	*	8
Some Other Way	57	41	6	6	15	9	35	26

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.53B Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	13.1 ^a	18.6	9.8	9.1	10.7	12.5	14.7 ^a	22.1
From More Than One Doctor	1.8	0.8	*	3.0	1.2	1.5	2.2 ^a	0.3
Wrote Fake Prescription	0.1	0.2	1.0	1.2	*	0.4	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3	1.2	*	1.0	0.3	0.9	*	*
From Friend or Relative for Free	59.3	57.9	42.4	41.0	52.5	50.6	64.7	62.6
Bought from Friend or Relative	20.3	20.9	17.4	26.1	27.0	25.8	17.2	18.3
Took from Friend or Relative without Asking	5.3	5.1	16.3 ^a	7.7	6.4	8.5	3.4	3.3
Bought from Drug Dealer or Other Stranger	9.9	8.1	15.1	15.6	14.5	16.7	7.0	3.8
Bought on the Internet	*	0.6	1.6	0.4	0.1	0.7	*	0.6
Some Other Way	3.2	2.3	5.2	6.0	2.8	1.8	3.2	2.2

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.54A Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	724	756	*	*	172	163	530	572
From More Than One Doctor	78	55	*	*	20	20	55	34
Wrote Fake Prescription	1	1	*	*	1	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	3	7	*	*	1	3	*	*
From Friend or Relative for Free	118	132	*	*	57	36	50	87
Bought from Friend or Relative	72	61	*	*	44 ^a	25	16	30
Took from Friend or Relative without Asking	15	23	*	*	3	8	7	*
Bought from Drug Dealer or Other Stranger	35	47	*	*	12	19	19	26
Bought on the Internet	1	6	*	*	1	*	0	*
Some Other Way	21	12	*	*	6	3	*	8

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.54B Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	73.8	77.9	*	*	63.5	70.5	79.8	81.8
From More Than One Doctor	8.0	5.7	*	*	7.6	8.6	8.3	4.9
Wrote Fake Prescription	0.1	0.1	*	*	0.2	0.4	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3	0.7	*	*	0.5	1.3	*	*
From Friend or Relative for Free	12.1	13.6	*	*	21.1	15.8	7.6	12.5
Bought from Friend or Relative	7.3	6.3	*	*	16.2	10.8	2.4	4.3
Took from Friend or Relative without Asking	1.6	2.4	*	*	1.2	3.4	1.1	*
Bought from Drug Dealer or Other Stranger	3.6	4.8	*	*	4.6	8.1	2.9	3.7
Bought on the Internet	0.1	0.7	*	*	0.4	*	0.0	*
Some Other Way	2.2	1.3	*	*	2.1	1.2	*	1.1

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.55A Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Stimulant Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	205	211	25	19	77	65	104	126
From More Than One Doctor	29	13	3	5	7	8	19	*
Wrote Fake Prescription	11	3	3	3	1	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	19	14	7	2	6	4	*	*
From Friend or Relative for Free	1,082	1,065	107	94	481	504	494	466
Bought from Friend or Relative	384	426	38	41	233	265	113	120
Took from Friend or Relative without Asking	120	86	27	18	31	30	62	38
Bought from Drug Dealer or Other Stranger	137	146	24	16	74	61	40	68
Bought on the Internet	33	21	2	1	11	6	21	14
Some Other Way	63	72	11	5	30	24	22	43

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.55B Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Stimulant Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	9.9	10.3	10.0	9.4	8.1	6.8	11.7	14.3
From More Than One Doctor	1.4	0.6	1.4	2.3	0.8	0.8	2.1	*
Wrote Fake Prescription	0.5	0.2	1.4	1.6	0.1	0.0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.9	0.7	2.9	1.0	0.7	0.4	*	*
From Friend or Relative for Free	51.9	51.8	43.4	46.1	50.6	52.1	55.7	52.8
Bought from Friend or Relative	18.4	20.7	15.5	20.0	24.5	27.4	12.7	13.5
Took from Friend or Relative without Asking	5.7	4.2	10.8	8.9	3.2	3.1	7.0	4.3
Bought from Drug Dealer or Other Stranger	6.6	7.1	9.6	8.0	7.8	6.3	4.5	7.7
Bought on the Internet	1.6	1.0	0.8	0.5	1.1	0.6	2.3	1.6
Some Other Way	3.0	3.5	4.3	2.4	3.1	2.5	2.5	4.9

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.56A Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Stimulants ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	702	766	60	48	327	338	315	380
From More Than One Doctor	34	30	4	6	11	20	19	4
Wrote Fake Prescription	*	1	*	*	*	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	2	1	*	3	2	*	*
From Friend or Relative for Free	92	70	10	9	43	43	39	18
Bought from Friend or Relative	51	55	9	6	32	37	10	12
Took from Friend or Relative without Asking	9	12	4	*	4	6	1	*
Bought from Drug Dealer or Other Stranger	28	26	4	4	11	13	*	*
Bought on the Internet	7	4	*	1	1	*	*	4
Some Other Way	17	2	1	1	3	1	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.56B Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Stimulants ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	74.4	79.1	65.1	58.1	75.3	73.6	75.5 ^a	89.2
From More Than One Doctor	3.6	3.1	4.6	7.7	2.4	4.3	4.6	0.8
Wrote Fake Prescription	*	0.1	*	*	*	0.1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.4	0.2	0.7	*	0.8	0.4	*	*
From Friend or Relative for Free	9.8	7.2	11.0	11.3	9.9	9.3	9.4	4.2
Bought from Friend or Relative	5.4	5.7	9.3	7.5	7.3	8.1	2.5	2.8
Took from Friend or Relative without Asking	0.9	1.2	4.2	*	0.9	1.3	0.2	*
Bought from Drug Dealer or Other Stranger	2.9	2.7	3.9	5.1	2.6	2.8	*	*
Bought on the Internet	0.7	0.5	*	0.8	0.1	*	*	0.9
Some Other Way	1.8	0.2	0.6	1.3	0.6	0.2	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.57A Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Stimulant Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	121	151	15	9	41	42	64	*
From More Than One Doctor	31	10	4	3	5	7	*	*
Wrote Fake Prescription	6	3	4	*	1	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	14	3	5	0	4	3	*	*
From Friend or Relative for Free	380	342	40	28	157	165	184	150
Bought from Friend or Relative	205	210	17	18	102	120	86	72
Took from Friend or Relative without Asking	80	54	13	8	19	25	*	20
Bought from Drug Dealer or Other Stranger	93	97	15	12	45	41	33	43
Bought on the Internet	21	10	1	1	4	4	*	5
Some Other Way	22	29	5	4	10	4	*	21

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*) and because respondents who were logically assigned through the editing process as reporting only methamphetamine use were not previously excluded.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.57B Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Stimulant Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	15.2	20.9	17.4	12.7	12.7	13.5	16.7	*
From More Than One Doctor	3.8	1.4	4.8	3.6	1.6	2.4	*	*
Wrote Fake Prescription	0.7	0.4	4.6	*	0.5	0.0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.8	0.4	5.1	0.6	1.3	0.9	*	*
From Friend or Relative for Free	47.8	47.2	44.6	39.7	48.6	53.1	47.9	43.4
Bought from Friend or Relative	25.8	29.0	19.4	26.0	31.7	38.9	22.4	20.8
Took from Friend or Relative without Asking	10.1	7.4	14.9	11.9	5.8	8.3	*	5.8
Bought from Drug Dealer or Other Stranger	11.6	13.4	16.4	17.8	13.8	13.4	8.7	12.4
Bought on the Internet	2.7	1.4	1.1	1.0	1.2	1.3	*	1.5
Some Other Way	2.8	4.2	6.0	6.1	3.2	1.3	*	6.3

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*) and because respondents who were logically assigned through the editing process as reporting only methamphetamine use were not previously excluded.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.58A Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Stimulants ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	240	257	*	*	107	119	*	*
From More Than One Doctor	16	17	*	*	5	12	*	*
Wrote Fake Prescription	*	0	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	1	*	*	*	*	*	*
From Friend or Relative for Free	75 ^a	30	*	*	25	16	*	*
Bought from Friend or Relative	42	22	*	*	13	14	*	*
Took from Friend or Relative without Asking	14	10	*	*	3	7	*	*
Bought from Drug Dealer or Other Stranger	22	18	*	*	8	6	*	*
Bought on the Internet	2	6	*	*	2	*	*	*
Some Other Way	*	2	*	*	0	0	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.58B Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Stimulants ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	68.7 ^a	82.3	*	*	75.0	80.8	*	*
From More Than One Doctor	4.5	5.4	*	*	3.3	8.4	*	*
Wrote Fake Prescription	*	0.0	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	0.5	*	*	*	*	*	*
From Friend or Relative for Free	21.5 ^a	9.9	*	*	17.4	10.9	*	*
Bought from Friend or Relative	12.1	7.4	*	*	9.0	9.6	*	*
Took from Friend or Relative without Asking	3.9	3.4	*	*	2.3	4.6	*	*
Bought from Drug Dealer or Other Stranger	6.2	6.0	*	*	5.7	4.4	*	*
Bought on the Internet	0.5	1.9	*	*	1.1	*	*	*
Some Other Way	*	0.7	*	*	0.3	0.1	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.59A Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	375	356	37	*	141	110	197	204
Bought from Friend or Relative	267	250	11	*	46	43	210	*
Took from Friend or Relative without Asking	36	26	8	2	7	*	*	13
Bought from Drug Dealer or Other Stranger	192	189	11	*	53	48	128	*
Bought on the Internet	6	9	*	*	2	3	*	*
Some Other Way	37	28	1	*	9	6	27	15

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.59B Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	41.0	41.5	51.7	*	54.4	49.6	33.8	36.7
Bought from Friend or Relative	29.3	29.2	15.6	*	17.8	19.5	36.0	*
Took from Friend or Relative without Asking	4.0	3.0	11.0 ^a	3.0	2.8	*	*	2.3
Bought from Drug Dealer or Other Stranger	21.0	22.0	14.9	*	20.6	21.7	21.9	*
Bought on the Internet	0.7	1.0	*	*	0.9	1.5	*	*
Some Other Way	4.0	3.3	1.0	*	3.4	2.9	4.7	2.7

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.60A Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	132	108	*	*	43	35	*	*
Bought from Friend or Relative	99	84	*	*	41	25	*	*
Took from Friend or Relative without Asking	7	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	61	48	*	*	22	20	*	*
Bought on the Internet	*	2	*	*	*	*	*	*
Some Other Way	10	18	*	*	8	9	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.60B Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	42.4	40.1	*	*	36.1	37.9	*	*
Bought from Friend or Relative	31.9	31.4	*	*	34.5	26.5	*	*
Took from Friend or Relative without Asking	2.1	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	19.4	17.8	*	*	18.4	21.0	*	*
Bought on the Internet	*	0.7	*	*	*	*	*	*
Some Other Way	3.3	6.9	*	*	6.6	9.3	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.61A Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	183	166	*	*	*	*	*	*
Bought from Friend or Relative	146	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	9	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	122	101	*	*	*	*	*	*
Bought on the Internet	1	7	*	*	*	*	*	*
Some Other Way	28	9	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.61B Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	49.2	51.1	*	*	*	*	*	*
Bought from Friend or Relative	39.3	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	2.9	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	32.9	31.0	*	*	*	*	*	*
Bought on the Internet	0.2	2.0	*	*	*	*	*	*
Some Other Way	7.9	2.7	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.62A Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.62B Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.63A Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Sedative Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	89	132	13	13	23	21	54	*
From More Than One Doctor	25	*	2	*	*	2	20	*
Wrote Fake Prescription	4	1	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	8	4	0	4	*	*	*	0
From Friend or Relative for Free	365	385	41	36	86	77	238	*
Bought from Friend or Relative	55	53	14	9	16	15	*	29
Took from Friend or Relative without Asking	37	25	14 ^a	4	16	9	8	*
Bought from Drug Dealer or Other Stranger	26	24	5	3	4	5	17	16
Bought on the Internet	*	0	*	0	0	*	*	*
Some Other Way	29	17	1	*	5	3	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.63B Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Sedative Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	13.8	20.1	13.9	16.9	14.3	16.0	13.5	*
From More Than One Doctor	3.8	*	2.2	*	*	1.4	5.1	*
Wrote Fake Prescription	0.7	0.2	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.3	0.7	0.4	5.7	*	*	*	0.1
From Friend or Relative for Free	56.2	58.9	44.9	48.7	53.8	58.5	59.9	*
Bought from Friend or Relative	8.5	8.1	15.0	12.4	10.1	11.7	*	6.4
Took from Friend or Relative without Asking	5.8	3.8	15.6 ^a	5.8	9.8	6.7	1.9	*
Bought from Drug Dealer or Other Stranger	4.0	3.7	5.4	3.6	2.8	3.7	4.2	3.7
Bought on the Internet	*	0.0	*	0.1	0.1	*	*	*
Some Other Way	4.5	2.5	1.1	*	3.3	2.0	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.64A Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Sedatives ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	273	326	*	*	*	*	*	*
From More Than One Doctor	3	6	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	6	2	*	*	*	*	*	*
From Friend or Relative for Free	33 ^a	7	*	*	*	*	*	*
Bought from Friend or Relative	2	5	*	*	*	*	*	*
Took from Friend or Relative without Asking	3	7	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	2	7	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	3	3	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.64B Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Sedatives ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	82.1	89.3	*	*	*	*	*	*
From More Than One Doctor	1.0	1.7	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.9	0.7	*	*	*	*	*	*
From Friend or Relative for Free	10.0 ^a	1.8	*	*	*	*	*	*
Bought from Friend or Relative	0.7	1.5	*	*	*	*	*	*
Took from Friend or Relative without Asking	1.0	1.8	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	0.7	2.0	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	1.0	0.7	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.65A Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Sedative Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	61	*	*	*	*	*	*	*
From More Than One Doctor	19 ^a	0	*	*	*	*	*	*
Wrote Fake Prescription	1	1	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	6	2	*	*	*	*	*	*
From Friend or Relative for Free	141	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	22 ^a	7	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	11	*	*	*	*	*	*
Bought on the Internet	1	0	*	*	*	*	*	*
Some Other Way	7	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.65B Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Sedative Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	22.3	*	*	*	*	*	*	*
From More Than One Doctor	6.9 ^a	0.0	*	*	*	*	*	*
Wrote Fake Prescription	0.3	0.4	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2.2	0.8	*	*	*	*	*	*
From Friend or Relative for Free	51.7	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	8.2 ^a	2.7	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	4.1	*	*	*	*	*	*
Bought on the Internet	0.2	0.1	*	*	*	*	*	*
Some Other Way	2.7	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.66A Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Sedatives ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.66B Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Sedatives ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2010-2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010-2011 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.