

Table 6.11C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2010 and 2011

| Drug | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|--|------------------------------------|------|--|------|------------------------------------|------|--------|------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 |
| ILLCIT DRUGS² | 217 | 214 | 246 | 228 | 247 | 222 | 227 | 225 |
| Marijuana and Hashish | 198 | 182 | 203 | 213 | 219 | 195 | 172 | 183 |
| Cocaine | 70 | 78 | 63 | 59 | 42 | 32 | 37 | 27 |
| Crack | 30 | 26 | 33 | 31 | 32 | 3 | 26 | 10 |
| Heroin | 17 | 20 | 13 | 17 | 27 | 16 | 26 | 32 |
| Hallucinogens | 62 | 33 | 40 | 35 | 22 | 27 | 26 | 22 |
| LSD | 14 | 10 | 12 | 11 | 7 | 6 | 4 | 16 |
| PCP | 3 | 3 | 4 | 9 | * | * | 20 | * |
| Ecstasy | 46 | 26 | 33 | 26 | 17 | 23 | 13 | 10 |
| Inhalants | 25 | 33 | 28 | 27 | 71 | 30 | 23 | 38 |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 104 | 104 | 134 | 105 | 132 | 113 | 146 | 131 |
| Pain Relievers | 94 | 76 | 101 | 95 | 112 | 89 | 114 | 116 |
| OxyContin [®] | 29 | 26 | 33 | 31 | 24 | 19 | 48 | 27 |
| Tranquilizers | 59 | 64 | 75 | 59 | 69 | 58 | 71 | 74 |
| Stimulants ⁴ | 49 | 51 | 45 | 36 | 32 | 28 | 42 | 53 |
| Methamphetamine ⁴ | 21 | 42 | 24 | 17 | 26 | 23 | 31 | 50 |
| Sedatives | 23 | 23 | 54 | 13 | 17 | 21 | 40 | 35 |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 138 | 128 | 150 | 121 | 142 | 122 | 152 | 142 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.11D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2010 and 2011

| Drug | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|--|------------------------------------|------|--|------|------------------------------------|------|--------|------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 |
| ILLCIT DRUGS² | 1.10 | 1.06 | 0.56 | 0.51 | 0.33 | 0.30 | 0.18 | 0.18 |
| Marijuana and Hashish | 1.01 | 0.96 | 0.47 | 0.47 | 0.29 | 0.26 | 0.14 | 0.15 |
| Cocaine | 0.40 | 0.47 | 0.15 | 0.14 | 0.06 | 0.04 | 0.03 | 0.02 |
| Crack | 0.17 | 0.17 | 0.08 | 0.07 | 0.04 | 0.00 | 0.02 | 0.01 |
| Heroin | 0.10 | 0.13 | 0.03 | 0.04 | 0.04 | 0.02 | 0.02 | 0.03 |
| Hallucinogens | 0.35 | 0.21 | 0.10 | 0.08 | 0.03 | 0.04 | 0.02 | 0.02 |
| LSD | 0.08 | 0.07 | 0.03 | 0.03 | 0.01 | 0.01 | 0.00 | 0.01 |
| PCP | 0.02 | 0.02 | 0.01 | 0.02 | * | * | 0.02 | * |
| Ecstasy | 0.27 | 0.17 | 0.08 | 0.06 | 0.02 | 0.03 | 0.01 | 0.01 |
| Inhalants | 0.14 | 0.21 | 0.07 | 0.06 | 0.10 | 0.04 | 0.02 | 0.03 |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 0.60 | 0.62 | 0.32 | 0.25 | 0.18 | 0.15 | 0.12 | 0.11 |
| Pain Relievers | 0.54 | 0.47 | 0.24 | 0.22 | 0.15 | 0.12 | 0.09 | 0.09 |
| OxyContin [®] | 0.17 | 0.16 | 0.08 | 0.07 | 0.03 | 0.03 | 0.04 | 0.02 |
| Tranquilizers | 0.35 | 0.40 | 0.18 | 0.14 | 0.09 | 0.08 | 0.06 | 0.06 |
| Stimulants ⁴ | 0.29 | 0.32 | 0.11 | 0.09 | 0.04 | 0.04 | 0.03 | 0.04 |
| Methamphetamine ⁴ | 0.13 | 0.26 | 0.06 | 0.04 | 0.04 | 0.03 | 0.03 | 0.04 |
| Sedatives | 0.13 | 0.14 | 0.13 | 0.03 | 0.02 | 0.03 | 0.03 | 0.03 |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 0.76 | 0.73 | 0.36 | 0.29 | 0.19 | 0.16 | 0.12 | 0.11 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.11P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2011 Versus 2010

| Drug | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|--|------------------------------------|------|--|------|------------------------------------|------|--------|------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 |
| ILLCIT DRUGS² | 0.7549 | | 0.4477 | | 0.8166 | | 0.6149 | |
| Marijuana and Hashish | 0.5218 | | 0.3237 | | 0.6287 | | 0.1313 | |
| Cocaine | 0.7056 | | 0.7486 | | 0.5110 | | 0.2482 | |
| Crack | 0.6154 | | 0.7489 | | 0.0412 | | 0.0891 | |
| Heroin | 0.8337 | | 0.4227 | | 0.4919 | | 0.3405 | |
| Hallucinogens | 0.0032 | | 0.4650 | | 0.4631 | | 0.8567 | |
| LSD | 0.3166 | | 0.7302 | | 0.8803 | | 0.4432 | |
| PCP | 0.7228 | | 0.2993 | | * | | * | |
| Ecstasy | 0.0531 | | 0.4076 | | 0.6063 | | 0.2339 | |
| Inhalants | 0.4679 | | 0.7067 | | 0.1954 | | 0.7322 | |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 0.6553 | | 0.0056 | | 0.1504 | | 0.7556 | |
| Pain Relievers | 0.1048 | | 0.0439 | | 0.1679 | | 0.7643 | |
| OxyContin [®] | 0.9619 | | 0.4339 | | 0.1616 | | 0.3608 | |
| Tranquilizers | 0.4197 | | 0.1070 | | 0.1429 | | 0.3267 | |
| Stimulants ⁴ | 0.8957 | | 0.0546 | | 0.7628 | | 0.8417 | |
| Methamphetamine ⁴ | 0.0555 | | 0.0569 | | 0.9182 | | 0.3897 | |
| Sedatives | 0.5428 | | 0.0858 | | 0.5011 | | 0.3724 | |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 0.3677 | | 0.0114 | | 0.1453 | | 0.7698 | |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.12C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2010 and 2011

| Drug | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|---|------------------------------------|------|--|------|------------------------------------|------|--------|------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 |
| ILLICIT DRUGS² | 24 | 20 | 41 | 32 | 25 | 30 | 48 | 50 |
| Marijuana and Hashish | 23 | 21 | 39 | 30 | 24 | 29 | 36 | 41 |
| Cocaine | 7 | 6 | 7 | 6 | 5 | 6 | 3 | 4 |
| Crack | 2 | 0 | 0 | 2 | * | * | * | * |
| Heroin | 1 | 1 | 2 | 2 | 1 | 7 | 2 | 1 |
| Hallucinogens | 10 | 9 | 13 | 12 | 8 | 11 | 10 | 12 |
| LSD | 7 | 3 | 5 | 7 | 3 | 1 | 3 | 3 |
| PCP | 1 | 1 | 1 | 1 | * | * | 2 | * |
| Ecstasy | 6 | 7 | 11 | 8 | 7 | 9 | 7 | 6 |
| Inhalants | 8 | 4 | 9 | 9 | 8 | 6 | 19 | 16 |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 12 | 12 | 21 | 16 | 10 | 11 | 28 | 25 |
| Pain Relievers | 12 | 10 | 19 | 15 | 9 | 9 | 25 | 23 |
| OxyContin [®] | 6 | 5 | 6 | 8 | 2 | 1 | 6 | 3 |
| Tranquilizers | 8 | 9 | 8 | 11 | 4 | 6 | 10 | 7 |
| Stimulants ⁴ | 7 | 6 | 10 | 5 | 3 | 6 | 7 | 7 |
| Methamphetamine ⁴ | 3 | 4 | 6 | 4 | 1 | 5 | 3 | 4 |
| Sedatives | 3 | 2 | 4 | 4 | * | 3 | 6 | 3 |
| ILLICIT DRUGS OTHER THAN MARIJUANA² | 16 | 13 | 25 | 21 | 14 | 17 | 34 | 31 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.12D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2010 and 2011

| Drug | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|--|------------------------------------|------|--|------|------------------------------------|------|--------|------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 |
| ILLCIT DRUGS² | 2.78 | 3.04 | 1.85 | 1.64 | 1.53 | 1.70 | 0.22 | 0.22 |
| Marijuana and Hashish | 3.06 | 3.08 | 1.80 | 1.63 | 1.48 | 1.64 | 0.17 | 0.19 |
| Cocaine | 1.69 | 1.52 | 0.45 | 0.40 | 0.33 | 0.42 | 0.01 | 0.02 |
| Crack | 0.45 | 0.03 | 0.02 | 0.14 | * | * | * | * |
| Heroin | 0.32 | 0.22 | 0.12 | 0.12 | 0.08 | 0.49 | 0.01 | 0.00 |
| Hallucinogens | 2.36 | 2.14 | 0.82 | 0.77 | 0.54 | 0.76 | 0.05 | 0.05 |
| LSD | 1.70 | 0.86 | 0.34 | 0.45 | 0.22 | 0.05 | 0.01 | 0.01 |
| PCP | 0.32 | 0.37 | 0.08 | 0.08 | * | * | 0.01 | * |
| Ecstasy | 1.41 | 1.86 | 0.72 | 0.54 | 0.48 | 0.60 | 0.03 | 0.03 |
| Inhalants | 1.78 | 0.93 | 0.58 | 0.58 | 0.59 | 0.43 | 0.09 | 0.08 |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 2.72 | 2.84 | 1.28 | 1.05 | 0.69 | 0.74 | 0.13 | 0.11 |
| Pain Relievers | 2.65 | 2.54 | 1.21 | 0.96 | 0.62 | 0.61 | 0.11 | 0.11 |
| OxyContin [®] | 1.46 | 1.20 | 0.38 | 0.50 | 0.12 | 0.07 | 0.03 | 0.02 |
| Tranquilizers | 1.83 | 2.38 | 0.56 | 0.72 | 0.26 | 0.40 | 0.05 | 0.03 |
| Stimulants ⁴ | 1.72 | 1.61 | 0.67 | 0.34 | 0.22 | 0.41 | 0.03 | 0.03 |
| Methamphetamine ⁴ | 0.83 | 1.03 | 0.42 | 0.26 | 0.10 | 0.33 | 0.01 | 0.02 |
| Sedatives | 0.79 | 0.42 | 0.27 | 0.27 | * | 0.18 | 0.03 | 0.01 |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 3.16 | 3.14 | 1.50 | 1.28 | 0.96 | 1.07 | 0.16 | 0.14 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.12P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2011 Versus 2010

| Drug | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|---|------------------------------------|------|--|------|------------------------------------|------|--------|------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 |
| ILLICIT DRUGS² | 0.9692 | | 0.0913 | | 0.0821 | | 0.4934 | |
| Marijuana and Hashish | 0.3624 | | 0.1346 | | 0.0061 | | 0.0286 | |
| Cocaine | 0.8419 | | 0.9587 | | 0.6310 | | 0.5451 | |
| Crack | 0.0974 | | 0.2233 | | * | | * | |
| Heroin | 0.7303 | | 0.6743 | | 0.2215 | | 0.6492 | |
| Hallucinogens | 0.8830 | | 0.7727 | | 0.7942 | | 0.6915 | |
| LSD | 0.2513 | | 0.4712 | | 0.0590 | | 0.5417 | |
| PCP | 0.9338 | | 0.8847 | | * | | * | |
| Ecstasy | 0.1943 | | 0.1793 | | 0.9306 | | 0.3845 | |
| Inhalants | 0.2481 | | 0.5027 | | 0.1375 | | 0.2955 | |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 0.3250 | | 0.3948 | | 0.5771 | | 0.2890 | |
| Pain Relievers | 0.6908 | | 0.3040 | | 0.6627 | | 0.4613 | |
| OxyContin [®] | 0.5240 | | 0.5392 | | 0.7308 | | 0.4678 | |
| Tranquilizers | 0.1515 | | 0.1601 | | 0.0849 | | 0.1596 | |
| Stimulants ⁴ | 0.3937 | | 0.0697 | | 0.0862 | | 0.4430 | |
| Methamphetamine ⁴ | 0.1784 | | 0.7944 | | 0.2528 | | 0.7941 | |
| Sedatives | 0.5077 | | 0.5848 | | * | | 0.1029 | |
| ILLICIT DRUGS OTHER THAN MARIJUANA² | 0.9289 | | 0.3869 | | 0.9022 | | 0.3036 | |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.13C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2010 and 2011

| Drug | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|--|------------------------------------|------|--|------|------------------------------------|------|--------|------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 |
| ILLCIT DRUGS² | 94 | 97 | 94 | 101 | 58 | 67 | 58 | 57 |
| Marijuana and Hashish | 88 | 86 | 86 | 96 | 55 | 62 | 54 | 50 |
| Cocaine | 30 | 33 | 23 | 24 | 10 | 8 | 7 | 7 |
| Crack | 8 | 3 | 12 | 5 | 5 | 1 | 2 | 1 |
| Heroin | 10 | 7 | 12 | 11 | 6 | 11 | 4 | 8 |
| Hallucinogens | 35 | 22 | 25 | 28 | 16 | 15 | 10 | 12 |
| LSD | 11 | 10 | 10 | 9 | 6 | 3 | 3 | 2 |
| PCP | 2 | 1 | 2 | 6 | * | 1 | * | * |
| Ecstasy | 26 | 17 | 19 | 21 | 15 | 11 | 8 | 8 |
| Inhalants | 10 | 18 | 16 | 14 | 7 | 17 | 6 | 5 |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 40 | 41 | 45 | 40 | 29 | 30 | 28 | 27 |
| Pain Relievers | 34 | 32 | 39 | 36 | 25 | 25 | 25 | 22 |
| OxyContin [®] | 11 | 13 | 18 | 13 | 6 | 8 | 12 | 6 |
| Tranquilizers | 25 | 22 | 23 | 22 | 16 | 17 | 10 | 16 |
| Stimulants ⁴ | 23 | 24 | 20 | 14 | 9 | 11 | 13 | 10 |
| Methamphetamine ⁴ | 5 | 8 | 14 | 6 | 6 | 6 | 4 | 9 |
| Sedatives | 7 | 6 | 5 | 5 | 3 | 6 | 8 | 4 |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 57 | 55 | 50 | 52 | 32 | 37 | 30 | 31 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.13D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2010 and 2011

| Drug | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|--|------------------------------------|------|--|------|------------------------------------|------|--------|------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 |
| ILLCIT DRUGS² | 1.24 | 1.33 | 0.80 | 0.81 | 0.74 | 0.81 | 0.41 | 0.39 |
| Marijuana and Hashish | 1.24 | 1.31 | 0.78 | 0.77 | 0.71 | 0.76 | 0.39 | 0.36 |
| Cocaine | 0.62 | 0.72 | 0.25 | 0.25 | 0.15 | 0.12 | 0.06 | 0.05 |
| Crack | 0.18 | 0.08 | 0.13 | 0.05 | 0.07 | 0.01 | 0.02 | 0.01 |
| Heroin | 0.22 | 0.18 | 0.13 | 0.11 | 0.09 | 0.15 | 0.03 | 0.06 |
| Hallucinogens | 0.70 | 0.50 | 0.26 | 0.29 | 0.23 | 0.21 | 0.07 | 0.09 |
| LSD | 0.24 | 0.23 | 0.11 | 0.09 | 0.08 | 0.05 | 0.02 | 0.02 |
| PCP | 0.05 | 0.03 | 0.02 | 0.06 | * | 0.01 | * | * |
| Ecstasy | 0.53 | 0.40 | 0.21 | 0.22 | 0.21 | 0.15 | 0.06 | 0.06 |
| Inhalants | 0.22 | 0.43 | 0.17 | 0.14 | 0.10 | 0.24 | 0.04 | 0.03 |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 0.77 | 0.85 | 0.46 | 0.40 | 0.39 | 0.42 | 0.21 | 0.20 |
| Pain Relievers | 0.68 | 0.71 | 0.41 | 0.36 | 0.35 | 0.34 | 0.19 | 0.16 |
| OxyContin [®] | 0.24 | 0.30 | 0.20 | 0.14 | 0.09 | 0.11 | 0.09 | 0.04 |
| Tranquilizers | 0.52 | 0.51 | 0.25 | 0.22 | 0.22 | 0.24 | 0.08 | 0.12 |
| Stimulants ⁴ | 0.46 | 0.55 | 0.22 | 0.14 | 0.13 | 0.15 | 0.10 | 0.08 |
| Methamphetamine ⁴ | 0.11 | 0.19 | 0.15 | 0.06 | 0.08 | 0.09 | 0.03 | 0.07 |
| Sedatives | 0.15 | 0.13 | 0.06 | 0.05 | 0.04 | 0.09 | 0.06 | 0.03 |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 1.01 | 1.03 | 0.50 | 0.50 | 0.44 | 0.50 | 0.22 | 0.22 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.13P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2011 Versus 2010

| Drug | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|---|------------------------------------|------|--|------|------------------------------------|------|--------|------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 |
| ILLICIT DRUGS² | 0.8813 | | 0.5540 | | 0.1031 | | 0.2888 | |
| Marijuana and Hashish | 0.9206 | | 0.0306 | | 0.1632 | | 0.6158 | |
| Cocaine | 0.7490 | | 0.8681 | | 0.6318 | | 0.7372 | |
| Crack | 0.0182 | | 0.1477 | | 0.1033 | | 0.3824 | |
| Heroin | 0.5761 | | 0.7004 | | 0.1692 | | 0.1099 | |
| Hallucinogens | 0.0141 | | 0.8760 | | 0.8134 | | 0.6630 | |
| LSD | 0.7784 | | 0.9865 | | 0.7662 | | 0.4832 | |
| PCP | 0.5908 | | 0.2817 | | * | | * | |
| Ecstasy | 0.0756 | | 0.9418 | | 0.2467 | | 0.8973 | |
| Inhalants | 0.4857 | | 0.6012 | | 0.4886 | | 0.4856 | |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 0.9721 | | 0.0013 | | 0.9542 | | 0.0582 | |
| Pain Relievers | 0.6716 | | 0.0052 | | 0.3952 | | 0.0146 | |
| OxyContin [®] | 0.8953 | | 0.0392 | | 0.9239 | | 0.0076 | |
| Tranquilizers | 0.9832 | | 0.0730 | | 0.4926 | | 0.1155 | |
| Stimulants ⁴ | 0.6814 | | 0.0158 | | 0.4896 | | 0.1738 | |
| Methamphetamine ⁴ | 0.2136 | | 0.0423 | | 0.7269 | | 0.1818 | |
| Sedatives | 0.6276 | | 0.2963 | | 0.2900 | | 0.1840 | |
| ILLICIT DRUGS OTHER THAN MARIJUANA² | 0.1996 | | 0.0308 | | 0.8664 | | 0.1860 | |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.14C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2010 and 2011

| Drug | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|--|------------------------------------|------|--|------|------------------------------------|------|--------|------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 |
| ILLCIT DRUGS² | 189 | 186 | 222 | 202 | 234 | 215 | 213 | 208 |
| Marijuana and Hashish | 172 | 165 | 181 | 186 | 205 | 187 | 156 | 170 |
| Cocaine | 61 | 69 | 59 | 54 | 40 | 31 | 36 | 25 |
| Crack | 28 | 26 | 31 | 30 | 32 | 3 | 26 | 10 |
| Heroin | 14 | 19 | 5 | 13 | 26 | 9 | 26 | 31 |
| Hallucinogens | 48 | 23 | 29 | 19 | 11 | 19 | 22 | 15 |
| LSD | 3 | 1 | * | * | * | 5 | * | 15 |
| PCP | * | 3 | 3 | 7 | * | * | 20 | * |
| Ecstasy | 37 | 19 | 24 | 14 | * | 18 | 7 | * |
| Inhalants | 22 | 28 | 22 | 21 | 70 | 24 | 13 | 34 |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 93 | 92 | 124 | 99 | 127 | 109 | 139 | 125 |
| Pain Relievers | 84 | 67 | 91 | 88 | 108 | 85 | 106 | 111 |
| OxyContin [®] | 26 | 22 | 27 | 27 | 24 | 17 | 46 | 26 |
| Tranquilizers | 50 | 59 | 71 | 53 | 67 | 55 | 70 | 73 |
| Stimulants ⁴ | 41 | 45 | 39 | 34 | 30 | 25 | 39 | 52 |
| Methamphetamine ⁴ | 20 | 41 | 18 | 16 | 26 | 22 | 30 | 49 |
| Sedatives | 21 | 22 | 54 | 11 | 16 | 20 | 39 | 35 |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 119 | 110 | 139 | 109 | 136 | 116 | 144 | 132 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.14D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2010 and 2011

| Drug | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|--|------------------------------------|------|--|------|------------------------------------|------|--------|------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 |
| ILLCIT DRUGS² | 1.39 | 1.35 | 0.68 | 0.61 | 0.35 | 0.32 | 0.24 | 0.23 |
| Marijuana and Hashish | 1.29 | 1.25 | 0.56 | 0.57 | 0.31 | 0.28 | 0.18 | 0.19 |
| Cocaine | 0.50 | 0.59 | 0.19 | 0.17 | 0.06 | 0.05 | 0.04 | 0.03 |
| Crack | 0.24 | 0.23 | 0.10 | 0.10 | 0.05 | 0.00 | 0.03 | 0.01 |
| Heroin | 0.12 | 0.17 | 0.02 | 0.04 | 0.04 | 0.01 | 0.03 | 0.04 |
| Hallucinogens | 0.40 | 0.20 | 0.09 | 0.06 | 0.02 | 0.03 | 0.02 | 0.02 |
| LSD | 0.02 | 0.01 | * | * | * | 0.01 | * | 0.02 |
| PCP | * | 0.03 | 0.01 | 0.02 | * | * | 0.02 | * |
| Ecstasy | 0.31 | 0.16 | 0.08 | 0.04 | * | 0.03 | 0.01 | * |
| Inhalants | 0.18 | 0.24 | 0.07 | 0.07 | 0.11 | 0.04 | 0.01 | 0.04 |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 0.77 | 0.78 | 0.40 | 0.31 | 0.20 | 0.16 | 0.16 | 0.14 |
| Pain Relievers | 0.70 | 0.58 | 0.29 | 0.28 | 0.17 | 0.13 | 0.12 | 0.12 |
| OxyContin [®] | 0.21 | 0.20 | 0.09 | 0.09 | 0.04 | 0.03 | 0.05 | 0.03 |
| Tranquilizers | 0.42 | 0.51 | 0.23 | 0.17 | 0.10 | 0.08 | 0.08 | 0.08 |
| Stimulants ⁴ | 0.35 | 0.39 | 0.13 | 0.11 | 0.05 | 0.04 | 0.04 | 0.06 |
| Methamphetamine ⁴ | 0.17 | 0.35 | 0.06 | 0.05 | 0.04 | 0.03 | 0.03 | 0.06 |
| Sedatives | 0.18 | 0.20 | 0.17 | 0.04 | 0.03 | 0.03 | 0.04 | 0.04 |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 0.95 | 0.90 | 0.44 | 0.35 | 0.21 | 0.17 | 0.16 | 0.15 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.14P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2011 Versus 2010

| Drug | LEVEL OF ALCOHOL USE IN PAST MONTH | | | | | | | |
|---|------------------------------------|------|--|------|------------------------------------|------|--------|------|
| | Heavy Use ¹ | | Binge Use But Not Heavy Use ¹ | | Use But Not Binge Use ¹ | | No Use | |
| | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 | 2010 | 2011 |
| ILLICIT DRUGS² | 0.9209 | | 0.3950 | | 0.4490 | | 0.5395 | |
| Marijuana and Hashish | 0.5461 | | 0.6502 | | 0.8996 | | 0.2166 | |
| Cocaine | 0.5440 | | 0.6802 | | 0.5304 | | 0.2374 | |
| Crack | 0.9353 | | 0.9579 | | 0.0710 | | 0.1023 | |
| Heroin | 0.5684 | | 0.0833 | | 0.0906 | | 0.5024 | |
| Hallucinogens | 0.0612 | | 0.4177 | | 0.1992 | | 0.4542 | |
| LSD | 0.4920 | | * | | * | | * | |
| PCP | * | | 0.5994 | | * | | * | |
| Ecstasy | 0.1752 | | 0.6011 | | * | | * | |
| Inhalants | 0.4668 | | 0.7218 | | 0.1935 | | 0.2101 | |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 0.5992 | | 0.0941 | | 0.1232 | | 0.7886 | |
| Pain Relievers | 0.1099 | | 0.3719 | | 0.2432 | | 0.3173 | |
| OxyContin [®] | 0.8254 | | 0.8717 | | 0.1488 | | 0.8369 | |
| Tranquilizers | 0.4615 | | 0.1793 | | 0.0683 | | 0.2640 | |
| Stimulants ⁴ | 0.9040 | | 0.5205 | | 0.8356 | | 0.7890 | |
| Methamphetamine ⁴ | 0.1210 | | 0.3508 | | 0.8791 | | 0.5313 | |
| Sedatives | 0.6774 | | 0.1224 | | 0.8272 | | 0.6298 | |
| ILLICIT DRUGS OTHER THAN MARIJUANA² | 0.6963 | | 0.0671 | | 0.1255 | | 0.8384 | |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.15C Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Numbers in Thousands, 2010 and 2011

| Drug | Total (2010) | Total (2011) | Aged 12-20 (2010) | Aged 12-20 (2011) | Aged 21+ (2010) | Aged 21+ (2011) |
|--|-------------------------|-------------------------|------------------------------|------------------------------|----------------------------|----------------------------|
| ILLCIT DRUGS¹ | 442 | 412 | 115 | 136 | 421 | 388 |
| Marijuana or Hashish | 377 | 367 | 108 | 127 | 356 | 347 |
| Cocaine (Including Crack) | 104 | 103 | 26 | 30 | 101 | 98 |
| Heroin | 34 | 35 | 12 | 14 | 32 | 31 |
| Hallucinogens | 79 | 56 | 35 | 31 | 68 | 45 |
| Inhalants | 81 | 57 | 17 | 28 | 78 | 45 |
| Pain Relievers ² | 178 | 148 | 48 | 40 | 172 | 142 |
| Tranquilizers ² | 119 | 109 | 25 | 29 | 115 | 105 |
| Stimulants ^{2,3} | 75 | 70 | 24 | 20 | 69 | 66 |
| Methamphetamine ³ | 42 | 50 | 15 | 10 | 39 | 50 |
| Sedatives ² | 61 | 34 | 8 | 7 | 60 | 33 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 259 | 211 | 65 | 66 | 251 | 197 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.15D Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Standard Errors of Percentages, 2010 and 2011

| Drug | Total (2010) | Total (2011) | Aged 12-20 (2010) | Aged 12-20 (2011) | Aged 21+ (2010) | Aged 21+ (2011) |
|--|-------------------------|-------------------------|------------------------------|------------------------------|----------------------------|----------------------------|
| ILLCIT DRUGS¹ | 0.31 | 0.29 | 0.83 | 0.90 | 0.32 | 0.29 |
| Marijuana or Hashish | 0.27 | 0.26 | 0.82 | 0.87 | 0.28 | 0.26 |
| Cocaine (Including Crack) | 0.08 | 0.08 | 0.26 | 0.29 | 0.08 | 0.08 |
| Heroin | 0.03 | 0.03 | 0.12 | 0.14 | 0.03 | 0.03 |
| Hallucinogens | 0.06 | 0.04 | 0.34 | 0.32 | 0.06 | 0.04 |
| Inhalants | 0.06 | 0.04 | 0.17 | 0.29 | 0.06 | 0.04 |
| Pain Relievers ² | 0.14 | 0.11 | 0.45 | 0.40 | 0.14 | 0.12 |
| Tranquilizers ² | 0.09 | 0.08 | 0.24 | 0.30 | 0.09 | 0.08 |
| Stimulants ^{2,3} | 0.06 | 0.05 | 0.23 | 0.20 | 0.06 | 0.05 |
| Methamphetamine ³ | 0.03 | 0.04 | 0.15 | 0.10 | 0.03 | 0.04 |
| Sedatives ² | 0.05 | 0.03 | 0.08 | 0.08 | 0.05 | 0.03 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 0.19 | 0.16 | 0.59 | 0.60 | 0.20 | 0.16 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.15P Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: P Values from Tests of Differences of Percentages, 2011 Versus 2010

| Drug | Total (2010) | Total (2011) | Aged 12-20 (2010) | Aged 12-20 (2011) | Aged 21+ (2010) | Aged 21+ (2011) |
|--|-------------------------|-------------------------|------------------------------|------------------------------|----------------------------|----------------------------|
| ILLCIT DRUGS¹ | 0.2036 | | 0.0932 | | 0.1400 | |
| Marijuana or Hashish | 0.9677 | | 0.0085 | | 0.7067 | |
| Cocaine (Including Crack) | 0.6267 | | 0.1393 | | 0.3944 | |
| Heroin | 0.9715 | | 0.9747 | | 0.9947 | |
| Hallucinogens | 0.0131 | | 0.1812 | | 0.0427 | |
| Inhalants | 0.3512 | | 0.6055 | | 0.2574 | |
| Pain Relievers ² | 0.0013 | | 0.0934 | | 0.0053 | |
| Tranquilizers ² | 0.1159 | | 0.0433 | | 0.0366 | |
| Stimulants ^{2,3} | 0.2832 | | 0.4316 | | 0.3958 | |
| Methamphetamine ³ | 0.6725 | | 0.6553 | | 0.5525 | |
| Sedatives ² | 0.1451 | | 0.7631 | | 0.1324 | |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 0.0013 | | 0.4616 | | 0.0023 | |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.16C Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: Standard Errors of Numbers in Thousands, 2010 and 2011

| Drug¹ | Total (2010) | Total (2011) | Aged 12-20 (2010) | Aged 12-20 (2011) | Aged 21+ (2010) | Aged 21+ (2011) |
|--|-------------------------|-------------------------|------------------------------|------------------------------|----------------------------|----------------------------|
| ILLCIT DRUGS² | 268 | 261 | 71 | 82 | 255 | 243 |
| Marijuana or Hashish | 246 | 246 | 69 | 76 | 234 | 230 |
| Cocaine (Including Crack) | 49 | 48 | 11 | 14 | 47 | 46 |
| Heroin | 18 | 15 | 8 | 7 | 17 | 13 |
| Hallucinogens | 26 | 22 | 10 | 8 | 24 | 20 |
| Inhalants | 14 | 22 | 8 | 16 | 12 | 15 |
| Pain Relievers ³ | 64 | 54 | 14 | 14 | 63 | 54 |
| Tranquilizers ³ | 32 | 19 | 10 | 8 | 31 | 17 |
| Stimulants ³ | 27 | 24 | 4 | 2 | 27 | 24 |
| Methamphetamine ⁴ | 26 | 31 | 2 | 4 | 26 | 31 |
| Sedatives ³ | 11 | 21 | 3 | 3 | 10 | 21 |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 97 | 89 | 24 | 30 | 95 | 85 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.16D Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: Standard Errors of Percentages, 2010 and 2011

| Drug¹ | Total (2010) | Total (2011) | Aged 12-20 (2010) | Aged 12-20 (2011) | Aged 21+ (2010) | Aged 21+ (2011) |
|--|-------------------------|-------------------------|------------------------------|------------------------------|----------------------------|----------------------------|
| ILLCIT DRUGS² | 0.20 | 0.20 | 0.62 | 0.70 | 0.20 | 0.20 |
| Marijuana or Hashish | 0.18 | 0.18 | 0.60 | 0.67 | 0.19 | 0.19 |
| Cocaine (Including Crack) | 0.04 | 0.04 | 0.11 | 0.14 | 0.04 | 0.04 |
| Heroin | 0.01 | 0.01 | 0.08 | 0.07 | 0.01 | 0.01 |
| Hallucinogens | 0.02 | 0.02 | 0.10 | 0.08 | 0.02 | 0.02 |
| Inhalants | 0.01 | 0.02 | 0.08 | 0.16 | 0.01 | 0.01 |
| Pain Relievers ³ | 0.05 | 0.04 | 0.14 | 0.14 | 0.05 | 0.04 |
| Tranquilizers ³ | 0.02 | 0.01 | 0.10 | 0.08 | 0.03 | 0.01 |
| Stimulants ³ | 0.02 | 0.02 | 0.04 | 0.02 | 0.02 | 0.02 |
| Methamphetamine ⁴ | 0.02 | 0.02 | 0.02 | 0.04 | 0.02 | 0.02 |
| Sedatives ³ | 0.01 | 0.02 | 0.03 | 0.03 | 0.01 | 0.02 |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 0.07 | 0.07 | 0.24 | 0.30 | 0.08 | 0.07 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.16P Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: P Values from Tests of Differences of Percentages, 2011 Versus 2010

| Drug¹ | Total (2010) | Total (2011) | Aged 12-20 (2010) | Aged 12-20 (2011) | Aged 21+ (2010) | Aged 21+ (2011) |
|---|-------------------------|-------------------------|------------------------------|------------------------------|----------------------------|----------------------------|
| ILLICIT DRUGS² | 0.4764 | | 0.2400 | | 0.3804 | |
| Marijuana or Hashish | 0.8731 | | 0.1991 | | 0.7579 | |
| Cocaine (Including Crack) | 0.6628 | | 0.7599 | | 0.6071 | |
| Heroin | 0.7361 | | 0.9074 | | 0.7624 | |
| Hallucinogens | 0.9573 | | 0.4490 | | 0.7777 | |
| Inhalants | 0.9770 | | 0.8987 | | 0.9012 | |
| Pain Relievers ³ | 0.1216 | | 0.3291 | | 0.1879 | |
| Tranquilizers ³ | 0.0199 | | 0.5100 | | 0.0272 | |
| Stimulants ³ | 0.2951 | | 0.0475 | | 0.4273 | |
| Methamphetamine ⁴ | 0.8855 | | 0.8946 | | 0.8714 | |
| Sedatives ³ | 0.4092 | | 0.9789 | | 0.4015 | |
| ILLICIT DRUGS OTHER THAN MARIJUANA² | 0.1992 | | 0.9383 | | 0.1940 | |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.17C Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2010 and 2011

| Drug | Total (2010) | Total (2011) | Aged 12-17 (2010) | Aged 12-17 (2011) | Aged 18-25 (2010) | Aged 18-25 (2011) | Aged 26+ (2010) | Aged 26+ (2011) |
|--|-------------------------|-------------------------|------------------------------|------------------------------|------------------------------|------------------------------|----------------------------|----------------------------|
| ILLCIT DRUGS¹ | 442 | 412 | 57 | 50 | 161 | 179 | 395 | 365 |
| Marijuana or Hashish | 377 | 367 | 54 | 48 | 149 | 165 | 333 | 331 |
| Cocaine (Including Crack) | 104 | 103 | 11 | 11 | 39 | 42 | 94 | 93 |
| Heroin | 34 | 35 | 2 | 7 | 16 | 18 | 30 | 28 |
| Hallucinogens | 79 | 56 | 18 | 19 | 48 | 39 | 57 | 35 |
| Inhalants | 81 | 57 | 14 | 11 | 20 | 36 | 76 | 43 |
| Pain Relievers ² | 178 | 148 | 25 | 21 | 62 | 56 | 163 | 136 |
| Tranquilizers ² | 119 | 109 | 11 | 15 | 38 | 36 | 110 | 100 |
| Stimulants ^{2,3} | 75 | 70 | 13 | 10 | 34 | 32 | 65 | 61 |
| Methamphetamine ³ | 42 | 50 | 7 | 7 | 16 | 12 | 38 | 49 |
| Sedatives ² | 61 | 34 | 5 | 5 | 9 | 10 | 60 | 32 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 259 | 211 | 33 | 30 | 89 | 90 | 237 | 188 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.17D Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2010 and 2011

| Drug | Total (2010) | Total (2011) | Aged 12-17 (2010) | Aged 12-17 (2011) | Aged 18-25 (2010) | Aged 18-25 (2011) | Aged 26+ (2010) | Aged 26+ (2011) |
|--|-------------------------|-------------------------|------------------------------|------------------------------|------------------------------|------------------------------|----------------------------|----------------------------|
| ILLCIT DRUGS¹ | 0.31 | 0.29 | 1.20 | 1.19 | 0.57 | 0.60 | 0.35 | 0.32 |
| Marijuana or Hashish | 0.27 | 0.26 | 1.17 | 1.16 | 0.55 | 0.57 | 0.30 | 0.29 |
| Cocaine (Including Crack) | 0.08 | 0.08 | 0.33 | 0.32 | 0.18 | 0.19 | 0.09 | 0.08 |
| Heroin | 0.03 | 0.03 | 0.07 | 0.22 | 0.08 | 0.09 | 0.03 | 0.03 |
| Hallucinogens | 0.06 | 0.04 | 0.53 | 0.54 | 0.22 | 0.18 | 0.05 | 0.03 |
| Inhalants | 0.06 | 0.04 | 0.42 | 0.33 | 0.10 | 0.17 | 0.07 | 0.04 |
| Pain Relievers ² | 0.14 | 0.11 | 0.71 | 0.61 | 0.28 | 0.25 | 0.15 | 0.13 |
| Tranquilizers ² | 0.09 | 0.08 | 0.33 | 0.46 | 0.18 | 0.17 | 0.10 | 0.09 |
| Stimulants ^{2,3} | 0.06 | 0.05 | 0.38 | 0.30 | 0.16 | 0.15 | 0.06 | 0.06 |
| Methamphetamine ³ | 0.03 | 0.04 | 0.22 | 0.22 | 0.08 | 0.06 | 0.04 | 0.04 |
| Sedatives ² | 0.05 | 0.03 | 0.16 | 0.15 | 0.04 | 0.05 | 0.06 | 0.03 |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 0.19 | 0.16 | 0.91 | 0.85 | 0.37 | 0.38 | 0.21 | 0.17 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.17P Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2011 Versus 2010

| Drug | Total (2010) | Total (2011) | Aged 12-17 (2010) | Aged 12-17 (2011) | Aged 18-25 (2010) | Aged 18-25 (2011) | Aged 26+ (2010) | Aged 26+ (2011) |
|--|-------------------------|-------------------------|------------------------------|------------------------------|------------------------------|------------------------------|----------------------------|----------------------------|
| ILLCIT DRUGS¹ | 0.2036 | | 0.6378 | | 0.5840 | | 0.1901 | |
| Marijuana or Hashish | 0.9677 | | 0.6257 | | 0.1108 | | 0.7075 | |
| Cocaine (Including Crack) | 0.6267 | | 0.8245 | | 0.5389 | | 0.7948 | |
| Heroin | 0.9715 | | 0.3583 | | 0.8825 | | 0.7926 | |
| Hallucinogens | 0.0131 | | 0.7660 | | 0.0463 | | 0.1214 | |
| Inhalants | 0.3512 | | 0.3250 | | 0.7166 | | 0.3302 | |
| Pain Relievers ² | 0.0013 | | 0.3391 | | 0.0069 | | 0.0234 | |
| Tranquilizers ² | 0.1159 | | 0.0169 | | 0.3057 | | 0.1110 | |
| Stimulants ^{2,3} | 0.2832 | | 0.8618 | | 0.3035 | | 0.5189 | |
| Methamphetamine ³ | 0.6725 | | 0.3273 | | 0.3935 | | 0.5588 | |
| Sedatives ² | 0.1451 | | 0.9472 | | 0.6652 | | 0.1652 | |
| ILLCIT DRUGS OTHER THAN MARIJUANA¹ | 0.0013 | | 0.3737 | | 0.0159 | | 0.0153 | |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.18C Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2010 and 2011

| Drug¹ | Total (2010) | Total (2011) | Aged 12-17 (2010) | Aged 12-17 (2011) | Aged 18-25 (2010) | Aged 18-25 (2011) | Aged 26+ (2010) | Aged 26+ (2011) |
|--|-------------------------|-------------------------|------------------------------|------------------------------|------------------------------|------------------------------|----------------------------|----------------------------|
| ILLCIT DRUGS² | 268 | 261 | 38 | 32 | 103 | 118 | 237 | 230 |
| Marijuana or Hashish | 246 | 246 | 38 | 31 | 99 | 111 | 218 | 217 |
| Cocaine (Including Crack) | 49 | 48 | 4 | 5 | 20 | 18 | 43 | 44 |
| Heroin | 18 | 15 | * | 4 | 10 | 10 | 16 | 10 |
| Hallucinogens | 26 | 22 | 5 | 4 | 13 | 15 | 22 | 15 |
| Inhalants | 14 | 22 | 7 | 1 | 10 | 16 | 8 | 15 |
| Pain Relievers ³ | 64 | 54 | 8 | 8 | 21 | 19 | 60 | 52 |
| Tranquilizers ³ | 32 | 19 | 4 | 4 | 12 | 9 | 30 | 16 |
| Stimulants ³ | 27 | 24 | 1 | 1 | 15 | 7 | 23 | 23 |
| Methamphetamine ⁴ | 26 | 31 | 2 | 4 | 6 | 3 | 26 | 31 |
| Sedatives ³ | 11 | 21 | 3 | 3 | 2 | * | 10 | 21 |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 97 | 89 | 12 | 12 | 36 | 38 | 89 | 82 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.18D Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2010 and 2011

| Drug¹ | Total (2010) | Total (2011) | Aged 12-17 (2010) | Aged 12-17 (2011) | Aged 18-25 (2010) | Aged 18-25 (2011) | Aged 26+ (2010) | Aged 26+ (2011) |
|--|-------------------------|-------------------------|------------------------------|------------------------------|------------------------------|------------------------------|----------------------------|----------------------------|
| ILLCIT DRUGS² | 0.20 | 0.20 | 0.99 | 0.90 | 0.43 | 0.47 | 0.22 | 0.21 |
| Marijuana or Hashish | 0.18 | 0.18 | 0.98 | 0.88 | 0.41 | 0.46 | 0.20 | 0.20 |
| Cocaine (Including Crack) | 0.04 | 0.04 | 0.13 | 0.15 | 0.10 | 0.09 | 0.04 | 0.04 |
| Heroin | 0.01 | 0.01 | * | 0.12 | 0.05 | 0.05 | 0.01 | 0.01 |
| Hallucinogens | 0.02 | 0.02 | 0.14 | 0.11 | 0.06 | 0.07 | 0.02 | 0.01 |
| Inhalants | 0.01 | 0.02 | 0.21 | 0.04 | 0.05 | 0.08 | 0.01 | 0.01 |
| Pain Relievers ³ | 0.05 | 0.04 | 0.23 | 0.25 | 0.10 | 0.09 | 0.06 | 0.05 |
| Tranquilizers ³ | 0.02 | 0.01 | 0.12 | 0.13 | 0.06 | 0.04 | 0.03 | 0.01 |
| Stimulants ³ | 0.02 | 0.02 | 0.04 | 0.04 | 0.07 | 0.03 | 0.02 | 0.02 |
| Methamphetamine ⁴ | 0.02 | 0.02 | 0.05 | 0.13 | 0.03 | 0.01 | 0.02 | 0.03 |
| Sedatives ³ | 0.01 | 0.02 | 0.09 | 0.09 | 0.01 | * | 0.01 | 0.02 |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 0.07 | 0.07 | 0.35 | 0.37 | 0.17 | 0.18 | 0.08 | 0.07 |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.18P Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2011 Versus 2010

| Drug¹ | Total (2010) | Total (2011) | Aged 12-17 (2010) | Aged 12-17 (2011) | Aged 18-25 (2010) | Aged 18-25 (2011) | Aged 26+ (2010) | Aged 26+ (2011) |
|---|-------------------------|-------------------------|------------------------------|------------------------------|------------------------------|------------------------------|----------------------------|----------------------------|
| ILLICIT DRUGS² | 0.4764 | | 0.8738 | | 0.9449 | | 0.5313 | |
| Marijuana or Hashish | 0.8731 | | 0.6584 | | 0.6753 | | 0.9294 | |
| Cocaine (Including Crack) | 0.6628 | | 0.6174 | | 0.4018 | | 0.8851 | |
| Heroin | 0.7361 | | * | | 0.8429 | | 0.6128 | |
| Hallucinogens | 0.9573 | | 0.6095 | | 0.7235 | | 0.8588 | |
| Inhalants | 0.9770 | | 0.1201 | | 0.9517 | | 0.5797 | |
| Pain Relievers ³ | 0.1216 | | 0.9212 | | 0.4920 | | 0.1811 | |
| Tranquilizers ³ | 0.0199 | | 0.8630 | | 0.2782 | | 0.0351 | |
| Stimulants ³ | 0.2951 | | 0.7787 | | 0.0961 | | 0.7625 | |
| Methamphetamine ⁴ | 0.8855 | | 0.4666 | | 0.1885 | | 0.9766 | |
| Sedatives ³ | 0.4092 | | 0.9949 | | * | | 0.3816 | |
| ILLICIT DRUGS OTHER THAN MARIJUANA² | 0.1992 | | 0.7963 | | 0.8980 | | 0.1830 | |

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.