

**Table 6.19C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2010 and 2011**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2010	2011	2010	2011	2010	2011	2010	2011
<b>ILLCIT DRUGS<sup>1</sup></b>	303	289	168	147	292	285	146	154
Marijuana and Hashish	273	268	143	126	241	247	94	119
Cocaine	95	82	30	26	40	60	22	5
Crack	53	41	22	10	16	*	14	*
Heroin	34	30	17	31	*	8	20	8
Hallucinogens	71	49	13	18	31	26	23	14
LSD	17	15	4	15	9	8	*	4
PCP	4	10	1	*	*	*	20	*
Ecstasy	54	40	10	7	26	21	8	7
Inhalants	39	44	5	8	71	36	23	37
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	151	129	87	83	160	131	110	104
Pain Relievers	131	103	74	73	123	109	84	89
OxyContin <sup>®</sup>	43	37	48	19	22	25	5	19
Tranquilizers	88	78	51	44	77	73	50	60
Stimulants <sup>3</sup>	61	60	26	29	40	34	32	44
Methamphetamine <sup>3</sup>	33	47	20	24	26	18	23	44
Sedatives	32	30	14	17	52	16	37	31
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	185	151	92	87	171	146	114	110

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.19D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2010 and 2011**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2010	2011	2010	2011	2010	2011	2010	2011
<b>ILLICIT DRUGS<sup>1</sup></b>	0.66	0.68	0.81	0.74	0.31	0.28	0.14	0.15
Marijuana and Hashish	0.62	0.64	0.70	0.65	0.26	0.25	0.09	0.11
Cocaine	0.24	0.21	0.15	0.14	0.04	0.06	0.02	0.01
Crack	0.13	0.11	0.11	0.05	0.02	*	0.01	*
Heroin	0.09	0.08	0.08	0.16	*	0.01	0.02	0.01
Hallucinogens	0.18	0.13	0.06	0.09	0.03	0.03	0.02	0.01
LSD	0.04	0.04	0.02	0.08	0.01	0.01	*	0.00
PCP	0.01	0.03	0.00	*	*	*	0.02	*
Ecstasy	0.14	0.11	0.05	0.04	0.03	0.02	0.01	0.01
Inhalants	0.10	0.12	0.03	0.04	0.08	0.04	0.02	0.04
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.37	0.33	0.43	0.42	0.17	0.14	0.11	0.10
Pain Relievers	0.33	0.27	0.37	0.37	0.13	0.11	0.08	0.09
OxyContin <sup>®</sup>	0.11	0.10	0.24	0.10	0.02	0.03	0.00	0.02
Tranquilizers	0.22	0.20	0.25	0.23	0.08	0.08	0.05	0.06
Stimulants <sup>3</sup>	0.16	0.16	0.13	0.15	0.04	0.04	0.03	0.04
Methamphetamine <sup>3</sup>	0.09	0.12	0.10	0.13	0.03	0.02	0.02	0.04
Sedatives	0.08	0.08	0.07	0.09	0.06	0.02	0.04	0.03
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.44	0.39	0.46	0.44	0.18	0.15	0.11	0.10

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.19P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2010	2011	2010	2011	2010	2011	2010	2011
<b>ILLCIT DRUGS<sup>1</sup></b>	0.4385		0.9298		0.7263		0.1936	
Marijuana and Hashish	0.8575		0.9674		0.5560		0.0020	
Cocaine	0.4812		0.9529		0.4774		0.0451	
Crack	0.2814		0.2712		*		*	
Heroin	0.9247		0.1210		*		0.5717	
Hallucinogens	0.0424		0.8203		0.2994		0.7504	
LSD	0.6527		0.4928		0.7291		*	
PCP	0.1108		*		*		*	
Ecstasy	0.1640		0.3638		0.3780		0.5274	
Inhalants	0.7863		0.4816		0.2038		0.8603	
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.0705		0.5350		0.0266		0.5049	
Pain Relievers	0.0111		0.2494		0.0945		0.7102	
OxyContin <sup>®</sup>	0.0885		0.1416		0.5844		0.1371	
Tranquilizers	0.1464		0.4066		0.5998		0.6620	
Stimulants <sup>3</sup>	0.9798		0.7940		0.0747		0.9744	
Methamphetamine <sup>3</sup>	0.2052		0.8369		0.2112		0.3448	
Sedatives	0.4342		0.3522		0.2399		0.1690	
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.0220		0.5600		0.0467		0.5501	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.20C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2010 and 2011**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2010	2011	2010	2011	2010	2011	2010	2011
<b>ILLCIT DRUGS<sup>1</sup></b>	44	37	22	26	35	33	42	42
Marijuana and Hashish	41	36	18	25	32	31	29	30
Cocaine	11	10	2	4	2	5	2	1
Crack	2	2	*	*	0	*	*	*
Heroin	2	7	*	1	1	0	2	*
Hallucinogens	17	16	5	5	8	10	9	11
LSD	9	7	3	1	3	3	2	3
PCP	2	2	0	*	1	0	2	*
Ecstasy	13	11	3	2	7	9	6	6
Inhalants	11	9	4	4	9	6	18	16
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	22	21	12	11	16	11	25	22
Pain Relievers	20	18	10	10	15	10	22	21
OxyContin <sup>®</sup>	8	8	4	3	2	3	4	2
Tranquilizers	11	15	8	4	5	5	7	6
Stimulants <sup>3</sup>	11	10	2	4	6	3	7	5
Methamphetamine <sup>3</sup>	6	7	2	4	5	2	2	1
Sedatives	5	3	2	2	2	4	5	2
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	28	26	13	12	19	17	31	29

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.20D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2010 and 2011**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2010	2011	2010	2011	2010	2011	2010	2011
<b>ILLICIT DRUGS<sup>1</sup></b>	2.02	1.83	2.19	2.43	1.37	1.33	0.20	0.19
Marijuana and Hashish	2.03	1.88	1.95	2.44	1.29	1.28	0.14	0.14
Cocaine	0.85	0.80	0.29	0.56	0.11	0.21	0.01	0.00
Crack	0.15	0.18	*	*	0.01	*	*	*
Heroin	0.17	0.63	*	0.12	0.06	0.01	0.01	*
Hallucinogens	1.31	1.25	0.55	0.59	0.36	0.46	0.04	0.05
LSD	0.70	0.59	0.31	0.07	0.14	0.13	0.01	0.01
PCP	0.12	0.15	0.05	*	0.05	0.02	0.01	*
Ecstasy	1.03	0.89	0.38	0.23	0.31	0.42	0.03	0.03
Inhalants	0.89	0.78	0.50	0.52	0.42	0.29	0.09	0.08
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	1.53	1.63	1.35	1.39	0.74	0.51	0.12	0.10
Pain Relievers	1.44	1.49	1.22	1.19	0.69	0.46	0.11	0.10
OxyContin <sup>®</sup>	0.66	0.71	0.53	0.38	0.10	0.13	0.02	0.01
Tranquilizers	0.86	1.21	0.92	0.49	0.25	0.22	0.03	0.03
Stimulants <sup>3</sup>	0.88	0.81	0.23	0.56	0.30	0.14	0.03	0.02
Methamphetamine <sup>3</sup>	0.46	0.61	0.20	0.51	0.23	0.08	0.01	0.00
Sedatives	0.40	0.26	0.28	0.23	0.09	0.19	0.03	0.01
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	1.88	1.86	1.49	1.48	0.86	0.75	0.15	0.13

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.20P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2011 Versus 2010**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2010	2011	2010	2011	2010	2011	2010	2011
<b>ILLCIT DRUGS<sup>1</sup></b>	0.2126		0.0395		0.2640		0.7842	
Marijuana and Hashish	0.1286		0.0024		0.9665		0.1338	
Cocaine	0.9242		0.3046		0.4284		0.4368	
Crack	0.9933		*		*		*	
Heroin	0.2388		*		0.1802		*	
Hallucinogens	0.6788		0.9708		0.6441		0.5820	
LSD	0.5363		0.3380		0.6055		0.9882	
PCP	0.8776		*		0.6205		*	
Ecstasy	0.3828		0.2902		0.4209		0.6498	
Inhalants	0.6796		0.7066		0.0402		0.3425	
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.1621		0.7125		0.0724		0.3888	
Pain Relievers	0.6223		0.5989		0.0417		0.6715	
OxyContin <sup>®</sup>	0.5944		0.9671		0.3269		0.3240	
Tranquilizers	0.0058		0.2608		0.9278		0.4471	
Stimulants <sup>3</sup>	0.6872		0.2489		0.3737		0.1462	
Methamphetamine <sup>3</sup>	0.1335		0.2838		0.6093		0.1598	
Sedatives	0.3368		0.8843		0.2344		0.0865	
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.4297		0.6983		0.0425		0.4197	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.21C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2010 and 2011**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2010	2011	2010	2011	2010	2011	2010	2011
<b>ILLCIT DRUGS<sup>1</sup></b>	125	125	45	37	85	101	38	43
Marijuana and Hashish	117	119	42	34	78	90	33	37
Cocaine	36	38	7	7	15	17	2	3
Crack	16	6	2	1	2	2	*	*
Heroin	16	18	3	5	0	3	1	6
Hallucinogens	42	33	8	9	20	20	5	8
LSD	14	12	3	2	8	5	1	*
PCP	3	6	*	0	*	*	*	*
Ecstasy	32	26	7	7	16	16	5	4
Inhalants	17	35	3	4	11	6	5	2
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	58	58	23	20	37	33	19	19
Pain Relievers	51	48	20	16	31	27	16	16
OxyContin <sup>®</sup>	21	18	11	4	5	9	3	5
Tranquilizers	35	31	9	12	16	18	4	10
Stimulants <sup>3</sup>	28	29	11	10	16	11	7	4
Methamphetamine <sup>3</sup>	16	11	4	9	4	4	1	2
Sedatives	8	8	3	4	5	5	7	2
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	73	76	24	22	43	41	19	21

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.21D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2010 and 2011**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2010	2011	2010	2011	2010	2011	2010	2011
<b>ILLICIT DRUGS<sup>1</sup></b>	0.90	0.92	1.47	1.29	0.62	0.66	0.34	0.37
Marijuana and Hashish	0.90	0.91	1.42	1.21	0.58	0.60	0.30	0.33
Cocaine	0.38	0.41	0.27	0.25	0.13	0.14	0.02	0.03
Crack	0.17	0.06	0.09	0.03	0.02	0.01	*	*
Heroin	0.18	0.20	0.14	0.19	0.00	0.03	0.01	0.06
Hallucinogens	0.43	0.36	0.32	0.34	0.17	0.17	0.05	0.07
LSD	0.15	0.14	0.10	0.09	0.07	0.04	0.01	*
PCP	0.04	0.07	*	0.01	*	*	*	*
Ecstasy	0.34	0.29	0.27	0.26	0.14	0.13	0.04	0.04
Inhalants	0.18	0.39	0.11	0.17	0.10	0.05	0.04	0.01
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.57	0.56	0.85	0.75	0.29	0.27	0.17	0.17
Pain Relievers	0.53	0.48	0.74	0.59	0.25	0.22	0.15	0.14
OxyContin <sup>®</sup>	0.23	0.20	0.43	0.14	0.04	0.07	0.03	0.04
Tranquilizers	0.37	0.33	0.37	0.46	0.13	0.15	0.04	0.09
Stimulants <sup>3</sup>	0.30	0.31	0.42	0.37	0.14	0.09	0.07	0.04
Methamphetamine <sup>3</sup>	0.17	0.12	0.16	0.33	0.03	0.03	0.01	0.02
Sedatives	0.09	0.09	0.11	0.14	0.04	0.04	0.07	0.02
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.67	0.73	0.88	0.83	0.34	0.32	0.18	0.19

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.



**Table 6.21P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2011 Versus 2010**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2010	2011	2010	2011	2010	2011	2010	2011
<b>ILLCIT DRUGS<sup>1</sup></b>	0.7104		0.0529		0.3166		0.6934	
Marijuana and Hashish	0.1990		0.0908		0.0965		0.3179	
Cocaine	0.8659		0.8724		0.4681		0.6306	
Crack	0.0058		0.3836		0.3509		*	
Heroin	0.9149		0.1487		0.3600		0.4107	
Hallucinogens	0.0379		0.8203		0.7910		0.6617	
LSD	0.8958		0.5518		0.5046		*	
PCP	0.3349		*		*		*	
Ecstasy	0.0880		0.9108		0.6033		0.6939	
Inhalants	0.4281		0.4491		0.2856		0.0714	
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.0566		0.1332		0.2187		0.2924	
Pain Relievers	0.0158		0.0332		0.2660		0.2245	
OxyContin <sup>®</sup>	0.0720		0.0043		0.5144		0.7696	
Tranquilizers	0.1736		0.2690		0.5282		0.2378	
Stimulants <sup>3</sup>	0.9323		0.4169		0.0200		0.1751	
Methamphetamine <sup>3</sup>	0.4343		0.2098		0.8702		0.5718	
Sedatives	0.9690		0.8092		0.4723		0.0883	
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.0870		0.2809		0.1257		0.4687	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.22C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2010 and 2011**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2010	2011	2010	2011	2010	2011	2010	2011
<b>ILLCIT DRUGS<sup>1</sup></b>	269	249	160	140	273	261	135	140
Marijuana and Hashish	239	232	135	119	223	227	82	108
Cocaine	86	73	30	25	37	57	21	4
Crack	50	40	21	10	16	*	14	*
Heroin	30	23	16	31	*	7	20	5
Hallucinogens	52	33	8	15	23	13	20	*
LSD	3	2	*	15	*	5	*	*
PCP	*	8	*	*	*	*	20	*
Ecstasy	40	28	7	*	19	10	*	*
Inhalants	32	25	*	5	69	35	13	33
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	134	111	82	79	155	126	106	98
Pain Relievers	115	88	69	70	118	105	80	84
OxyContin <sup>®</sup>	36	32	46	19	22	23	*	18
Tranquilizers	78	69	49	43	74	70	49	59
Stimulants <sup>3</sup>	52	52	24	27	37	32	30	44
Methamphetamine <sup>3</sup>	28	45	20	23	25	17	23	44
Sedatives	30	29	13	16	52	14	36	31
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	163	124	87	83	164	138	108	102

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.22D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2010 and 2011**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2010	2011	2010	2011	2010	2011	2010	2011
<b>ILLICIT DRUGS<sup>1</sup></b>	0.82	0.82	0.93	0.85	0.34	0.31	0.19	0.19
Marijuana and Hashish	0.75	0.77	0.79	0.74	0.28	0.27	0.11	0.15
Cocaine	0.30	0.26	0.18	0.16	0.05	0.07	0.03	0.01
Crack	0.18	0.14	0.13	0.07	0.02	*	0.02	*
Heroin	0.11	0.08	0.10	0.20	*	0.01	0.03	0.01
Hallucinogens	0.19	0.12	0.05	0.10	0.03	0.02	0.03	*
LSD	0.01	0.01	*	0.09	*	0.01	*	*
PCP	*	0.03	*	*	*	*	0.03	*
Ecstasy	0.14	0.10	0.04	*	0.02	0.01	*	*
Inhalants	0.11	0.09	*	0.03	0.09	0.04	0.02	0.05
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.45	0.40	0.49	0.49	0.19	0.15	0.15	0.13
Pain Relievers	0.39	0.32	0.42	0.44	0.15	0.13	0.11	0.12
OxyContin <sup>®</sup>	0.13	0.11	0.28	0.12	0.03	0.03	*	0.02
Tranquilizers	0.27	0.25	0.30	0.27	0.09	0.09	0.07	0.08
Stimulants <sup>3</sup>	0.19	0.19	0.15	0.17	0.05	0.04	0.04	0.06
Methamphetamine <sup>3</sup>	0.10	0.16	0.12	0.14	0.03	0.02	0.03	0.06
Sedatives	0.11	0.10	0.08	0.10	0.07	0.02	0.05	0.04
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.54	0.44	0.52	0.51	0.21	0.17	0.15	0.14

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.22P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2010	2011	2010	2011	2010	2011	2010	2011
<b>ILLCIT DRUGS<sup>1</sup></b>	0.3009		0.9509		0.6413		0.2270	
Marijuana and Hashish	0.7394		0.9414		0.8426		0.0096	
Cocaine	0.4896		0.8740		0.3459		0.0581	
Crack	0.6837		0.3124		*		*	
Heroin	0.6882		0.1959		*		0.4644	
Hallucinogens	0.3391		0.9492		0.1307		*	
LSD	0.9487		*		*		*	
PCP	*		*		*		*	
Ecstasy	0.8552		*		0.2362		*	
Inhalants	0.6376		*		0.4017		0.2664	
Nonmedical Use of Psychotherapeutics <sup>2,3</sup>	0.1567		0.2883		0.0709		0.7474	
Pain Relievers	0.0630		0.0732		0.2283		0.9501	
OxyContin <sup>®</sup>	0.2693		0.3848		0.7900		*	
Tranquilizers	0.1469		0.3148		0.4960		0.6086	
Stimulants <sup>3</sup>	0.9035		0.8855		0.3964		0.6304	
Methamphetamine <sup>3</sup>	0.1419		0.7258		0.2420		0.3249	
Sedatives	0.5011		0.3544		0.2335		0.3542	
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	0.0534		0.3577		0.1681		0.7738	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>3</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.23C Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: Standard Errors of Numbers in Thousands, 2010 and 2011**

<b>Cigarette Use/Level of Alcohol Use</b>	<b>Total (2010)</b>	<b>Total (2011)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 12-17 (2011)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 18-25 (2011)</b>	<b>Aged 26+ (2010)</b>	<b>Aged 26+ (2011)</b>
<b>TOTAL</b>	488	460	70	68	136	142	441	420
Heavy Alcohol Use <sup>1</sup>	217	214	24	20	94	97	189	186
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	246	228	41	32	94	101	222	202
Alcohol Use But Not Binge Use <sup>1</sup>	247	222	25	30	58	67	234	215
No Alcohol Use	227	225	48	50	58	57	213	208
<b>PAST MONTH CIGARETTE USE</b>	355	342	48	47	132	133	318	299
Heavy Alcohol Use <sup>1</sup>	173	166	22	17	76	70	156	144
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	172	161	31	26	75	75	149	139
Alcohol Use But Not Binge Use <sup>1</sup>	150	142	16	20	41	46	140	130
No Alcohol Use	168	147	22	26	45	37	160	140
<b>NO PAST MONTH CIGARETTE USE</b>	330	330	57	52	95	113	308	300
Heavy Alcohol Use <sup>1</sup>	109	126	11	10	44	50	98	116
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	169	162	25	20	55	62	159	148
Alcohol Use But Not Binge Use <sup>1</sup>	195	168	20	22	41	47	188	161
No Alcohol Use	146	154	42	42	38	43	135	140

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.23D Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: Standard Errors of Percentages, 2010 and 2011**

<b>Cigarette Use/Level of Alcohol Use</b>	<b>Total (2010)</b>	<b>Total (2011)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 12-17 (2011)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 18-25 (2011)</b>	<b>Aged 26+ (2010)</b>	<b>Aged 26+ (2011)</b>
<b>TOTAL</b>	0.19	0.18	0.29	0.27	0.40	0.42	0.23	0.21
Heavy Alcohol Use <sup>1</sup>	1.10	1.06	2.78	3.04	1.24	1.33	1.39	1.35
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	0.56	0.51	1.85	1.64	0.80	0.81	0.68	0.61
Alcohol Use But Not Binge Use <sup>1</sup>	0.33	0.30	1.53	1.70	0.74	0.81	0.35	0.32
No Alcohol Use	0.18	0.18	0.22	0.22	0.41	0.39	0.24	0.23
<b>PAST MONTH CIGARETTE USE</b>	0.54	0.54	1.57	1.49	0.78	0.80	0.64	0.64
Heavy Alcohol Use <sup>1</sup>	1.52	1.54	3.16	3.13	1.51	1.55	2.14	2.05
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	0.95	0.95	2.70	2.37	1.30	1.28	1.19	1.23
Alcohol Use But Not Binge Use <sup>1</sup>	1.08	1.01	4.22	4.12	1.75	1.86	1.20	1.11
No Alcohol Use	0.81	0.74	2.19	2.43	1.47	1.29	0.93	0.85
<b>NO PAST MONTH CIGARETTE USE</b>	0.17	0.16	0.24	0.21	0.37	0.41	0.20	0.19
Heavy Alcohol Use <sup>1</sup>	1.32	1.48	5.65	5.52	1.80	2.16	1.58	1.83
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	0.65	0.58	2.30	2.04	0.97	0.98	0.78	0.68
Alcohol Use But Not Binge Use <sup>1</sup>	0.32	0.27	1.55	1.70	0.73	0.81	0.34	0.29
No Alcohol Use	0.14	0.15	0.20	0.19	0.34	0.37	0.19	0.19

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.23P Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: *P* Values from Tests of Differences of Percentages, 2011 Versus 2010**

<b>Cigarette Use/Level of Alcohol Use</b>	<b>Total (2010)</b>	<b>Total (2011)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 12-17 (2011)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 18-25 (2011)</b>	<b>Aged 26+ (2010)</b>	<b>Aged 26+ (2011)</b>
<b>TOTAL</b>	0.4074		0.9273		0.8206		0.4496	
Heavy Alcohol Use <sup>1</sup>	0.7549		0.9692		0.8813		0.9209	
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	0.4477		0.0913		0.5540		0.3950	
Alcohol Use But Not Binge Use <sup>1</sup>	0.8166		0.0821		0.1031		0.4490	
No Alcohol Use	0.6149		0.4934		0.2888		0.5395	
<b>PAST MONTH CIGARETTE USE</b>	0.5443		0.0223		0.5297		0.4582	
Heavy Alcohol Use <sup>1</sup>	0.2677		0.1711		0.5891		0.2145	
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	0.5813		0.5182		0.9049		0.7863	
Alcohol Use But Not Binge Use <sup>1</sup>	0.7504		0.2973		0.4173		0.5193	
No Alcohol Use	0.9298		0.0395		0.0529		0.9509	
<b>NO PAST MONTH CIGARETTE USE</b>	0.7422		0.6525		0.3475		0.8585	
Heavy Alcohol Use <sup>1</sup>	0.3967		0.3158		0.3871		0.1963	
Binge Alcohol Use But Not Heavy Use <sup>1</sup>	0.4257		0.0092		0.2111		0.3753	
Alcohol Use But Not Binge Use <sup>1</sup>	0.9066		0.2026		0.1524		0.5868	
No Alcohol Use	0.1936		0.7842		0.6934		0.2270	

\*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.