

**Table 6.24C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2010 and 2011**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	371	367
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	176	161	256	223
Cigars	263	231	258	290
Pipe Tobacco	90	95	123	110
<b>ALCOHOL</b>	707	716	1,447	1,529
Binge Alcohol Use <sup>2</sup>	515	491	688	717
Heavy Alcohol Use <sup>2</sup>	295	282	326	324

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.24D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2010 and 2011**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	0.18	0.18
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.30	0.27	0.13	0.11
Cigars	0.42	0.39	0.13	0.14
Pipe Tobacco	0.15	0.16	0.06	0.05
<b>ALCOHOL</b>	0.70	0.71	0.46	0.47
Binge Alcohol Use <sup>2</sup>	0.70	0.67	0.30	0.30
Heavy Alcohol Use <sup>2</sup>	0.47	0.46	0.16	0.16

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.24P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A		0.6206	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.2804		0.1776	
Cigars	0.3125		0.8646	
Pipe Tobacco	0.9975		0.9329	
<b>ALCOHOL</b>	0.7406		0.8664	
Binge Alcohol Use <sup>2</sup>	0.2281		0.8104	
Heavy Alcohol Use <sup>2</sup>	0.4647		0.0917	

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.25C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2010 and 2011**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	31	31
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	22	22	22	19
Cigars	28	32	23	24
Pipe Tobacco	15	14	9	11
<b>ALCOHOL</b>	55	46	66	70
Binge Alcohol Use <sup>2</sup>	46	37	44	44
Heavy Alcohol Use <sup>2</sup>	24	19	14	15

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.25D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Percentages, 2010 and 2011**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	0.14	0.13
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.98	1.04	0.10	0.08
Cigars	1.28	1.35	0.10	0.10
Pipe Tobacco	0.71	0.71	0.04	0.05
<b>ALCOHOL</b>	1.54	1.53	0.27	0.28
Binge Alcohol Use <sup>2</sup>	1.51	1.42	0.19	0.19
Heavy Alcohol Use <sup>2</sup>	1.05	0.94	0.06	0.07

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.25P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A		0.3666	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.8185		0.0546	
Cigars	0.0069		0.7399	
Pipe Tobacco	0.4488		0.2499	
<b>ALCOHOL</b>	0.8250		0.9195	
Binge Alcohol Use <sup>2</sup>	0.3879		0.8306	
Heavy Alcohol Use <sup>2</sup>	0.4298		0.9462	

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.26C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2010 and 2011**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	90	82
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	63	58	51	42
Cigars	83	94	72	66
Pipe Tobacco	34	37	23	33
<b>ALCOHOL</b>	188	204	259	290
Binge Alcohol Use <sup>2</sup>	163	171	179	203
Heavy Alcohol Use <sup>2</sup>	105	91	87	92

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.26D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Percentages, 2010 and 2011**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	0.35	0.33
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.50	0.47	0.22	0.18
Cigars	0.61	0.65	0.30	0.27
Pipe Tobacco	0.29	0.31	0.10	0.15
<b>ALCOHOL</b>	0.65	0.69	0.61	0.67
Binge Alcohol Use <sup>2</sup>	0.72	0.77	0.58	0.63
Heavy Alcohol Use <sup>2</sup>	0.70	0.65	0.35	0.36

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.26P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A		0.0374	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.0920		0.0046	
Cigars	0.6380		0.0717	
Pipe Tobacco	0.6199		0.2722	
<b>ALCOHOL</b>	0.3705		0.6988	
Binge Alcohol Use <sup>2</sup>	0.1640		0.8902	
Heavy Alcohol Use <sup>2</sup>	0.0176		0.1000	

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.27C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2010 and 2011**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	356	360
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	161	146	245	218
Cigars	245	206	246	281
Pipe Tobacco	82	87	121	104
<b>ALCOHOL</b>	668	659	1,406	1,459
Binge Alcohol Use <sup>2</sup>	479	445	658	671
Heavy Alcohol Use <sup>2</sup>	278	266	310	302

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.27D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2010 and 2011**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	0.23	0.23
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.36	0.33	0.16	0.14
Cigars	0.52	0.46	0.16	0.18
Pipe Tobacco	0.18	0.20	0.08	0.07
<b>ALCOHOL</b>	0.89	0.89	0.58	0.59
Binge Alcohol Use <sup>2</sup>	0.87	0.81	0.37	0.38
Heavy Alcohol Use <sup>2</sup>	0.59	0.57	0.20	0.19

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

**Table 6.27P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2011 Versus 2010**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2010	2011	2010	2011
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A		0.9297	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.5354		0.5265	
Cigars	0.1208		0.7536	
Pipe Tobacco	0.9414		0.6414	
<b>ALCOHOL</b>	0.6359		0.7825	
Binge Alcohol Use <sup>2</sup>	0.3505		0.7832	
Heavy Alcohol Use <sup>2</sup>	0.9697		0.1891	

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.