

Table 6.28C Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2010 and 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	327	319	463	431	464	477	572	550
Cigarettes	295	282	407	385	434	450	528	516
Smokeless Tobacco	141	125	157	147	132	123	173	148
Cigars	164	136	212	209	166	215	178	175
Pipe Tobacco	44	39	62	70	79	85	103	86

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.28D Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2010 and 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	1.24	1.27	0.81	0.80	0.58	0.60	0.42	0.40
Cigarettes	1.27	1.32	0.77	0.77	0.55	0.58	0.40	0.38
Smokeless Tobacco	0.78	0.73	0.37	0.34	0.18	0.16	0.14	0.12
Cigars	0.88	0.77	0.49	0.47	0.23	0.28	0.14	0.14
Pipe Tobacco	0.26	0.25	0.15	0.16	0.11	0.11	0.08	0.07

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.28P Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2011 Versus 2010

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	0.5756		0.0163		0.4835		0.0424	
Cigarettes	0.8607		0.0125		0.8381		0.1346	
Smokeless Tobacco	0.8977		0.2584		0.2408		0.4213	
Cigars	0.0940		0.0360		0.0346		0.7238	
Pipe Tobacco	0.7289		0.3872		0.7526		0.1947	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.29C Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2010 and 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	25	21	42	36	25	29	45	47
Cigarettes	24	19	38	32	23	25	38	40
Smokeless Tobacco	10	14	19	14	12	13	19	17
Cigars	13	14	23	23	13	17	22	25
Pipe Tobacco	7	7	11	13	6	4	10	10

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.29D Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2010 and 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	2.77	2.71	1.86	1.71	1.54	1.64	0.21	0.21
Cigarettes	2.98	3.16	1.82	1.67	1.45	1.53	0.18	0.19
Smokeless Tobacco	2.33	3.15	1.17	0.96	0.81	0.85	0.09	0.08
Cigars	2.91	2.96	1.41	1.39	0.90	1.06	0.11	0.11
Pipe Tobacco	1.64	1.86	0.68	0.84	0.46	0.30	0.05	0.05

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.29P Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	0.7667		0.3317		0.6845		0.0784	
Cigarettes	0.3602		0.3960		0.6130		0.1809	
Smokeless Tobacco	0.0500		0.0724		0.8236		0.1341	
Cigars	0.3267		0.5791		0.1980		0.6796	
Pipe Tobacco	0.6077		0.0463		0.5460		0.7623	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.30C Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2010 and 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	123	103	126	142	79	88	93	100
Cigarettes	105	91	111	129	69	79	87	90
Smokeless Tobacco	51	39	48	47	24	18	31	28
Cigars	67	58	62	70	38	42	44	48
Pipe Tobacco	19	20	27	31	16	21	17	21

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.30D Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2010 and 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	1.10	1.30	0.83	0.88	0.93	1.00	0.61	0.65
Cigarettes	1.30	1.36	0.86	0.90	0.84	0.93	0.58	0.60
Smokeless Tobacco	0.92	0.85	0.49	0.47	0.33	0.25	0.24	0.21
Cigars	1.10	1.09	0.61	0.64	0.52	0.57	0.32	0.34
Pipe Tobacco	0.40	0.48	0.29	0.31	0.23	0.29	0.13	0.16

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.30P Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2011 Versus 2010

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	0.3995		0.0360		0.8006		0.9681	
Cigarettes	0.8426		0.2240		0.7690		0.8822	
Smokeless Tobacco	0.1056		0.1341		0.0584		0.2797	
Cigars	0.6590		0.0533		0.1218		0.8011	
Pipe Tobacco	0.5914		0.6597		0.2419		0.7411	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.31C Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2010 and 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	307	303	441	391	455	465	551	533
Cigarettes	278	266	389	353	427	438	512	496
Smokeless Tobacco	132	121	146	137	128	122	168	145
Cigars	151	125	204	189	161	209	168	167
Pipe Tobacco	41	33	56	60	77	82	101	83

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.31D Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2010 and 2011

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	1.71	1.70	1.05	1.02	0.64	0.67	0.57	0.54
Cigarettes	1.71	1.74	0.99	0.97	0.61	0.64	0.54	0.51
Smokeless Tobacco	1.05	1.00	0.47	0.43	0.20	0.18	0.19	0.16
Cigars	1.16	1.01	0.63	0.59	0.25	0.31	0.19	0.19
Pipe Tobacco	0.35	0.29	0.18	0.19	0.12	0.12	0.11	0.09

*Low precision; no estimate reported.

NOTE: Some 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.

Table 6.31P Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2011 Versus 2010

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2010	2011	2010	2011	2010	2011	2010	2011
TOBACCO PRODUCTS²	0.3833		0.0504		0.4444		0.0620	
Cigarettes	0.7074		0.0236		0.8681		0.1812	
Smokeless Tobacco	0.7985		0.5995		0.3902		0.6686	
Cigars	0.1070		0.0886		0.0796		0.6118	
Pipe Tobacco	0.9296		0.4192		0.9148		0.2204	

*Low precision; no estimate reported.

NOTE: Some 2010 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010 and 2011.