

Table 6.47C Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	98	100	17	15	28	27	92	93
From More Than One Doctor	29	36	6	5	9	8	27	35
Wrote Fake Prescription	11	8	2	3	2	4	10	7
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	11	15	5	4	6	5	7	14
From Friend or Relative for Free	151	158	23	24	58	54	138	146
Bought from Friend or Relative	62	70	10	10	28	27	54	63
Took from Friend or Relative without Asking	41	37	13	11	14	14	37	31
Bought from Drug Dealer or Other Stranger	46	30	6	7	17	17	44	22
Bought on the Internet	21	14	1	0	1	1	21	14
Some Other Way	49	47	8	10	12	13	46	44

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.47D Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	0.77	0.83	1.09	1.08	0.66	0.75	1.32	1.35
From More Than One Doctor	0.26	0.33	0.43	0.40	0.24	0.23	0.43	0.57
Wrote Fake Prescription	0.10	0.08	0.15	0.24	0.05	0.11	0.17	0.12
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.10	0.14	0.34	0.33	0.17	0.16	0.12	0.22
From Friend or Relative for Free	0.89	0.96	1.30	1.43	0.91	0.98	1.53	1.61
Bought from Friend or Relative	0.52	0.60	0.73	0.76	0.67	0.70	0.84	0.93
Took from Friend or Relative without Asking	0.36	0.33	0.87	0.84	0.37	0.40	0.58	0.50
Bought from Drug Dealer or Other Stranger	0.41	0.27	0.48	0.58	0.44	0.48	0.69	0.35
Bought on the Internet	0.18	0.13	0.06	0.03	0.02	0.04	0.33	0.22
Some Other Way	0.43	0.42	0.57	0.77	0.31	0.38	0.73	0.69

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.47P Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	0.6776		0.5418		0.3104		0.8706	
From More Than One Doctor	0.9075		0.9499		0.3448		0.9386	
Wrote Fake Prescription	0.9236		0.8347		0.0664		0.5813	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.6829		0.4344		0.3712		0.7370	
From Friend or Relative for Free	0.4235		0.4373		0.0993		0.6140	
Bought from Friend or Relative	0.0039		0.6543		0.2038		0.0060	
Took from Friend or Relative without Asking	0.1807		0.0420		0.9667		0.4662	
Bought from Drug Dealer or Other Stranger	0.0659		0.9072		0.6579		0.0883	
Bought on the Internet	0.4147		0.4593		0.2291		0.3563	
Some Other Way	0.8784		0.0784		0.4143		0.6826	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.48C Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	132	137	17	16	46	44	123	130
From More Than One Doctor	30	22	5	6	9	11	28	17
Wrote Fake Prescription	11	11	0	1	1	2	11	10
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	3	5	1	4	3	4	*	2
From Friend or Relative for Free	30	34	7	6	14	13	26	31
Bought from Friend or Relative	23	20	6	5	13	13	18	14
Took from Friend or Relative without Asking	14	13	7	7	5	8	10	8
Bought from Drug Dealer or Other Stranger	14	15	3	5	9	8	10	12
Bought on the Internet	4	5	1	0	2	1	4	5
Some Other Way	11	15	3	5	6	5	8	14

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.48D Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	0.92	0.95	2.09	2.25	1.15	1.38	1.33	1.29
From More Than One Doctor	0.53	0.41	0.89	1.22	0.48	0.67	0.84	0.55
Wrote Fake Prescription	0.19	0.20	0.06	0.12	0.06	0.12	0.33	0.33
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.06	0.10	0.21	0.74	0.16	0.22	*	0.05
From Friend or Relative for Free	0.55	0.65	1.37	1.23	0.76	0.80	0.80	0.96
Bought from Friend or Relative	0.40	0.38	1.08	1.07	0.66	0.79	0.55	0.43
Took from Friend or Relative without Asking	0.25	0.25	1.27	1.33	0.29	0.47	0.32	0.27
Bought from Drug Dealer or Other Stranger	0.25	0.30	0.67	1.00	0.48	0.51	0.31	0.37
Bought on the Internet	0.07	0.09	0.11	0.05	0.10	0.04	0.11	0.15
Some Other Way	0.20	0.29	0.64	0.95	0.34	0.30	0.25	0.43

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.48P Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	0.2306		0.5787		0.6135		0.0854	
From More Than One Doctor	0.0665		0.3980		0.4447		0.0105	
Wrote Fake Prescription	0.6656		0.4254		0.4551		0.7864	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.1933		0.4278		0.4245		*	
From Friend or Relative for Free	0.4150		0.1656		0.5245		0.8267	
Bought from Friend or Relative	0.4590		0.7997		0.5964		0.2585	
Took from Friend or Relative without Asking	0.7097		0.8184		0.0865		0.4406	
Bought from Drug Dealer or Other Stranger	0.9616		0.6192		0.7529		0.8974	
Bought on the Internet	0.9929		0.5536		0.3055		0.7318	
Some Other Way	0.3122		0.1068		0.2744		0.2424	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.49C Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	81	75	11	10	18	20	79	72
From More Than One Doctor	34	33	4	5	10	8	32	31
Wrote Fake Prescription	8	12	2	2	2	2	7	12
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	8	15	4	2	3	5	7	14
From Friend or Relative for Free	104	103	15	15	35	36	98	95
Bought from Friend or Relative	67	66	9	8	26	24	60	59
Took from Friend or Relative without Asking	38	46	10	8	14	16	34	42
Bought from Drug Dealer or Other Stranger	41	39	7	7	18	20	35	32
Bought on the Internet	20	15	0	1	3	5	20	14
Some Other Way	30	33	6	5	9	10	29	31

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.49D Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	1.42	1.47	1.91	1.87	1.13	1.43	2.31	2.35
From More Than One Doctor	0.73	0.75	0.86	0.98	0.67	0.63	1.17	1.16
Wrote Fake Prescription	0.17	0.28	0.39	0.51	0.12	0.14	0.27	0.45
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.19	0.35	0.82	0.39	0.24	0.36	0.25	0.55
From Friend or Relative for Free	1.42	1.50	2.11	2.25	1.32	1.59	2.39	2.35
Bought from Friend or Relative	1.30	1.29	1.56	1.53	1.44	1.57	2.05	1.98
Took from Friend or Relative without Asking	0.82	1.00	1.69	1.56	0.91	1.13	1.27	1.53
Bought from Drug Dealer or Other Stranger	0.85	0.86	1.33	1.33	1.12	1.29	1.29	1.21
Bought on the Internet	0.44	0.34	0.03	0.12	0.21	0.43	0.75	0.53
Some Other Way	0.67	0.76	1.14	1.08	0.64	0.79	1.09	1.19

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.49P Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	0.5446		0.3241		0.0846		0.2726	
From More Than One Doctor	0.4948		0.2990		0.2715		0.5524	
Wrote Fake Prescription	0.5408		0.6179		0.6968		0.5005	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.5403		0.0206		0.3360		0.3180	
From Friend or Relative for Free	0.7152		0.3243		0.8606		0.5489	
Bought from Friend or Relative	0.7311		0.2817		1.0000		0.5411	
Took from Friend or Relative without Asking	0.4407		0.0102		0.0663		0.4296	
Bought from Drug Dealer or Other Stranger	0.6133		0.1532		0.1051		0.3426	
Bought on the Internet	0.8150		0.1202		0.1771		0.5068	
Some Other Way	0.3984		0.5169		0.6376		0.3822	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.50C Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	89	90	10	9	27	28	84	85
From More Than One Doctor	34	22	4	6	10	11	32	18
Wrote Fake Prescription	13	15	2	0	4	2	12	15
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	8	7	2	1	4	4	7	6
From Friend or Relative for Free	31	32	7	4	12	13	28	29
Bought from Friend or Relative	29	27	6	4	12	14	26	23
Took from Friend or Relative without Asking	16	16	5	5	6	8	14	13
Bought from Drug Dealer or Other Stranger	24	25	5	5	11	14	21	20
Bought on the Internet	4	3	1	0	3	2	3	1
Some Other Way	14	16	3	4	6	6	12	14

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.50D Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	1.42	1.60	3.11	3.41	1.70	2.10	2.08	2.33
From More Than One Doctor	1.28	0.92	2.08	2.56	1.12	1.35	2.04	1.29
Wrote Fake Prescription	0.51	0.64	0.83	0.05	0.49	0.28	0.82	1.07
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.34	0.31	1.07	0.31	0.44	0.59	0.49	0.41
From Friend or Relative for Free	1.21	1.29	2.93	2.11	1.35	1.59	1.86	1.99
Bought from Friend or Relative	1.15	1.14	2.73	2.05	1.33	1.68	1.74	1.62
Took from Friend or Relative without Asking	0.66	0.67	2.28	2.15	0.71	1.08	0.96	0.93
Bought from Drug Dealer or Other Stranger	0.96	1.05	2.25	2.37	1.28	1.68	1.44	1.43
Bought on the Internet	0.18	0.11	0.70	0.09	0.35	0.32	0.20	0.07
Some Other Way	0.58	0.68	1.56	1.97	0.74	0.80	0.84	1.02

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.50P Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Source Where Friend or Relative Obtained Pain Relievers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	0.7893		0.1932		0.3403		0.6626	
From More Than One Doctor	0.0245		0.9961		0.8346		0.0167	
Wrote Fake Prescription	0.7376		0.0234		0.2345		0.4196	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.9416		0.0383		0.6931		0.8533	
From Friend or Relative for Free	0.8310		0.0011		0.8374		0.7364	
Bought from Friend or Relative	0.9721		0.0450		0.2243		0.8780	
Took from Friend or Relative without Asking	0.9430		0.4753		0.2143		0.7587	
Bought from Drug Dealer or Other Stranger	0.9361		0.3049		0.7253		0.8488	
Bought on the Internet	0.1555		0.0772		0.6239		0.3881	
Some Other Way	0.7040		0.8631		0.9828		0.6404	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.51C Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	43	63	5	5	12	12	41	62
From More Than One Doctor	17	15	2	2	6	5	16	14
Wrote Fake Prescription	1	1	1	1	1	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	6	2	1	3	5	*	4
From Friend or Relative for Free	115	115	12	14	35	36	110	107
Bought from Friend or Relative	41	48	9	8	21	20	34	43
Took from Friend or Relative without Asking	28	19	6	5	9	9	25	16
Bought from Drug Dealer or Other Stranger	27	28	6	6	14	14	22	22
Bought on the Internet	41	14	1	1	1	2	41	*
Some Other Way	34	19	3	4	9	8	32	18

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.51D Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	0.82	1.19	1.25	1.31	0.72	0.78	1.37	1.88
From More Than One Doctor	0.34	0.31	0.56	0.63	0.35	0.34	0.54	0.47
Wrote Fake Prescription	0.03	0.03	0.24	0.27	0.05	0.06	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.08	0.13	0.58	0.18	0.21	0.34	*	0.12
From Friend or Relative for Free	1.46	1.53	2.37	2.63	1.38	1.49	2.32	2.30
Bought from Friend or Relative	0.81	0.92	1.93	1.90	1.07	1.14	1.15	1.35
Took from Friend or Relative without Asking	0.56	0.39	1.39	1.30	0.54	0.58	0.84	0.53
Bought from Drug Dealer or Other Stranger	0.53	0.56	1.34	1.61	0.78	0.85	0.77	0.73
Bought on the Internet	0.83	0.29	0.28	0.14	0.07	0.12	1.40	*
Some Other Way	0.67	0.39	0.81	1.01	0.55	0.49	1.09	0.58

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.51P Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	0.0041		0.6144		0.5244		0.0074	
From More Than One Doctor	0.8164		0.9947		0.9235		0.8276	
Wrote Fake Prescription	0.7311		0.6734		0.8580		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.8101		0.2291		0.7879		*	
From Friend or Relative for Free	0.6723		0.4427		0.9218		0.4619	
Bought from Friend or Relative	0.9750		0.5787		0.4698		0.4916	
Took from Friend or Relative without Asking	0.3193		0.2318		0.7064		0.3536	
Bought from Drug Dealer or Other Stranger	0.9281		0.9706		0.4761		0.7720	
Bought on the Internet	0.2162		0.5871		0.8112		*	
Some Other Way	0.0655		0.5274		0.1633		0.1190	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.52C Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	97	103	8	9	25	28	95	98
From More Than One Doctor	17	14	3	3	5	7	16	12
Wrote Fake Prescription	2	1	1	*	1	1	2	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	4	1	*	3	1	3	4
From Friend or Relative for Free	25	27	4	5	12	12	22	24
Bought from Friend or Relative	17	15	3	4	10	7	13	12
Took from Friend or Relative without Asking	5	6	4	2	3	5	2	3
Bought from Drug Dealer or Other Stranger	8	13	2	4	6	7	4	10
Bought on the Internet	3	4	*	*	2	1	2	4
Some Other Way	9	8	3	3	3	2	8	7

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.52D Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	1.43	1.31	3.63	4.10	1.95	1.99	1.84	1.74
From More Than One Doctor	0.66	0.51	2.04	1.79	0.69	0.92	0.98	0.66
Wrote Fake Prescription	0.09	0.06	0.91	*	0.07	0.20	0.12	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.17	0.15	0.68	*	0.39	0.08	0.18	0.23
From Friend or Relative for Free	0.94	0.98	2.54	2.73	1.46	1.55	1.29	1.31
Bought from Friend or Relative	0.64	0.53	2.08	2.54	1.26	0.88	0.76	0.69
Took from Friend or Relative without Asking	0.20	0.23	2.28	1.50	0.40	0.62	0.10	0.16
Bought from Drug Dealer or Other Stranger	0.30	0.47	0.98	2.31	0.81	0.92	0.25	0.55
Bought on the Internet	0.11	0.15	*	*	0.26	0.11	0.12	0.21
Some Other Way	0.35	0.31	1.69	2.11	0.35	0.25	0.49	0.40

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.52P Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	0.0468		0.3481		0.0818		0.3191	
From More Than One Doctor	0.0242		0.8616		0.1831		0.0032	
Wrote Fake Prescription	0.3105		*		0.3833		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.5221		*		0.1625		0.6980	
From Friend or Relative for Free	0.6097		0.6135		0.2219		0.7840	
Bought from Friend or Relative	0.0918		0.5977		0.0063		0.9267	
Took from Friend or Relative without Asking	0.7260		0.0893		0.4209		0.6121	
Bought from Drug Dealer or Other Stranger	0.2794		0.0965		0.6077		0.5225	
Bought on the Internet	0.6457		*		0.4634		0.4880	
Some Other Way	0.5695		0.6429		0.3480		0.6122	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.53C Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	27	36	3	3	9	9	26	36
From More Than One Doctor	10	5	*	2	3	4	9	2
Wrote Fake Prescription	1	1	1	1	*	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	3	13	*	1	1	3	*	*
From Friend or Relative for Free	72	75	7	6	20	20	68	72
Bought from Friend or Relative	33	41	4	5	14	12	30	38
Took from Friend or Relative without Asking	15	18	4	3	6	8	12	16
Bought from Drug Dealer or Other Stranger	21	16	4	5	10	10	19	12
Bought on the Internet	*	7	1	0	1	2	*	7
Some Other Way	15	11	2	2	4	3	15	11

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.53D Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	1.48	1.85	2.42	2.65	1.45	1.63	2.34	2.73
From More Than One Doctor	0.54	0.29	*	1.81	0.55	0.84	0.82	0.18
Wrote Fake Prescription	0.04	0.08	0.56	0.68	*	0.24	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.14	0.72	*	0.55	0.25	0.54	*	*
From Friend or Relative for Free	2.37	2.43	3.94	4.16	2.37	2.52	3.51	3.46
Bought from Friend or Relative	1.77	1.97	2.84	3.85	2.09	2.07	2.62	2.86
Took from Friend or Relative without Asking	0.79	0.99	3.04	2.72	1.10	1.47	1.10	1.34
Bought from Drug Dealer or Other Stranger	1.17	0.92	2.70	3.85	1.65	1.76	1.68	1.01
Bought on the Internet	*	0.39	0.97	0.33	0.13	0.49	*	0.56
Some Other Way	0.87	0.59	1.57	2.02	0.67	0.57	1.34	0.89

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.53P Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	0.0214		0.8497		0.3739		0.0449	
From More Than One Doctor	0.0918		*		0.7889		0.0259	
Wrote Fake Prescription	0.1752		0.7849		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2160		*		0.3728		*	
From Friend or Relative for Free	0.6625		0.8049		0.5692		0.6636	
Bought from Friend or Relative	0.8020		0.0694		0.6671		0.7695	
Took from Friend or Relative without Asking	0.8791		0.0404		0.2502		0.9737	
Bought from Drug Dealer or Other Stranger	0.2285		0.9214		0.3601		0.1008	
Bought on the Internet	*		0.2439		0.2454		*	
Some Other Way	0.3786		0.7451		0.2458		0.5214	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.54C Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	61	65	*	*	15	16	58	64
From More Than One Doctor	17	15	*	*	5	5	16	14
Wrote Fake Prescription	1	1	*	*	1	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	4	*	*	1	2	*	*
From Friend or Relative for Free	18	25	*	*	10	7	14	24
Bought from Friend or Relative	10	13	*	*	7	4	6	12
Took from Friend or Relative without Asking	6	10	*	*	2	3	5	*
Bought from Drug Dealer or Other Stranger	12	14	*	*	3	5	11	13
Bought on the Internet	1	4	*	*	1	*	0	*
Some Other Way	10	6	*	*	3	2	*	6

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.54D Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	3.08	2.55	*	*	3.55	3.60	4.21	3.36
From More Than One Doctor	1.66	1.51	*	*	1.75	2.25	2.33	1.96
Wrote Fake Prescription	0.05	0.07	*	*	0.19	0.27	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.17	0.43	*	*	0.46	0.79	*	*
From Friend or Relative for Free	1.78	2.42	*	*	3.11	2.82	2.10	3.23
Bought from Friend or Relative	1.12	1.30	*	*	2.53	1.92	0.98	1.63
Took from Friend or Relative without Asking	0.60	1.00	*	*	0.73	1.38	0.77	*
Bought from Drug Dealer or Other Stranger	1.17	1.43	*	*	1.11	1.99	1.67	1.82
Bought on the Internet	0.08	0.40	*	*	0.29	*	0.01	*
Some Other Way	1.06	0.67	*	*	1.06	0.71	*	0.88

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.54P Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Source Where Friend or Relative Obtained Tranquilizers ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	0.3009		*		0.1698		0.7062	
From More Than One Doctor	0.2882		*		0.7302		0.2639	
Wrote Fake Prescription	0.6907		*		0.5578		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3672		*		0.3702		*	
From Friend or Relative for Free	0.6116		*		0.2137		0.2030	
Bought from Friend or Relative	0.5312		*		0.0944		0.3133	
Took from Friend or Relative without Asking	0.4699		*		0.1524		*	
Bought from Drug Dealer or Other Stranger	0.4994		*		0.1288		0.7600	
Bought on the Internet	0.1700		*		*		*	
Some Other Way	0.4495		*		0.4825		*	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.55C Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Stimulant Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	27	28	5	3	10	9	24	26
From More Than One Doctor	11	4	1	2	3	3	11	*
Wrote Fake Prescription	7	2	2	2	1	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	7	8	3	1	3	2	*	*
From Friend or Relative for Free	58	62	10	9	29	28	49	54
Bought from Friend or Relative	31	31	5	6	20	19	22	24
Took from Friend or Relative without Asking	20	15	4	4	7	6	19	13
Bought from Drug Dealer or Other Stranger	16	30	4	4	10	9	13	29
Bought on the Internet	11	7	2	1	4	3	10	7
Some Other Way	11	17	3	2	6	5	9	16

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.55D Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Stimulant Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	1.22	1.25	1.85	1.64	0.98	0.90	2.57	2.74
From More Than One Doctor	0.54	0.19	0.58	1.13	0.34	0.32	1.19	*
Wrote Fake Prescription	0.34	0.09	0.73	0.89	0.07	0.01	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.34	0.40	1.19	0.54	0.31	0.19	*	*
From Friend or Relative for Free	1.89	1.96	2.94	3.44	1.97	1.73	3.81	4.18
Bought from Friend or Relative	1.36	1.43	1.96	2.80	1.75	1.62	2.33	2.51
Took from Friend or Relative without Asking	0.92	0.71	1.63	1.69	0.68	0.59	2.00	1.52
Bought from Drug Dealer or Other Stranger	0.75	1.39	1.64	1.89	0.99	0.87	1.35	3.08
Bought on the Internet	0.52	0.36	0.64	0.35	0.41	0.30	1.12	0.75
Some Other Way	0.54	0.81	1.16	0.97	0.65	0.55	1.00	1.77

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

**Table 6.55P Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009**

Source for Most Recent Nonmedical Use among Past Year Stimulant Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	0.8178		0.8332		0.3311		0.4983	
From More Than One Doctor	0.1580		0.4608		0.9397		*	
Wrote Fake Prescription	0.2766		0.8717		0.1456		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.6395		0.1375		0.4670		*	
From Friend or Relative for Free	0.9637		0.5476		0.5567		0.6132	
Bought from Friend or Relative	0.2612		0.2000		0.2144		0.8113	
Took from Friend or Relative without Asking	0.1769		0.4160		0.8541		0.2791	
Bought from Drug Dealer or Other Stranger	0.7573		0.5200		0.2573		0.3325	
Bought on the Internet	0.3497		0.6765		0.2898		0.5856	
Some Other Way	0.5982		0.2057		0.5026		0.2509	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.56C Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Stimulants ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	45	56	8	7	23	23	38	50
From More Than One Doctor	10	6	2	2	3	6	9	2
Wrote Fake Prescription	*	1	*	*	*	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	1	0	*	2	1	*	*
From Friend or Relative for Free	15	12	3	3	7	7	13	10
Bought from Friend or Relative	9	9	3	2	7	7	5	5
Took from Friend or Relative without Asking	3	4	2	*	2	2	1	*
Bought from Drug Dealer or Other Stranger	9	10	1	2	4	5	*	*
Bought on the Internet	6	3	*	0	1	*	*	3
Some Other Way	10	1	0	1	2	1	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.56D Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Stimulants ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	2.31	2.00	4.65	5.04	2.34	2.43	4.57	3.13
From More Than One Doctor	1.02	0.68	1.86	2.80	0.75	1.18	2.14	0.52
Wrote Fake Prescription	*	0.07	*	*	*	0.10	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.25	0.14	0.47	*	0.53	0.29	*	*
From Friend or Relative for Free	1.52	1.23	2.96	3.08	1.56	1.46	2.95	2.19
Bought from Friend or Relative	0.94	0.95	2.72	2.27	1.54	1.51	1.27	1.26
Took from Friend or Relative without Asking	0.34	0.38	2.11	*	0.44	0.53	0.20	*
Bought from Drug Dealer or Other Stranger	0.98	0.97	1.53	2.05	0.85	1.00	*	*
Bought on the Internet	0.62	0.29	*	0.55	0.15	*	*	0.65
Some Other Way	1.06	0.10	0.43	0.98	0.35	0.13	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.56P Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Source Where Friend or Relative Obtained Stimulants ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	0.1320		0.3120		0.6262		0.0137	
From More Than One Doctor	0.6535		0.3492		0.1873		0.0864	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.5366		*		0.5823		*	
From Friend or Relative for Free	0.1849		0.9462		0.7636		0.1568	
Bought from Friend or Relative	0.8114		0.6201		0.7277		0.8806	
Took from Friend or Relative without Asking	0.5642		*		0.5789		*	
Bought from Drug Dealer or Other Stranger	0.8689		0.6236		0.9066		*	
Bought on the Internet	0.6670		*		*		*	
Some Other Way	0.1244		0.4746		0.2383		*	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.57C Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Stimulant Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	21	25	4	3	8	8	19	*
From More Than One Doctor	12	3	1	1	3	3	*	*
Wrote Fake Prescription	2	2	2	*	1	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	7	2	2	0	3	2	*	*
From Friend or Relative for Free	36	31	6	4	15	17	32	26
Bought from Friend or Relative	24	23	3	4	12	13	21	19
Took from Friend or Relative without Asking	18	11	3	2	5	6	*	9
Bought from Drug Dealer or Other Stranger	15	16	3	4	8	8	13	13
Bought on the Internet	11	4	1	0	2	3	*	3
Some Other Way	6	9	2	2	3	2	*	9

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*) and because respondents who were logically assigned through the editing process as reporting only methamphetamine use were not previously excluded.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.57D Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Stimulant Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	2.40	2.98	3.66	3.71	2.18	2.26	4.45	*
From More Than One Doctor	1.43	0.44	1.71	1.99	0.93	0.91	*	*
Wrote Fake Prescription	0.27	0.24	2.13	*	0.27	0.03	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.81	0.23	2.35	0.46	0.77	0.53	*	*
From Friend or Relative for Free	3.25	3.16	4.59	5.30	3.20	3.14	6.00	5.81
Bought from Friend or Relative	2.73	2.68	3.32	4.82	2.90	3.04	4.90	4.65
Took from Friend or Relative without Asking	2.12	1.43	3.30	2.98	1.56	1.89	*	2.57
Bought from Drug Dealer or Other Stranger	1.84	1.96	3.35	4.60	2.28	2.16	3.15	3.55
Bought on the Internet	1.37	0.57	0.77	0.63	0.57	0.82	*	0.95
Some Other Way	0.76	1.25	2.00	2.66	0.98	0.57	*	2.53

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*) and because respondents who were logically assigned through the editing process as reporting only methamphetamine use were not previously excluded.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.57P Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Source for Past Month Nonmedical Use among Past Month Stimulant Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	0.1338		0.3704		0.7947		*	
From More Than One Doctor	0.1010		0.6594		0.5054		*	
Wrote Fake Prescription	0.3530		*		0.1156		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.1059		0.0585		0.6700		*	
From Friend or Relative for Free	0.8936		0.4693		0.3079		0.5919	
Bought from Friend or Relative	0.4051		0.2576		0.0842		0.8138	
Took from Friend or Relative without Asking	0.3080		0.5050		0.3235		*	
Bought from Drug Dealer or Other Stranger	0.5231		0.8058		0.9222		0.4409	
Bought on the Internet	0.3920		0.9306		0.9287		*	
Some Other Way	0.3692		0.9692		0.0985		*	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*) and because respondents who were logically assigned through the editing process as reporting only methamphetamine use were not previously excluded.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.58C Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Stimulants ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	26	28	*	*	11	14	*	*
From More Than One Doctor	6	5	*	*	2	5	*	*
Wrote Fake Prescription	*	0	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	1	*	*	*	*	*	*
From Friend or Relative for Free	17	8	*	*	6	5	*	*
Bought from Friend or Relative	12	6	*	*	5	4	*	*
Took from Friend or Relative without Asking	6	4	*	*	2	3	*	*
Bought from Drug Dealer or Other Stranger	7	7	*	*	3	2	*	*
Bought on the Internet	1	3	*	*	1	*	*	*
Some Other Way	*	1	*	*	0	0	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.58D Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Stimulants ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	4.79	3.26	*	*	4.13	4.06	*	*
From More Than One Doctor	1.77	1.69	*	*	1.32	3.16	*	*
Wrote Fake Prescription	*	0.03	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	0.27	*	*	*	*	*	*
From Friend or Relative for Free	4.32	2.43	*	*	3.59	3.11	*	*
Bought from Friend or Relative	3.28	2.01	*	*	3.35	2.60	*	*
Took from Friend or Relative without Asking	1.66	1.16	*	*	1.38	2.15	*	*
Bought from Drug Dealer or Other Stranger	1.96	2.11	*	*	2.00	1.58	*	*
Bought on the Internet	0.34	1.09	*	*	0.82	*	*	*
Some Other Way	*	0.46	*	*	0.26	0.09	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.58P Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Source Where Friend or Relative Obtained Stimulants ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	0.0217		*		0.3168		*	
From More Than One Doctor	0.7111		*		0.1315		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	0.0196		*		0.1750		*	
Bought from Friend or Relative	0.2167		*		0.8976		*	
Took from Friend or Relative without Asking	0.7953		*		0.3743		*	
Bought from Drug Dealer or Other Stranger	0.9272		*		0.6229		*	
Bought on the Internet	0.2226		*		*		*	
Some Other Way	*		*		0.5449		*	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.59C Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	32	36	6	*	14	12	28	33
Bought from Friend or Relative	40	50	3	*	7	8	39	*
Took from Friend or Relative without Asking	13	9	3	1	3	*	*	7
Bought from Drug Dealer or Other Stranger	25	35	3	*	8	8	24	*
Bought on the Internet	4	5	*	*	2	2	*	*
Some Other Way	11	8	0	*	3	3	11	6

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.59D Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	2.94	3.69	5.89	*	3.53	4.13	4.18	5.15
Bought from Friend or Relative	3.40	4.41	3.72	*	2.42	3.40	4.91	*
Took from Friend or Relative without Asking	1.44	1.05	3.70	1.30	1.13	*	*	1.21
Bought from Drug Dealer or Other Stranger	2.52	3.61	3.98	*	2.76	3.29	3.79	*
Bought on the Internet	0.47	0.55	*	*	0.62	0.78	*	*
Some Other Way	1.19	0.92	0.48	*	1.16	1.43	1.80	1.16

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.59P Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	0.9177		*		0.3525		0.6669	
Bought from Friend or Relative	0.9915		*		0.6788		*	
Took from Friend or Relative without Asking	0.5815		0.0404		*		*	
Bought from Drug Dealer or Other Stranger	0.8206		*		0.8123		*	
Bought on the Internet	0.6806		*		0.5467		*	
Some Other Way	0.5996		*		0.7794		0.3529	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.60C Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	21	20	*	*	9	8	*	*
Bought from Friend or Relative	15	18	*	*	7	5	*	*
Took from Friend or Relative without Asking	3	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	12	10	*	*	5	4	*	*
Bought on the Internet	*	1	*	*	*	*	*	*
Some Other Way	4	5	*	*	3	3	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.60D Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	4.72	5.38	*	*	5.75	5.72	*	*
Bought from Friend or Relative	4.24	5.26	*	*	5.12	5.01	*	*
Took from Friend or Relative without Asking	1.03	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	3.58	3.46	*	*	3.85	4.43	*	*
Bought on the Internet	*	0.49	*	*	*	*	*	*
Some Other Way	1.18	1.88	*	*	2.70	2.81	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.60P Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	0.7523		*		0.8277		*	
Bought from Friend or Relative	0.9329		*		0.2857		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	0.7466		*		0.6592		*	
Bought on the Internet	*		*		*		*	
Some Other Way	0.1054		*		0.5028		*	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.61C Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	27	26	*	*	*	*	*	*
Bought from Friend or Relative	27	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	4	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	26	19	*	*	*	*	*	*
Bought on the Internet	1	3	*	*	*	*	*	*
Some Other Way	11	4	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.61D Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	5.29	5.60	*	*	*	*	*	*
Bought from Friend or Relative	5.44	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	1.18	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	5.38	5.30	*	*	*	*	*	*
Bought on the Internet	0.15	1.02	*	*	*	*	*	*
Some Other Way	2.94	1.24	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.61P Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	0.8075		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	0.8084		*		*		*	
Bought on the Internet	0.0753		*		*		*	
Some Other Way	0.0953		*		*		*	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.62C Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.62D Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.62P Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Source Where Friend or Relative Obtained Methamphetamine ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

**Table 6.63C Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011**

Source for Most Recent Nonmedical Use among Past Year Sedative Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	17	26	3	3	5	5	16	*
From More Than One Doctor	10	*	1	*	*	1	9	*
Wrote Fake Prescription	3	1	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	2	0	2	*	*	*	0
From Friend or Relative for Free	37	49	7	7	11	10	34	*
Bought from Friend or Relative	14	13	3	3	5	4	*	12
Took from Friend or Relative without Asking	9	9	4	2	6	4	5	*
Bought from Drug Dealer or Other Stranger	8	9	2	1	2	2	8	8
Bought on the Internet	*	0	*	0	0	*	*	*
Some Other Way	12	8	1	*	2	1	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.63D Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Most Recent Nonmedical Use among Past Year Sedative Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	2.48	3.67	3.32	4.04	2.80	3.61	3.75	*
From More Than One Doctor	1.44	*	1.19	*	*	0.85	2.27	*
Wrote Fake Prescription	0.41	0.15	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.68	0.31	0.37	2.72	*	*	*	0.06
From Friend or Relative for Free	3.63	4.45	5.02	5.58	4.60	4.82	5.56	*
Bought from Friend or Relative	2.02	1.93	3.12	3.72	2.81	3.17	*	2.54
Took from Friend or Relative without Asking	1.34	1.38	3.90	2.17	3.61	2.84	1.20	*
Bought from Drug Dealer or Other Stranger	1.25	1.29	2.45	1.52	1.03	1.48	1.91	1.81
Bought on the Internet	*	0.01	*	0.05	0.06	*	*	*
Some Other Way	1.81	1.16	0.62	*	1.25	1.07	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

**Table 6.63P Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009**

Source for Most Recent Nonmedical Use among Past Year Sedative Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	0.1494		0.5679		0.7088		*	
From More Than One Doctor	*		*		*		*	
Wrote Fake Prescription	0.2700		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.4421		0.0563		*		*	
From Friend or Relative for Free	0.6472		0.6168		0.4647		*	
Bought from Friend or Relative	0.8911		0.5882		0.7105		*	
Took from Friend or Relative without Asking	0.3011		0.0243		0.4961		*	
Bought from Drug Dealer or Other Stranger	0.8494		0.5346		0.6186		0.8449	
Bought on the Internet	*		*		*		*	
Some Other Way	0.3542		*		0.4585		*	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.64C Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Sedatives ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	32	48	*	*	*	*	*	*
From More Than One Doctor	2	3	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	3	2	*	*	*	*	*	*
From Friend or Relative for Free	12	3	*	*	*	*	*	*
Bought from Friend or Relative	1	2	*	*	*	*	*	*
Took from Friend or Relative without Asking	3	3	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	2	4	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	2	2	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.64D Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Sedatives ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	3.85	2.41	*	*	*	*	*	*
From More Than One Doctor	0.53	0.84	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.96	0.52	*	*	*	*	*	*
From Friend or Relative for Free	3.33	0.76	*	*	*	*	*	*
Bought from Friend or Relative	0.43	0.65	*	*	*	*	*	*
Took from Friend or Relative without Asking	0.81	0.88	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	0.46	1.17	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	0.59	0.49	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.64P Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Source Where Friend or Relative Obtained Sedatives ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	0.1063		*		*		*	
From More Than One Doctor	0.4439		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2789		*		*		*	
From Friend or Relative for Free	0.0172		*		*		*	
Bought from Friend or Relative	0.2742		*		*		*	
Took from Friend or Relative without Asking	0.5078		*		*		*	
Bought from Drug Dealer or Other Stranger	0.3231		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	0.7098		*		*		*	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

¹ Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.65C Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Sedative Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	15	*	*	*	*	*	*	*
From More Than One Doctor	8	0	*	*	*	*	*	*
Wrote Fake Prescription	1	1	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	1	*	*	*	*	*	*
From Friend or Relative for Free	22	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	7	3	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	5	*	*	*	*	*	*
Bought on the Internet	0	0	*	*	*	*	*	*
Some Other Way	4	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.65D Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source for Past Month Nonmedical Use among Past Month Sedative Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	4.74	*	*	*	*	*	*	*
From More Than One Doctor	2.85	0.04	*	*	*	*	*	*
Wrote Fake Prescription	0.20	0.34	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.40	0.47	*	*	*	*	*	*
From Friend or Relative for Free	5.56	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	2.54	1.15	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	1.96	*	*	*	*	*	*
Bought on the Internet	0.18	0.04	*	*	*	*	*	*
Some Other Way	1.51	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.65P Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Source for Past Month Nonmedical Use among Past Month Sedative Users ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	*		*		*		*	
From More Than One Doctor	0.0163		*		*		*	
Wrote Fake Prescription	0.8124		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3752		*		*		*	
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	0.0466		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	0.3912		*		*		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.66C Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Sedatives ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.66D Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Source Where Friend or Relative Obtained Sedatives ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.66P Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Source Where Friend or Relative Obtained Sedatives ¹	Total (2008-2009)	Total (2010-2011)	Aged 12-17 (2008-2009)	Aged 12-17 (2010-2011)	Aged 18-25 (2008-2009)	Aged 18-25 (2010-2011)	Aged 26+ (2008-2009)	Aged 26+ (2010-2011)
From One Doctor	*		*		*		*	
From More Than One Doctor	*		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.