

Table 6.71C Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
ILLCIT DRUGS²	143	147	16	16	142	146
Marijuana and Hashish	119	120	15	15	118	118
Cocaine	39	34	2	1	38	34
Crack	19	18	*	1	19	18
Heroin	14	15	1	3	14	14
Hallucinogens	33	29	11	1	31	29
LSD	9	4	4	*	8	4
PCP	7	2	4	0	6	2
Ecstasy	28	22	11	1	26	22
Inhalants	17	18	4	1	17	18
Nonmedical Use of Psychotherapeutics ^{3,4}	88	85	6	6	87	84
Pain Relievers	74	65	6	6	73	64
OxyContin [®]	17	18	2	1	17	18
Tranquilizers	51	49	6	1	50	49
Stimulants ⁴	35	31	4	2	34	31
Methamphetamine ⁴	22	20	4	*	21	20
Sedatives	14	13	4	0	14	13
ILLCIT DRUGS OTHER THAN MARIJUANA²	99	93	12	7	97	93

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.71D Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
ILLCIT DRUGS²	0.22	0.22	0.61	0.67	0.23	0.22
Marijuana and Hashish	0.19	0.18	0.58	0.63	0.20	0.19
Cocaine	0.06	0.05	0.06	0.05	0.06	0.06
Crack	0.03	0.03	*	0.05	0.03	0.03
Heroin	0.02	0.02	0.06	0.15	0.02	0.02
Hallucinogens	0.05	0.05	0.43	0.04	0.05	0.05
LSD	0.01	0.01	0.15	*	0.01	0.01
PCP	0.01	0.00	0.15	0.01	0.01	0.00
Ecstasy	0.04	0.04	0.43	0.04	0.04	0.04
Inhalants	0.03	0.03	0.16	0.04	0.03	0.03
Nonmedical Use of Psychotherapeutics ^{3,4}	0.14	0.13	0.26	0.27	0.15	0.14
Pain Relievers	0.12	0.10	0.24	0.25	0.12	0.11
OxyContin [®]	0.03	0.03	0.08	0.05	0.03	0.03
Tranquilizers	0.08	0.08	0.23	0.06	0.09	0.08
Stimulants ⁴	0.06	0.05	0.16	0.08	0.06	0.05
Methamphetamine ⁴	0.04	0.03	0.16	*	0.04	0.03
Sedatives	0.02	0.02	0.15	0.01	0.02	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.16	0.14	0.48	0.29	0.16	0.15

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.71P Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
ILLCIT DRUGS²	0.6066		0.6135		0.6551	
Marijuana and Hashish	0.1972		0.8039		0.2159	
Cocaine	0.0844		0.9755		0.0939	
Crack	0.2839		*		0.2609	
Heroin	0.9098		0.3469		0.8234	
Hallucinogens	0.4937		0.1496		0.6693	
LSD	0.0688		*		0.1352	
PCP	0.1401		0.2763		0.2848	
Ecstasy	0.5507		0.1649		0.8001	
Inhalants	0.9302		0.2026		0.9258	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.2002		0.9650		0.2004	
Pain Relievers	0.0316		1.0000		0.0329	
OxyContin [®]	0.7910		0.6049		0.7354	
Tranquilizers	0.3483		0.0517		0.4335	
Stimulants ⁴	0.2189		0.6628		0.2805	
Methamphetamine ⁴	0.8753		*		0.9854	
Sedatives	0.3693		0.3568		0.4618	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.0995		0.4757		0.1103	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.72C Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOTAL	143	147	16	16	142	146
AGE						
15-17	29	30	3	3	28	29
18-25	59	66	8	11	69	81
26-44	117	116	13	11	116	114
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	136	135	15	13	135	134
White	120	116	8	12	120	115
Black or African American	57	59	13	6	54	59
American Indian or Alaska Native	6	9	*	*	6	9
Native Hawaiian or Other Pacific Islander	5	8	*	*	5	8
Asian	19	23	*	*	19	23
Two or More Races	14	14	*	*	14	14
Hispanic or Latino	48	58	5	8	48	57
TRIMESTER²						
First	N/A	N/A	14	12	N/A	N/A
Second	N/A	N/A	7	9	N/A	N/A
Third	N/A	N/A	4	4	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.72D Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOTAL	0.22	0.22	0.61	0.67	0.23	0.22
AGE						
15-17	0.42	0.43	3.13	4.72	0.43	0.44
18-25	0.36	0.39	0.86	1.21	0.37	0.41
26-44	0.29	0.28	0.86	0.77	0.30	0.29
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.25	0.24	0.71	0.72	0.26	0.25
White	0.29	0.29	0.54	0.88	0.31	0.30
Black or African American	0.59	0.62	3.37	1.73	0.59	0.64
American Indian or Alaska Native	1.97	2.04	*	*	2.03	2.10
Native Hawaiian or Other Pacific Islander	2.01	2.89	*	*	2.02	2.96
Asian	0.58	0.66	*	*	0.61	0.68
Two or More Races	1.83	1.46	*	*	1.92	1.52
Hispanic or Latino	0.44	0.47	1.07	1.76	0.46	0.49
TRIMESTER²						
First	N/A	N/A	1.70	1.69	N/A	N/A
Second	N/A	N/A	0.75	1.12	N/A	N/A
Third	N/A	N/A	0.50	0.48	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.72P Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOTAL	0.6066		0.6135		0.6551	
AGE						
15-17	0.2308		0.3634		0.1857	
18-25	0.1497		0.4542		0.2202	
26-44	0.5009		0.9682		0.5058	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.7238		0.9217		0.7162	
White	0.5477		0.4141		0.4608	
Black or African American	0.5983		0.3517		0.3985	
American Indian or Alaska Native	0.8878		*		0.7332	
Native Hawaiian or Other Pacific Islander	0.8302		*		0.8577	
Asian	0.4073		*		0.4254	
Two or More Races	0.2912		*		0.2247	
Hispanic or Latino	0.0118		0.3385		0.0170	
TRIMESTER²						
First	N/A		0.7713		N/A	
Second	N/A		0.3874		N/A	
Third	N/A		0.4942		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.73C Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOTAL	119	120	15	15	118	118
AGE						
15-17	25	26	3	3	25	26
18-25	54	62	8	10	62	74
26-44	93	93	12	10	93	91
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	113	112	14	13	113	111
White	98	97	8	11	98	96
Black or African American	49	52	12	6	46	52
American Indian or Alaska Native	5	5	*	*	5	5
Native Hawaiian or Other Pacific Islander	5	6	*	*	5	6
Asian	15	14	*	*	15	14
Two or More Races	14	13	*	*	14	13
Hispanic or Latino	36	42	5	7	35	41
TRIMESTER²						
First	N/A	N/A	13	12	N/A	N/A
Second	N/A	N/A	7	8	N/A	N/A
Third	N/A	N/A	4	3	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.73D Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOTAL	0.19	0.18	0.58	0.63	0.20	0.19
AGE						
15-17	0.37	0.39	2.88	4.53	0.38	0.39
18-25	0.33	0.36	0.83	1.11	0.34	0.38
26-44	0.24	0.23	0.82	0.75	0.25	0.24
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.21	0.21	0.68	0.69	0.22	0.21
White	0.25	0.25	0.50	0.85	0.26	0.26
Black or African American	0.51	0.54	3.29	1.71	0.51	0.57
American Indian or Alaska Native	1.67	1.23	*	*	1.72	1.27
Native Hawaiian or Other Pacific Islander	1.88	2.27	*	*	1.89	2.32
Asian	0.46	0.39	*	*	0.48	0.40
Two or More Races	1.81	1.33	*	*	1.90	1.39
Hispanic or Latino	0.33	0.35	1.06	1.50	0.34	0.37
TRIMESTER²						
First	N/A	N/A	1.63	1.67	N/A	N/A
Second	N/A	N/A	0.74	0.95	N/A	N/A
Third	N/A	N/A	0.47	0.47	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.73P Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOTAL	0.1972		0.8039		0.2159	
AGE						
15-17	0.1524		0.3381		0.1221	
18-25	0.0224		0.8908		0.0305	
26-44	0.6384		0.9303		0.6271	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.5642		0.8427		0.5804	
White	0.5342		0.3629		0.6167	
Black or African American	0.5814		0.4247		0.4397	
American Indian or Alaska Native	0.4665		*		0.6906	
Native Hawaiian or Other Pacific Islander	0.9258		*		0.8815	
Asian	0.4300		*		0.4466	
Two or More Races	0.1020		*		0.0777	
Hispanic or Latino	0.0091		0.8399		0.0109	
TRIMESTER²						
First	N/A		0.6852		N/A	
Second	N/A		0.9077		N/A	
Third	N/A		0.6350		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.74C Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOBACCO PRODUCTS²	252	274	29	34	247	268
Cigarettes	246	261	29	33	241	255
Smokeless Tobacco	29	33	4	2	28	33
Cigars	73	90	5	8	73	89
Pipe Tobacco	22	25	4	1	22	25
ALCOHOL	419	436	26	24	415	433
Binge Alcohol Use ³	243	258	18	10	242	257
Heavy Alcohol Use ³	102	116	6	4	101	116

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.74D Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOBACCO PRODUCTS²	0.36	0.36	1.11	1.36	0.37	0.37
Cigarettes	0.35	0.35	1.11	1.35	0.36	0.36
Smokeless Tobacco	0.05	0.05	0.17	0.08	0.05	0.06
Cigars	0.12	0.14	0.21	0.34	0.12	0.14
Pipe Tobacco	0.04	0.04	0.17	0.04	0.04	0.04
ALCOHOL	0.40	0.40	0.98	1.02	0.42	0.41
Binge Alcohol Use ³	0.33	0.33	0.68	0.43	0.35	0.34
Heavy Alcohol Use ³	0.16	0.17	0.23	0.15	0.17	0.18

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.74P Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: *P* Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOBACCO PRODUCTS²	0.0008		0.1557		0.0003	
Cigarettes	0.0001		0.1603		0.0000	
Smokeless Tobacco	0.3313		0.6695		0.2922	
Cigars	0.0762		0.5609		0.1073	
Pipe Tobacco	0.0432		0.2455		0.0299	
ALCOHOL	0.1403		0.7023		0.2378	
Binge Alcohol Use ³	0.8914		0.0234		0.8410	
Heavy Alcohol Use ³	0.3529		0.1282		0.3532	

*Low precision; no estimate reported.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.75C Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOTAL	246	261	29	33	241	255
AGE						
15-17	29	26	4	*	29	26
18-25	70	74	15	15	88	90
26-44	215	230	25	29	211	226
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	230	242	28	33	225	236
White	212	217	25	30	207	211
Black or African American	85	85	13	10	84	84
American Indian or Alaska Native	22	17	*	*	22	17
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	34	40	*	*	34	40
Two or More Races	21	30	*	*	21	30
Hispanic or Latino	84	95	6	4	84	95
TRIMESTER²						
First	N/A	N/A	20	24	N/A	N/A
Second	N/A	N/A	17	17	N/A	N/A
Third	N/A	N/A	13	16	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.75D Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOTAL	0.35	0.35	1.11	1.35	0.36	0.36
AGE						
15-17	0.44	0.39	3.67	*	0.43	0.39
18-25	0.43	0.44	1.49	1.53	0.45	0.46
26-44	0.51	0.50	1.59	2.01	0.52	0.51
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.39	0.39	1.31	1.66	0.40	0.40
White	0.46	0.46	1.56	2.04	0.48	0.47
Black or African American	0.82	0.82	3.42	2.96	0.84	0.84
American Indian or Alaska Native	3.99	3.19	*	*	4.13	3.29
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	0.99	1.10	*	*	1.02	1.14
Two or More Races	2.34	2.22	*	*	2.39	2.27
Hispanic or Latino	0.70	0.75	1.27	0.84	0.74	0.78
TRIMESTER²						
First	N/A	N/A	2.34	3.03	N/A	N/A
Second	N/A	N/A	1.77	1.93	N/A	N/A
Third	N/A	N/A	1.60	2.03	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.75P Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOTAL	0.0001		0.1603		0.0000	
AGE						
15-17	0.0017		*		0.0025	
18-25	0.0007		0.9278		0.0006	
26-44	0.0076		0.1282		0.0030	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0009		0.0592		0.0003	
White	0.0064		0.1070		0.0029	
Black or African American	0.0612		0.8637		0.0499	
American Indian or Alaska Native	0.0247		*		0.0241	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.4876		*		0.4884	
Two or More Races	0.8997		*		0.7708	
Hispanic or Latino	0.1216		0.1709		0.1207	
TRIMESTER²						
First	N/A		0.3212		N/A	
Second	N/A		0.3082		N/A	
Third	N/A		0.6362		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.76C Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOTAL	419	436	26	24	415	433
AGE						
15-17	39	39	3	2	39	39
18-25	79	86	10	10	158	174
26-44	360	368	24	22	358	366
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	386	391	22	22	382	389
White	340	341	17	19	338	339
Black or African American	144	157	14	6	144	156
American Indian or Alaska Native	19	20	*	*	19	20
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	82	100	*	*	82	99
Two or More Races	28	39	*	*	27	39
Hispanic or Latino	146	167	13	10	146	167
TRIMESTER²						
First	N/A	N/A	21	17	N/A	N/A
Second	N/A	N/A	12	15	N/A	N/A
Third	N/A	N/A	8	7	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.76D Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOTAL	0.40	0.40	0.98	1.02	0.42	0.41
AGE						
15-17	0.54	0.54	3.22	2.89	0.55	0.55
18-25	0.48	0.51	1.02	1.10	0.49	0.51
26-44	0.57	0.57	1.50	1.56	0.59	0.59
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.41	0.41	1.02	1.14	0.43	0.43
White	0.45	0.47	1.06	1.39	0.46	0.48
Black or African American	1.05	1.05	3.59	1.85	1.10	1.08
American Indian or Alaska Native	3.93	3.54	*	*	4.07	3.64
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.95	2.00	*	*	2.00	2.07
Two or More Races	2.72	2.54	*	*	2.74	2.61
Hispanic or Latino	1.01	1.04	2.66	2.14	1.06	1.06
TRIMESTER²						
First	N/A	N/A	2.42	2.37	N/A	N/A
Second	N/A	N/A	1.22	1.75	N/A	N/A
Third	N/A	N/A	0.94	0.92	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.76P Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOTAL	0.1403		0.7023		0.2378	
AGE						
15-17	0.0646		0.0639		0.0896	
18-25	0.6406		0.2889		0.4162	
26-44	0.0501		0.7926		0.0886	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2064		0.5991		0.3649	
White	0.1852		0.2895		0.3954	
Black or African American	0.7533		0.0468		0.9217	
American Indian or Alaska Native	0.4631		*		0.4201	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.0169		*		0.0292	
Two or More Races	0.3031		*		0.3628	
Hispanic or Latino	0.1411		0.0923		0.1369	
TRIMESTER²						
First	N/A		0.5490		N/A	
Second	N/A		0.4044		N/A	
Third	N/A		0.7130		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.77C Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOTAL	243	258	18	10	242	257
AGE						
15-17	28	28	3	2	28	28
18-25	72	78	7	5	106	116
26-44	203	218	16	8	203	217
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	220	224	14	10	219	223
White	202	205	10	9	201	204
Black or African American	82	82	11	3	81	82
American Indian or Alaska Native	13	17	*	*	13	17
Native Hawaiian or Other Pacific Islander	*	13	*	*	*	13
Asian	37	51	*	*	37	51
Two or More Races	19	19	*	*	19	19
Hispanic or Latino	98	113	10	1	98	113
TRIMESTER²						
First	N/A	N/A	17	9	N/A	N/A
Second	N/A	N/A	3	4	N/A	N/A
Third	N/A	N/A	4	2	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.77D Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOTAL	0.33	0.33	0.68	0.43	0.35	0.34
AGE						
15-17	0.42	0.41	3.35	2.65	0.42	0.42
18-25	0.43	0.46	0.76	0.58	0.46	0.48
26-44	0.47	0.47	1.02	0.61	0.49	0.49
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.37	0.35	0.68	0.54	0.39	0.36
White	0.45	0.43	0.61	0.69	0.47	0.44
Black or African American	0.80	0.79	2.96	0.95	0.83	0.83
American Indian or Alaska Native	3.20	3.28	*	*	3.35	3.37
Native Hawaiian or Other Pacific Islander	*	4.46	*	*	*	4.57
Asian	1.11	1.32	*	*	1.16	1.36
Two or More Races	2.24	1.89	*	*	2.34	1.99
Hispanic or Latino	0.78	0.81	2.07	0.28	0.82	0.85
TRIMESTER²						
First	N/A	N/A	2.01	1.29	N/A	N/A
Second	N/A	N/A	0.34	0.53	N/A	N/A
Third	N/A	N/A	0.45	0.20	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.77P Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOTAL	0.8914		0.0234		0.8410	
AGE						
15-17	0.0013		0.0616		0.0031	
18-25	0.1131		0.0441		0.0909	
26-44	0.4721		0.2508		0.5300	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.7086		0.2272		0.6416	
White	0.6361		0.4915		0.5195	
Black or African American	0.7787		0.2450		0.6438	
American Indian or Alaska Native	0.9516		*		0.9106	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.0700		*		0.0763	
Two or More Races	0.2641		*		0.2136	
Hispanic or Latino	0.5111		0.0172		0.4873	
TRIMESTER²						
First	N/A		0.0303		N/A	
Second	N/A		0.5149		N/A	
Third	N/A		0.3701		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.78C Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2008-2009 and 2010-2011

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOTAL	102	116	6	4	101	116
AGE						
15-17	14	12	1	1	13	12
18-25	45	48	4	3	50	56
26-44	87	96	4	*	87	96
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	97	107	4	3	96	107
White	92	97	4	3	92	97
Black or African American	29	36	2	1	28	36
American Indian or Alaska Native	3	12	*	*	3	12
Native Hawaiian or Other Pacific Islander	3	*	*	*	3	*
Asian	9	15	*	*	9	15
Two or More Races	7	7	*	*	7	7
Hispanic or Latino	32	39	4	1	31	39
TRIMESTER²						
First	N/A	N/A	6	2	N/A	N/A
Second	N/A	N/A	*	3	N/A	N/A
Third	N/A	N/A	0	*	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.78D Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2008-2009 and 2010-2011

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOTAL	0.16	0.17	0.23	0.15	0.17	0.18
AGE						
15-17	0.21	0.18	1.60	1.09	0.21	0.19
18-25	0.27	0.28	0.38	0.39	0.28	0.30
26-44	0.22	0.24	0.29	*	0.23	0.25
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.18	0.20	0.21	0.18	0.19	0.20
White	0.23	0.24	0.25	0.22	0.24	0.25
Black or African American	0.32	0.40	0.51	0.39	0.33	0.42
American Indian or Alaska Native	0.90	2.82	*	*	0.95	2.91
Native Hawaiian or Other Pacific Islander	1.30	*	*	*	1.30	*
Asian	0.28	0.40	*	*	0.29	0.41
Two or More Races	0.89	0.75	*	*	0.94	0.79
Hispanic or Latino	0.29	0.34	0.85	0.28	0.30	0.35
TRIMESTER²						
First	N/A	N/A	0.75	0.29	N/A	N/A
Second	N/A	N/A	*	0.35	N/A	N/A
Third	N/A	N/A	0.04	*	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2008-2009 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.

Table 6.78P Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2010-2011 Versus 2008-2009

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2008-2009	2010-2011	2008-2009	2010-2011	2008-2009	2010-2011
TOTAL	0.3529		0.1282		0.3532	
AGE						
15-17	0.0028		0.1647		0.0060	
18-25	0.1570		0.8040		0.1481	
26-44	0.9603		*		0.9336	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.2467		0.2106		0.2435	
White	0.1859		0.1800		0.1803	
Black or African American	0.5980		0.7630		0.6198	
American Indian or Alaska Native	0.0064		*		0.0069	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.3624		*		0.3720	
Two or More Races	0.8641		*		0.9093	
Hispanic or Latino	0.4401		0.3864		0.4211	
TRIMESTER²						
First	N/A		0.0177		N/A	
Second	N/A		*		N/A	
Third	N/A		*		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Some 2008-2009 estimates used in the comparisons may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2011.