

Table 7.22A GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	1,635	1,326	1,198	1,443	1,698	1,389
AGE GROUP						
12 to 17	40 ^a	21	21	30	21	17
18 to 25	475 ^b	307 ^b	202 ^a	255 ^b	217 ^a	142
26 or Older	1,120	998	976	1,158	1,459	1,230
GENDER						
Male	986	864	765	965	1,211	976
Female	650 ^b	462	433	479	486	413
PAST YEAR USE						
TOTAL	130	68	100	99	170	108
AGE GROUP						
12 to 17	16	7	8	14	3	11
18 to 25	46	42	22	31	30	23
26 or Older	68	19	70	55	137	74
GENDER						
Male	105	55	81	78	123	96
Female	26	14	19	21	47	12
PAST MONTH USE						
TOTAL	28	24	28	18	70	43
AGE GROUP						
12 to 17	5	3	1	5	*	1
18 to 25	6	21 ^a	5	4	6	3
26 or Older	17	*	22	*	63	39
GENDER						
Male	24	22	22	8	40	43
Female	*	*	*	9	30	*

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.22B GHB Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	0.7	0.5	0.5	0.6	0.7	0.5
AGE GROUP						
12 to 17	0.2 ^a	0.1	0.1	0.1	0.1	0.1
18 to 25	1.5 ^b	0.9 ^b	0.6 ^a	0.8 ^b	0.6 ^a	0.4
26 or Older	0.6	0.5	0.5	0.6	0.7	0.6
GENDER						
Male	0.8	0.7	0.6	0.8	1.0	0.8
Female	0.5 ^b	0.4	0.3	0.4	0.4	0.3
PAST YEAR USE						
TOTAL	0.1	0.0	0.0	0.0	0.1	0.0
AGE GROUP						
12 to 17	0.1	0.0	0.0	0.1	0.0	0.0
18 to 25	0.1	0.1	0.1	0.1	0.1	0.1
26 or Older	0.0	0.0	0.0	0.0	0.1	0.0
GENDER						
Male	0.1	0.0	0.1	0.1	0.1	0.1
Female	0.0	0.0	0.0	0.0	0.0	0.0
PAST MONTH USE						
TOTAL	0.0	0.0	0.0	0.0	0.0	0.0
AGE GROUP						
12 to 17	0.0	0.0	0.0	0.0	*	0.0
18 to 25	0.0	0.1 ^a	0.0	0.0	0.0	0.0
26 or Older	0.0	*	0.0	*	0.0	0.0
GENDER						
Male	0.0	0.0	0.0	0.0	0.0	0.0
Female	*	*	*	0.0	0.0	*

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: GHB is also known as "G," "Georgia Home Boy," "Grievous Bodily Harm," "Liquid G," and gamma hydroxybutyrate.

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

**Table 7.23A Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:
Numbers in Thousands, 2006-2011**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	4,704 ^b	4,951 ^b	5,366 ^b	6,751 ^b	7,936	8,198
AGE GROUP						
12 to 17	701	673	613	631	664	621
18 to 25	2,778 ^b	2,868 ^b	3,111 ^b	3,701 ^a	4,038	4,079
26 or Older	1,224 ^b	1,410 ^b	1,642 ^b	2,419 ^b	3,234	3,498
GENDER						
Male	2,662 ^b	2,798 ^b	2,959 ^b	4,012	4,707	4,327
Female	2,042 ^b	2,153 ^b	2,407 ^b	2,739 ^b	3,228 ^b	3,870
PAST YEAR USE						
TOTAL	1,991 ^b	2,003 ^b	2,099 ^b	2,779	2,904	3,076
AGE GROUP						
12 to 17	434	395	382	398	414	409
18 to 25	1,212 ^b	1,216 ^b	1,218 ^b	1,608	1,774	1,766
26 or Older	345 ^b	391 ^b	499 ^b	773	716	902
GENDER						
Male	1,055 ^b	1,157 ^b	1,155 ^b	1,649	1,661	1,684
Female	937 ^b	846 ^b	944 ^b	1,129 ^a	1,243	1,392
PAST MONTH USE						
TOTAL	551 ^b	529 ^b	613 ^b	858	921	836
AGE GROUP						
12 to 17	119	134	102	118	126	122
18 to 25	339 ^b	344 ^b	346 ^b	466	501	490
26 or Older	94 ^a	51 ^b	165	274	293	224
GENDER						
Male	299 ^b	300 ^b	344 ^b	512	547	502
Female	252	229 ^a	269	346	373	334

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.23B Nonmedical Use of Adderall® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	1.9 ^b	2.0 ^b	2.1 ^b	2.7 ^b	3.1	3.2
AGE GROUP						
12 to 17	2.8	2.7	2.5	2.6	2.7	2.5
18 to 25	8.5 ^b	8.8 ^b	9.4 ^b	11.0	11.9	11.9
26 or Older	0.7 ^b	0.7 ^b	0.9 ^b	1.2 ^b	1.7	1.8
GENDER						
Male	2.2 ^b	2.3 ^b	2.4 ^b	3.3	3.8	3.5
Female	1.6 ^b	1.7 ^b	1.9 ^b	2.1 ^b	2.5 ^b	2.9
PAST YEAR USE						
TOTAL	0.8 ^b	0.8 ^b	0.8 ^b	1.1	1.1	1.2
AGE GROUP						
12 to 17	1.7	1.6	1.5	1.6	1.7	1.6
18 to 25	3.7 ^b	3.7 ^b	3.7 ^b	4.8	5.2	5.1
26 or Older	0.2 ^b	0.2 ^b	0.3 ^b	0.4	0.4	0.5
GENDER						
Male	0.9 ^b	1.0 ^b	1.0 ^b	1.3	1.3	1.4
Female	0.7 ^b	0.7 ^b	0.7 ^b	0.9 ^a	1.0	1.0
PAST MONTH USE						
TOTAL	0.2 ^b	0.2 ^b	0.2 ^a	0.3	0.4	0.3
AGE GROUP						
12 to 17	0.5	0.5	0.4	0.5	0.5	0.5
18 to 25	1.0 ^b	1.1 ^a	1.1 ^a	1.4	1.5	1.4
26 or Older	0.0 ^a	0.0 ^b	0.1	0.1	0.2	0.1
GENDER						
Male	0.3 ^b	0.2 ^b	0.3 ^a	0.4	0.4	0.4
Female	0.2	0.2	0.2	0.3	0.3	0.3

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

**Table 7.24A Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender:
Numbers in Thousands, 2006-2011**

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	4,440 ^b	5,197 ^b	5,525	6,231	6,444	6,123
AGE GROUP						
12 to 17	315	297	268	287	231	273
18 to 25	1,187	1,087	1,182	1,284 ^a	1,288 ^a	1,113
26 or Older	2,939 ^b	3,813 ^b	4,075 ^a	4,660	4,925	4,738
GENDER						
Male	2,141 ^b	2,366	2,345	2,932	3,227	2,703
Female	2,299 ^b	2,831 ^a	3,180	3,299	3,217	3,421
PAST YEAR USE						
TOTAL	2,034	2,048	2,065	2,163	2,264	1,904
AGE GROUP						
12 to 17	174	152	139	142	123	150
18 to 25	536 ^b	476	480 ^a	530 ^b	453	382
26 or Older	1,324	1,420	1,446	1,492	1,688	1,371
GENDER						
Male	997	995	940	1,089 ^a	1,160 ^a	789
Female	1,037	1,053	1,125	1,074	1,104	1,115
PAST MONTH USE						
TOTAL	544	377	442	613	815 ^a	486
AGE GROUP						
12 to 17	46	30	28	35	35	35
18 to 25	98	108	133	132	109	92
26 or Older	400	238	281	446	671 ^a	359
GENDER						
Male	329	135	206	308	460 ^a	205
Female	215	241	236	305	355	281

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.24B Nonmedical Use of Ambien® in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	1.8 ^b	2.1 ^a	2.2	2.5	2.5	2.4
AGE GROUP						
12 to 17	1.2	1.2	1.1	1.2	0.9	1.1
18 to 25	3.6	3.3	3.6	3.8 ^b	3.8 ^a	3.2
26 or Older	1.6 ^b	2.0 ^a	2.1	2.4	2.5	2.4
GENDER						
Male	1.8 ^a	2.0	1.9	2.4	2.6 ^a	2.2
Female	1.8 ^b	2.2	2.5	2.5	2.5	2.6
PAST YEAR USE						
TOTAL	0.8	0.8	0.8	0.9	0.9	0.7
AGE GROUP						
12 to 17	0.7	0.6	0.6	0.6	0.5	0.6
18 to 25	1.6 ^b	1.5 ^a	1.5 ^a	1.6 ^b	1.3	1.1
26 or Older	0.7	0.7	0.8	0.8	0.9	0.7
GENDER						
Male	0.8	0.8	0.8	0.9 ^a	0.9 ^a	0.6
Female	0.8	0.8	0.9	0.8	0.8	0.8
PAST MONTH USE						
TOTAL	0.2	0.2	0.2	0.2	0.3 ^a	0.2
AGE GROUP						
12 to 17	0.2	0.1	0.1	0.1	0.1	0.1
18 to 25	0.3	0.3	0.4	0.4	0.3	0.3
26 or Older	0.2	0.1	0.1	0.2	0.3 ^a	0.2
GENDER						
Male	0.3	0.1	0.2	0.3	0.4 ^a	0.2
Female	0.2	0.2	0.2	0.2	0.3	0.2

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.25A Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	7,770	7,108	7,532	7,419	8,044	7,678
AGE GROUP						
12 to 17	932 ^b	855 ^b	795	750	760	704
18 to 25	2,134	2,040 ^a	2,183	2,325	2,324	2,270
26 or Older	4,705	4,213	4,554	4,344	4,959	4,704
GENDER						
Male	4,879	4,211	4,428	4,262	5,030	4,696
Female	2,891	2,898	3,103	3,157	3,014	2,982
PAST YEAR USE						
TOTAL	1,582	1,354	1,434	1,470	1,781	1,577
AGE GROUP						
12 to 17	486 ^b	433 ^b	404 ^a	373	374	332
18 to 25	511	482	462	485	489	519
26 or Older	586	439 ^a	568	611	918	726
GENDER						
Male	887	690	779	727	1,102	807
Female	696	664	655	743	679	771
PAST MONTH USE						
TOTAL	423	403	404	412	467	395
AGE GROUP						
12 to 17	123 ^b	125 ^a	104	108	94	77
18 to 25	107	130	107	124	127	94
26 or Older	193	147	192	181	246	224
GENDER						
Male	263	205	220	195	248	189
Female	160	197	184	217	219	206

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.25B Use of Non-Prescription Cough or Cold Medicine to Get High in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	3.2	2.9	3.0	2.9	3.2	3.0
AGE GROUP						
12 to 17	3.7 ^b	3.4 ^a	3.2	3.0	3.1	2.8
18 to 25	6.5	6.2	6.6	6.9	6.8	6.6
26 or Older	2.5	2.2	2.4	2.2	2.5	2.4
GENDER						
Male	4.1	3.5	3.7	3.5	4.1	3.8
Female	2.3	2.3	2.4	2.4	2.3	2.2
PAST YEAR USE						
TOTAL	0.6	0.5	0.6	0.6	0.7	0.6
AGE GROUP						
12 to 17	1.9 ^b	1.7 ^a	1.6 ^a	1.5	1.5	1.3
18 to 25	1.6	1.5	1.4	1.4	1.4	1.5
26 or Older	0.3	0.2 ^a	0.3	0.3	0.5	0.4
GENDER						
Male	0.7	0.6	0.6	0.6	0.9	0.6
Female	0.5	0.5	0.5	0.6	0.5	0.6
PAST MONTH USE						
TOTAL	0.2	0.2	0.2	0.2	0.2	0.2
AGE GROUP						
12 to 17	0.5 ^a	0.5 ^a	0.4	0.4	0.4	0.3
18 to 25	0.3	0.4	0.3	0.4	0.4	0.3
26 or Older	0.1	0.1	0.1	0.1	0.1	0.1
GENDER						
Male	0.2	0.2	0.2	0.2	0.2	0.2
Female	0.1	0.2	0.1	0.2	0.2	0.2

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.26A Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	2,283	2,191	2,148	2,309	2,411	2,379
AGE GROUP						
12 to 17	83	71	74	78	80	88
18 to 25	919 ^b	796 ^b	636 ^a	595	555	503
26 or Older	1,281 ^a	1,324 ^a	1,439	1,637	1,776	1,788
GENDER						
Male	1,468	1,436	1,354	1,547	1,747	1,552
Female	814	755	794	762	663	827
PAST YEAR USE						
TOTAL	203 ^a	212 ^a	150 ^b	202 ^a	246	336
AGE GROUP						
12 to 17	33	32	34	35	51	54
18 to 25	74 ^b	122	70 ^b	92 ^a	136	152
26 or Older	95	58	46	75	59	130
GENDER						
Male	141	143	105 ^a	115 ^a	175	224
Female	62	69	45 ^a	88	71	112
PAST MONTH USE						
TOTAL	53	39	27 ^a	86	61	82
AGE GROUP						
12 to 17	11	12	10	17	14	25
18 to 25	14	17	17	26	22	29
26 or Older	28	10	*	43	24	28
GENDER						
Male	34	27	16	42	39	38
Female	19	12	10	43	21	44

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Ketamine is also known as "Special K" or "Super K."

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.26B Ketamine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	0.9	0.9	0.9	0.9	1.0	0.9
AGE GROUP						
12 to 17	0.3	0.3	0.3	0.3	0.3	0.4
18 to 25	2.8 ^b	2.4 ^b	1.9 ^b	1.8	1.6	1.5
26 or Older	0.7 ^a	0.7 ^a	0.7	0.8	0.9	0.9
GENDER						
Male	1.2	1.2	1.1	1.3	1.4	1.2
Female	0.6	0.6	0.6	0.6	0.5	0.6
PAST YEAR USE						
TOTAL	0.1	0.1 ^a	0.1 ^b	0.1 ^a	0.1	0.1
AGE GROUP						
12 to 17	0.1	0.1	0.1	0.1	0.2	0.2
18 to 25	0.2 ^a	0.4	0.2 ^b	0.3	0.4	0.4
26 or Older	0.1	0.0	0.0	0.0	0.0	0.1
GENDER						
Male	0.1	0.1	0.1 ^a	0.1 ^a	0.1	0.2
Female	0.0	0.1	0.0 ^a	0.1	0.1	0.1
PAST MONTH USE						
TOTAL	0.0	0.0	0.0 ^a	0.0	0.0	0.0
AGE GROUP						
12 to 17	0.0	0.0	0.0	0.1	0.1	0.1
18 to 25	0.0	0.1	0.1	0.1	0.1	0.1
26 or Older	0.0	0.0	*	0.0	0.0	0.0
GENDER						
Male	0.0	0.0	0.0	0.0	0.0	0.0
Female	0.0	0.0	0.0	0.0	0.0	0.0

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: Ketamine is also known as "Special K" or "Super K."

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.27A DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	688 ^b	678 ^b	731 ^b	885	1,212	1,082
AGE GROUP						
12 to 17	45	44	49	58	61	65
18 to 25	307 ^b	270 ^b	280 ^b	367 ^a	444	483
26 or Older	336 ^a	363	402	460	707	535
GENDER						
Male	508 ^b	521 ^b	474 ^b	644	883	825
Female	180	157	258	241	329	257
PAST YEAR USE						
TOTAL	104 ^b	82 ^b	150 ^b	194	234	265
AGE GROUP						
12 to 17	16 ^a	20	29	38	38	36
18 to 25	64 ^b	52 ^b	73 ^b	111 ^a	118 ^a	183
26 or Older	24	10	47	46	79	47
GENDER						
Male	83 ^b	63 ^b	111 ^b	117 ^b	161	213
Female	21	19	39	77	73	53
PAST MONTH USE						
TOTAL	20	*	46	57	81	47
AGE GROUP						
12 to 17	*	*	8	11	17	7
18 to 25	15	10	19	12	21	25
26 or Older	*	*	19	34	43	14
GENDER						
Male	13	7 ^a	40	22	50	27
Female	7	*	*	35	31	20

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: DMT is also known as dimethyltryptamine.

NOTE: AMT is also known as alpha-methyltryptamine.

NOTE: 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy".

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.27B DMT, AMT, or 5-MeO-DIPT ("Foxy") Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	0.3 ^b	0.3 ^b	0.3 ^b	0.4	0.5	0.4
AGE GROUP						
12 to 17	0.2	0.2	0.2	0.2	0.3	0.3
18 to 25	0.9 ^b	0.8 ^b	0.8 ^b	1.1 ^a	1.3	1.4
26 or Older	0.2	0.2	0.2	0.2	0.4	0.3
GENDER						
Male	0.4 ^b	0.4 ^b	0.4 ^b	0.5	0.7	0.7
Female	0.1	0.1	0.2	0.2	0.3	0.2
PAST YEAR USE						
TOTAL	0.0 ^b	0.0 ^b	0.1 ^b	0.1	0.1	0.1
AGE GROUP						
12 to 17	0.1 ^a	0.1	0.1	0.2	0.2	0.1
18 to 25	0.2 ^b	0.2 ^b	0.2 ^b	0.3 ^a	0.3 ^a	0.5
26 or Older	0.0	0.0	0.0	0.0	0.0	0.0
GENDER						
Male	0.1 ^b	0.1 ^b	0.1 ^b	0.1 ^b	0.1	0.2
Female	0.0	0.0	0.0	0.1	0.1	0.0
PAST MONTH USE						
TOTAL	0.0	*	0.0	0.0	0.0	0.0
AGE GROUP						
12 to 17	*	*	0.0	0.0	0.1	0.0
18 to 25	0.0	0.0	0.1	0.0	0.1	0.1
26 or Older	*	*	0.0	0.0	0.0	0.0
GENDER						
Male	0.0	0.0 ^a	0.0	0.0	0.0	0.0
Female	0.0	*	*	0.0	0.0	0.0

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: DMT is also known as dimethyltryptamine.

NOTE: AMT is also known as alpha-methyltryptamine.

NOTE: 5-MeO-DIPT is also known as N, N-diisopropyl-5-methoxytryptamine and "Foxy".

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.28A Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Numbers in Thousands, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	1,802 ^b	2,522 ^b	3,201 ^b	4,406 ^b	5,235	5,285
AGE GROUP						
12 to 17	229 ^b	298 ^a	394	418	399	375
18 to 25	1,188 ^b	1,599 ^b	1,962 ^b	2,698 ^a	2,999	3,015
26 or Older	386 ^b	624 ^b	846 ^b	1,289 ^b	1,837	1,895
GENDER						
Male	1,436 ^b	1,873 ^b	2,296 ^b	3,391	3,907	3,733
Female	366 ^b	649 ^b	905 ^b	1,015 ^b	1,329	1,551
PAST YEAR USE						
TOTAL	761	1,007	1,213 ^b	1,283 ^b	1,191 ^b	871
AGE GROUP						
12 to 17	149 ^a	192	264	244	229	208
18 to 25	552	656 ^a	738 ^b	784 ^b	651 ^a	522
26 or Older	60 ^a	158	211	255 ^a	311 ^a	141
GENDER						
Male	600	737	877 ^b	965 ^b	929 ^b	629
Female	161 ^a	270	337 ^a	318	262	242
PAST MONTH USE						
TOTAL	136	200 ^a	198 ^a	194	162	136
AGE GROUP						
12 to 17	28	42	59	50	44	37
18 to 25	104	134	129	95	93	85
26 or Older	*	24	10	49	25	13
GENDER						
Male	112	151 ^a	160 ^a	134	134	95
Female	24	49	37	60	28	41

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.

Table 7.28B Salvia divinorum Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Age Group and Gender: Percentages, 2006-2011

Drug Use/Demographic Characteristic	2006	2007	2008	2009	2010	2011
LIFETIME USE						
TOTAL	0.7 ^b	1.0 ^b	1.3 ^b	1.7 ^b	2.1	2.1
AGE GROUP						
12 to 17	0.9 ^b	1.2 ^a	1.6	1.7	1.6	1.5
18 to 25	3.6 ^b	4.9 ^b	6.0 ^b	8.0 ^a	8.8	8.8
26 or Older	0.2 ^b	0.3 ^b	0.4 ^b	0.7 ^b	0.9	1.0
GENDER						
Male	1.2 ^b	1.6 ^b	1.9 ^b	2.8	3.2	3.0
Female	0.3 ^b	0.5 ^b	0.7 ^b	0.8 ^b	1.0	1.2
PAST YEAR USE						
TOTAL	0.3	0.4 ^a	0.5 ^b	0.5 ^b	0.5 ^b	0.3
AGE GROUP						
12 to 17	0.6 ^a	0.8	1.1	1.0	0.9	0.8
18 to 25	1.7	2.0 ^b	2.2 ^b	2.3 ^b	1.9 ^a	1.5
26 or Older	0.0 ^a	0.1	0.1	0.1 ^a	0.2 ^b	0.1
GENDER						
Male	0.5	0.6	0.7 ^b	0.8 ^b	0.8 ^b	0.5
Female	0.1 ^a	0.2	0.3 ^a	0.2	0.2	0.2
PAST MONTH USE						
TOTAL	0.1	0.1 ^a	0.1 ^a	0.1	0.1	0.1
AGE GROUP						
12 to 17	0.1	0.2	0.2	0.2	0.2	0.1
18 to 25	0.3	0.4 ^a	0.4	0.3	0.3	0.2
26 or Older	*	0.0	0.0	0.0	0.0	0.0
GENDER						
Male	0.1	0.1 ^a	0.1 ^a	0.1	0.1	0.1
Female	0.0	0.0	0.0	0.0	0.0	0.0

*Low precision, no estimate reported.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

^a Difference between estimate and 2011 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2011 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2006-2011.