

**Table 1.11A Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	124,808	127,458	41,479	41,591	23,863	24,573
12	327	299	204	191	100	87
13	518	473	355 <sup>a</sup>	275	184 <sup>b</sup>	119
14	802	802	557	567	285	291
15	1,134	1,060	811	776	399	373
16	1,464	1,439	1,154	1,127	620	646
17	1,800	1,721	1,387	1,350	789 <sup>a</sup>	680
18	2,322	2,258	1,805	1,765	1,061	993
19	2,221	2,120	1,625	1,591	1,034	948
20	2,513	2,441	1,829	1,705	1,079	1,036
21	2,598	2,524	1,665	1,646	946	1,022
22	2,710	2,633	1,674	1,558	1,004	980
23	2,665	2,708	1,409	1,455	793	847
24	2,505	2,646	1,339	1,460	782 <sup>a</sup>	921
25	2,444	2,496	1,205	1,268	683	742
26-29	10,587	10,631	4,529	4,808	2,506	2,743
30-34	11,853	12,425	4,248	4,692	2,605	2,957
35-39	10,781	10,149	3,048	2,826	1,718	1,780
40-44	10,998	11,565	2,775	2,855	1,471	1,649
45-49	12,626	11,631	2,862	2,578	1,640	1,454
50-54	14,212	14,530	2,829	2,877	1,681	1,812
55-59	11,686	11,704	2,216	1,757	1,355	1,165
60-64	7,972	8,861	1,010	1,150	604	692
65 or Older	8,073 <sup>b</sup>	10,340	943	1,311	525	634

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.11B Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	48.0	48.6	16.0	15.9	9.2	9.4
12	8.1	8.0	5.0	5.1	2.5	2.3
13	12.9	11.2	8.8 <sup>b</sup>	6.5	4.6 <sup>b</sup>	2.8
14	19.1	18.3	13.3	12.9	6.8	6.6
15	27.1	25.6	19.3	18.8	9.5	9.0
16	34.8	33.7	27.5	26.4	14.7	15.1
17	42.1	41.7	32.4	32.7	18.4	16.5
18	49.3	47.7	38.4	37.3	22.5	21.0
19	52.2	50.9	38.2	38.2	24.3	22.8
20	58.3	56.8	42.4	39.7	25.0	24.1
21	58.8	59.6	37.7	38.8	21.4	24.1
22	60.9	58.8	37.6	34.8	22.6	21.9
23	61.0	61.9	32.2	33.2	18.1	19.4
24	61.3	60.2	32.8	33.2	19.2	21.0
25	61.2	60.7	30.2	30.9	17.1	18.1
26-29	61.7	61.8	26.4	27.9	14.6	15.9
30-34	60.0	61.7	21.5	23.3	13.2	14.7
35-39	55.5	53.3	15.7	14.8	8.8	9.3
40-44	54.5	53.9	13.8	13.3	7.3	7.7
45-49	59.0	58.2	13.4	12.9	7.7	7.3
50-54	60.7	63.0	12.1	12.5	7.2	7.9
55-59	56.8	57.0	10.8	8.6	6.6	5.7
60-64	47.6	49.8	6.0	6.5	3.6	3.9
65 or Older	19.3 <sup>b</sup>	23.8	2.3	3.0	1.3	1.5

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.12A Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	111,239 <sup>a</sup>	114,712	31,513	32,952	18,855	19,810
12	57	72	44	59	9	21
13	204	168	164	138	84	60
14	437	445	352	371	183	196
15	788	765	615	604	332	296
16	1,205	1,182	968	974	508	568
17	1,540	1,460	1,219	1,200	684	621
18	2,040	2,004	1,609	1,566	940	897
19	1,963	1,906	1,465	1,436	939	873
20	2,301	2,226	1,609	1,523	968	915
21	2,337	2,338	1,468	1,486	831	919
22	2,477	2,420	1,429	1,350	856	860
23	2,438	2,448	1,209	1,244	687	734
24	2,282	2,435	1,129	1,285	684	788
25	2,214	2,274	969	1,111	579	650
26-29	9,635	9,816	3,447	3,984	2,041	2,363
30-34	10,732	11,447	3,259 <sup>a</sup>	3,849	2,130	2,358
35-39	9,651	9,075	2,099	1,997	1,291	1,332
40-44	9,642	10,049	1,836	2,008	1,027	1,208
45-49	11,507	10,658	2,008	1,740	1,119	1,044
50-54	13,271	13,787	1,866	2,215	1,198	1,272
55-59	10,913	10,965	1,517	1,347	986	910
60-64	7,427	8,264	739	842	404	548
65 or Older	6,177 <sup>b</sup>	8,508	494	623	375	377

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.12B Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2012 and 2013**

<b>Age Category</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	42.8	43.7	12.1	12.6	7.3	7.5
12	1.4	1.9	1.1	1.6	0.2	0.6
13	5.1	4.0	4.1	3.3	2.1	1.4
14	10.4	10.2	8.4	8.5	4.4	4.5
15	18.8	18.5	14.7	14.6	7.9	7.1
16	28.7	27.7	23.0	22.8	12.1	13.3
17	36.0	35.4	28.5	29.1	16.0	15.0
18	43.3	42.3	34.2	33.1	20.0	18.9
19	46.1	45.7	34.4	34.5	22.1	20.9
20	53.4	51.8	37.3	35.5	22.5	21.3
21	52.9	55.2	33.2	35.1	18.8	21.7
22	55.7	54.1	32.1	30.2	19.3	19.2
23	55.8	55.9	27.7	28.4	15.7	16.8
24	55.9	55.4	27.6	29.3	16.7	18.0
25	55.4	55.3	24.3	27.0	14.5	15.8
26-29	56.2	57.0	20.1 <sup>a</sup>	23.1	11.9	13.7
30-34	54.3	56.9	16.5 <sup>a</sup>	19.1	10.8	11.7
35-39	49.6	47.6	10.8	10.5	6.6	7.0
40-44	47.8	46.8	9.1	9.4	5.1	5.6
45-49	53.7	53.3	9.4	8.7	5.2	5.2
50-54	56.7	59.7	8.0	9.6	5.1	5.5
55-59	53.0	53.4	7.4	6.6	4.8	4.4
60-64	44.4	46.5	4.4	4.7	2.4	3.1
65 or Older	14.8 <sup>b</sup>	19.6	1.2	1.4	0.9	0.9

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.13A Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	37,688	37,634	4,671	4,182	1,650	1,549
12	3	3	2	3	1	*
13	8	6	8	4	*	2
14	20	10	13	8	1	8
15	39	27	21	18	8	4
16	64	68	47	37	5	10
17	132	101	90	62	15	20
18	279 <sup>a</sup>	186	168	115	54	33
19	346	283	191	159	41	46
20	457	405	235	176	45	45
21	535	452	201	208	46	47
22	639	649	260	262	78	79
23	656	680	203	228	53	40
24	700	727	164	227	38	56
25	655	658	179	169	42	52
26-29	3,330	3,158	608	615	174	189
30-34	3,730	3,895	551	660	250	292
35-39	3,220	2,707	386	319	167	132
40-44	3,774	3,632	330 <sup>a</sup>	167	120	68
45-49	5,356 <sup>a</sup>	4,649	412 <sup>a</sup>	216	211	103
50-54	6,281	6,849	231	294	109	181
55-59	4,258	4,125	256	113	117	48
60-64	2,093	2,669	116	99	73	73
65 or Older	1,115 <sup>a</sup>	1,693	*	26	*	22

\*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.13B Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	14.5	14.3	1.8	1.6	0.6	0.6
12	0.1	0.1	0.1	0.1	0.0	*
13	0.2	0.1	0.2	0.1	*	0.0
14	0.5	0.2	0.3	0.2	0.0	0.2
15	0.9	0.6	0.5	0.4	0.2	0.1
16	1.5	1.6	1.1	0.9	0.1	0.2
17	3.1	2.5	2.1	1.5	0.4	0.5
18	5.9 <sup>b</sup>	3.9	3.6	2.4	1.2	0.7
19	8.1	6.8	4.5	3.8	1.0	1.1
20	10.6	9.4	5.5	4.1	1.0	1.0
21	12.1	10.7	4.6	4.9	1.1	1.1
22	14.4	14.5	5.8	5.8	1.8	1.8
23	15.0	15.5	4.6	5.2	1.2	0.9
24	17.1	16.5	4.0	5.2	0.9	1.3
25	16.4	16.0	4.5	4.1	1.1	1.3
26-29	19.4	18.4	3.5	3.6	1.0	1.1
30-34	18.9	19.3	2.8	3.3	1.3	1.4
35-39	16.6	14.2	2.0	1.7	0.9	0.7
40-44	18.7	16.9	1.6 <sup>a</sup>	0.8	0.6	0.3
45-49	25.0	23.3	1.9 <sup>a</sup>	1.1	1.0	0.5
50-54	26.8	29.7	1.0	1.3	0.5	0.8
55-59	20.7	20.1	1.2	0.5	0.6	0.2
60-64	12.5	15.0	0.7	0.6	0.4	0.4
65 or Older	2.7 <sup>a</sup>	3.9	*	0.1	*	0.1

\*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.14A Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	9,015	8,870	921 <sup>a</sup>	632	443	377
12	*	*	*	*	*	*
13	1	2	0	2	*	2
14	2	1	1	*	*	*
15	8	5	3	2	*	*
16	12	6	7	3	1	*
17	12	9	7	4	*	1
18	39	29	17	8	7	2
19	48	27	20	10	2	2
20	67	40	27 <sup>a</sup>	6	5	*
21	59	49	12	11	5	6
22	91 <sup>a</sup>	54	17	11	5	5
23	113	111	11	10	3	1
24	116	119	19	9	7	3
25	121	134	16	30	3	17
26-29	717	809	79	49	15	17
30-34	1,192	957	129	101	78	67
35-39	918	784	73	35	53	20
40-44	1,126	939	88	44	27	37
45-49	1,467	1,217	172 <sup>a</sup>	40	97	21
50-54	1,455	1,760	117	121	57	103
55-59	836	1,128	71	81	60	24
60-64	484	386	32	29	17	29
65 or Older	132	303	*	26	*	21

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.14B Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	3.5	3.4	0.4 <sup>a</sup>	0.2	0.2	0.1
12	*	*	*	*	*	*
13	0.0	0.0	0.0	0.0	*	0.0
14	0.0	0.0	0.0	*	*	*
15	0.2	0.1	0.1	0.1	*	*
16	0.3	0.1	0.2	0.1	0.0	*
17	0.3	0.2	0.2	0.1	*	0.0
18	0.8	0.6	0.4	0.2	0.2	0.0
19	1.1	0.7	0.5	0.2	0.0	0.1
20	1.6	0.9	0.6 <sup>a</sup>	0.1	0.1	*
21	1.3	1.2	0.3	0.3	0.1	0.2
22	2.0 <sup>a</sup>	1.2	0.4	0.3	0.1	0.1
23	2.6	2.5	0.3	0.2	0.1	0.0
24	2.8	2.7	0.5	0.2	0.2	0.1
25	3.0	3.3	0.4	0.7	0.1	0.4
26-29	4.2	4.7	0.5	0.3	0.1	0.1
30-34	6.0	4.8	0.7	0.5	0.4	0.3
35-39	4.7	4.1	0.4	0.2	0.3	0.1
40-44	5.6	4.4	0.4	0.2	0.1	0.2
45-49	6.8	6.1	0.8 <sup>a</sup>	0.2	0.5	0.1
50-54	6.2	7.6	0.5	0.5	0.2	0.4
55-59	4.1	5.5	0.3	0.4	0.3	0.1
60-64	2.9	2.2	0.2	0.2	0.1	0.2
65 or Older	0.3	0.7	*	0.1	*	0.0

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.



**Table 1.15A Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	37,908	39,736	4,306	4,430	1,134	1,333
12	14	25	8	16	6	3
13	35	34	25	19	13	8
14	89 <sup>a</sup>	54	61	38	17	18
15	127	97	87	61	24	19
16	220	200	171	140	45	32
17	339	293	200	208	54	74
18	551	486	344	305	137	123
19	586	564	375	309	105	93
20	721	687	371	313	78	65
21	776	785	277	357	71	65
22	909	908	273	333	75	88
23	858	936	224	303	69	62
24	857	975	188	251	34 <sup>a</sup>	79
25	845	783	184	167	31	42
26-29	3,833	3,863	545	473	110	147
30-34	4,653	4,903	351	444	85	90
35-39	3,895	3,741	249	218	56	117
40-44	3,173 <sup>a</sup>	3,778	82	153	21	21
45-49	3,885	3,395	112	57	49	12
50-54	4,332	5,215	35	87	35	77
55-59	3,822	3,804	16	8	4	*
60-64	2,229	2,542	69	13	*	*
65 or Older	1,159	1,671	57	157	13	97

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.15B Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	14.6	15.1	1.7	1.7	0.4	0.5
12	0.4	0.7	0.2	0.4	0.2	0.1
13	0.9	0.8	0.6	0.5	0.3	0.2
14	2.1 <sup>a</sup>	1.2	1.5	0.9	0.4	0.4
15	3.0	2.4	2.1	1.5	0.6	0.5
16	5.2	4.7	4.1	3.3	1.1	0.7
17	7.9	7.1	4.7	5.0	1.3	1.8
18	11.7	10.3	7.3	6.4	2.9	2.6
19	13.8	13.5	8.8	7.4	2.5	2.2
20	16.7	16.0	8.6	7.3	1.8	1.5
21	17.6	18.5	6.3 <sup>a</sup>	8.4	1.6	1.5
22	20.4	20.3	6.1	7.4	1.7	2.0
23	19.6	21.4	5.1 <sup>a</sup>	6.9	1.6	1.4
24	21.0	22.2	4.6	5.7	0.8 <sup>a</sup>	1.8
25	21.2	19.1	4.6	4.1	0.8	1.0
26-29	22.3	22.4	3.2	2.7	0.6	0.9
30-34	23.6	24.4	1.8	2.2	0.4	0.4
35-39	20.0	19.6	1.3	1.1	0.3	0.6
40-44	15.7	17.6	0.4	0.7	0.1	0.1
45-49	18.1	17.0	0.5	0.3	0.2	0.1
50-54	18.5 <sup>a</sup>	22.6	0.1	0.4	0.1	0.3
55-59	18.6	18.5	0.1	0.0	0.0	*
60-64	13.3	14.3	0.4	0.1	*	*
65 or Older	2.8	3.8	0.1	0.4	0.0	0.2

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.16A Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	21,095	21,068	1,693	1,533	493	496
12	172	139	73	71	31	27
13	222	197	104	74	43 <sup>a</sup>	20
14	316	265	142 <sup>a</sup>	90	38	18
15	324	287	108	107	18	31
16	296 <sup>a</sup>	215	125 <sup>b</sup>	63	27	12
17	283 <sup>a</sup>	224	95	76	31	13
18	358	280	107	74	28	16
19	312	277	68	66	16	15
20	325	309	64	78	16	20
21	350	341	88	84	35	16
22	399	376	56	54	12	17
23	386	367	30	48	8	6
24	374	386	15 <sup>a</sup>	45	1	7
25	413 <sup>b</sup>	277	45	30	7	10
26-29	1,891	1,809	121	113	39	22
30-34	2,735	2,768	66	115	18	62
35-39	2,180	2,108	94	43	14	19
40-44	2,275	2,299	90	77	5	51
45-49	2,097	2,167	117	74	43	55
50-54	1,951 <sup>b</sup>	2,888	39	50	26	31
55-59	1,878	1,391	37	93	37	19
60-64	762	971	9	7	*	7
65 or Older	795	727	*	3	*	3

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.16B Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	8.1	8.0	0.7	0.6	0.2	0.2
12	4.2	3.7	1.8	1.9	0.8	0.7
13	5.5	4.7	2.6	1.7	1.1 <sup>a</sup>	0.5
14	7.5	6.0	3.4 <sup>b</sup>	2.0	0.9 <sup>a</sup>	0.4
15	7.7	6.9	2.6	2.6	0.4	0.8
16	7.0 <sup>b</sup>	5.0	3.0 <sup>b</sup>	1.5	0.6	0.3
17	6.6	5.4	2.2	1.8	0.7	0.3
18	7.6	5.9	2.3	1.6	0.6	0.3
19	7.3	6.7	1.6	1.6	0.4	0.4
20	7.5	7.2	1.5	1.8	0.4	0.5
21	7.9	8.0	2.0	2.0	0.8	0.4
22	9.0	8.4	1.3	1.2	0.3	0.4
23	8.8	8.4	0.7	1.1	0.2	0.1
24	9.2	8.8	0.4 <sup>a</sup>	1.0	0.0	0.2
25	10.3 <sup>b</sup>	6.7	1.1	0.7	0.2	0.2
26-29	11.0	10.5	0.7	0.7	0.2	0.1
30-34	13.8	13.7	0.3	0.6	0.1	0.3
35-39	11.2	11.1	0.5	0.2	0.1	0.1
40-44	11.3	10.7	0.4	0.4	0.0	0.2
45-49	9.8	10.8	0.5	0.4	0.2	0.3
50-54	8.3 <sup>b</sup>	12.5	0.2	0.2	0.1	0.1
55-59	9.1	6.8	0.2	0.5	0.2	0.1
60-64	4.6	5.5	0.1	0.0	*	0.0
65 or Older	1.9	1.7	*	0.0	*	0.0

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.17A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	54,389	53,172	16,666 <sup>a</sup>	15,348	6,831	6,484
12	164	130	119	88	66	50
13	219	199	149 <sup>a</sup>	99	70	50
14	305	338	198	229	100	96
15	469 <sup>b</sup>	356	302	248	114	89
16	580	549	394	371	159	140
17	745 <sup>a</sup>	615	479	406	185 <sup>a</sup>	124
18	1,037	904	634	527	224	182
19	990	871	556	490	217	165
20	1,213	1,117	701	584	252	236
21	1,211	1,178	563	589	204	227
22	1,405	1,289	743 <sup>a</sup>	603	297	234
23	1,328	1,340	546	535	235	243
24	1,266	1,324	525	507	205	215
25	1,259	1,216	460	409	198	159
26-29	5,650	5,448	1,951	1,877	730	745
30-34	5,814	6,014	1,807	1,966	773	913
35-39	4,791	4,593	1,304	1,185	559	493
40-44	4,623	4,765	1,296	1,113	516	590
45-49	5,329 <sup>b</sup>	4,319	1,074	1,026	445	419
50-54	5,912	6,168	1,289	967	564	482
55-59	4,774	3,999	832	542	380	247
60-64	2,654	2,792	300	324	200	95
65 or Older	2,652 <sup>a</sup>	3,648	442	662	137	290

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.17B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	20.9	20.3	6.4 <sup>a</sup>	5.8	2.6	2.5
12	4.0	3.5	3.0	2.3	1.6	1.3
13	5.5	4.7	3.7 <sup>b</sup>	2.3	1.7	1.2
14	7.2	7.7	4.7	5.2	2.4	2.2
15	11.2 <sup>b</sup>	8.6	7.2	6.0	2.7	2.1
16	13.8	12.8	9.4	8.7	3.8	3.3
17	17.4 <sup>a</sup>	14.9	11.2	9.8	4.3 <sup>a</sup>	3.0
18	22.0 <sup>a</sup>	19.1	13.5 <sup>a</sup>	11.1	4.8	3.9
19	23.3	20.9	13.1	11.8	5.1	4.0
20	28.1	26.0	16.3	13.6	5.8	5.5
21	27.4	27.8	12.7	13.9	4.6	5.4
22	31.6	28.8	16.7 <sup>a</sup>	13.5	6.7	5.2
23	30.4	30.6	12.5	12.2	5.4	5.5
24	31.0	30.1	12.8	11.6	5.0	4.9
25	31.5	29.6	11.5	9.9	5.0	3.9
26-29	32.9	31.7	11.4	10.9	4.3	4.3
30-34	29.4	29.9	9.1	9.8	3.9	4.5
35-39	24.6	24.1	6.7	6.2	2.9	2.6
40-44	22.9	22.2	6.4	5.2	2.6	2.7
45-49	24.9 <sup>a</sup>	21.6	5.0	5.1	2.1	2.1
50-54	25.3	26.7	5.5	4.2	2.4	2.1
55-59	23.2 <sup>a</sup>	19.5	4.0	2.6	1.8	1.2
60-64	15.9	15.7	1.8	1.8	1.2	0.5
65 or Older	6.3 <sup>a</sup>	8.4	1.1	1.5	0.3	0.7

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.18A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	37,045	35,473	12,489 <sup>b</sup>	11,082	4,862	4,521
12	147 <sup>a</sup>	99	107 <sup>a</sup>	64	59	34
13	185	168	125	86	59	42
14	246	294	162	198	88	88
15	383 <sup>a</sup>	296	247	204	93	70
16	497	450	329	297	134	106
17	611 <sup>a</sup>	499	362	308	124 <sup>a</sup>	87
18	871 <sup>a</sup>	705	499 <sup>a</sup>	392	171 <sup>a</sup>	116
19	772	702	410	362	148	100
20	979	895	528	425	190	174
21	957	926	361	407	137	144
22	1,134	1,039	550	440	208	155
23	1,031	1,066	410	377	167	183
24	1,020	1,014	394	357	138	148
25	996	904	354	300	152	113
26-29	4,498	4,092	1,459	1,266	559	484
30-34	4,337	4,643	1,288	1,400	551	631
35-39	3,586	3,291	994	854	397	344
40-44	3,273	3,373	997	847	382	414
45-49	3,255 <sup>a</sup>	2,660	810	708	346	306
50-54	3,269	3,185	919	769	348	318
55-59	2,694 <sup>a</sup>	2,043	613	392	163	133
60-64	1,259	1,445	233	196	135	69
65 or Older	1,047 <sup>a</sup>	1,687	336	432	111	264

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.18B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2012 and 2013**

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOTAL</b>	14.2 <sup>a</sup>	13.5	4.8 <sup>b</sup>	4.2	1.9	1.7
12	3.6	2.6	2.7 <sup>a</sup>	1.7	1.5	0.9
13	4.6	4.0	3.1 <sup>a</sup>	2.0	1.5	1.0
14	5.9	6.7	3.9	4.5	2.1	2.0
15	9.1 <sup>a</sup>	7.2	5.9	4.9	2.2	1.7
16	11.8	10.5	7.8	6.9	3.2	2.5
17	14.3 <sup>a</sup>	12.1	8.5	7.5	2.9	2.1
18	18.5 <sup>b</sup>	14.9	10.6 <sup>a</sup>	8.3	3.6 <sup>a</sup>	2.4
19	18.1	16.8	9.6	8.7	3.5	2.4
20	22.7	20.8	12.3	9.9	4.4	4.0
21	21.7	21.8	8.2	9.6	3.1	3.4
22	25.5	23.2	12.4 <sup>a</sup>	9.8	4.7	3.5
23	23.6	24.3	9.4	8.6	3.8	4.2
24	25.0	23.1	9.6	8.1	3.4	3.4
25	24.9	22.0	8.9	7.3	3.8	2.7
26-29	26.2	23.8	8.5	7.4	3.3	2.8
30-34	22.0	23.1	6.5	7.0	2.8	3.1
35-39	18.4	17.3	5.1	4.5	2.0	1.8
40-44	16.2	15.7	4.9	3.9	1.9	1.9
45-49	15.2	13.3	3.8	3.5	1.6	1.5
50-54	14.0	13.8	3.9	3.3	1.5	1.4
55-59	13.1 <sup>a</sup>	9.9	3.0	1.9	0.8	0.6
60-64	7.5	8.1	1.4	1.1	0.8	0.4
65 or Older	2.5 <sup>a</sup>	3.9	0.8	1.0	0.3	0.6

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.