Table 1.74A Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2012 and 2013

Geographic Characteristic	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	9,015	8,870	921 <sup>a</sup>	632	443	377
GEOGRAPHIC DIVISION	9,013	0,070	921	032	443	311
Northeast	1,331	1,229	177	100	86	66
New England	314	412	27	29	12	23
Middle Atlantic	1,017	818	150	71	74	42
Midwest	1,883	1,814	183	178	84	98
East North Central	1,334	1,204	163	153	77	80
West North Central	549	610	20	25	7	18
South	3,410	3,421	477 <sup>a</sup>	254	241	146
South Atlantic	1,875	2,005	319	160	194	90
East South Central	544	579	47	44	37	33
West South Central	991	838	111	50	11	22
West	2,391	2,405	84	100	32	68
Mountain	839	781	54	50	26	36
Pacific	1,552	1,624	30	50	6	31
COUNTY TYPE						
Large Metro	4,605	4,435	504 <sup>a</sup>	252	276	189
Small Metro	3,132	2,783	294	252	121	123
250K – 1 Mil. Pop.	2,052	1,993	185	208	102	101
< 250K Pop.	1,079	791	110	44	19	22
Nonmetro	1,279 <sup>a</sup>	1,651	124	128	46	65
Urbanized	546	679	43	42	27	14
Less Urbanized	602	780	54	42	9	24
Completely Rural	131	193	27	44	10	28

<sup>\*</sup>Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

<sup>&</sup>lt;sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>&</sup>lt;sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Table 1.74B Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2012 and 2013

Geographic Characteristic	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	3.5	3.4	$0.4^{a}$	0.2	0.2	0.1
GEOGRAPHIC DIVISION						
Northeast	2.8	2.6	0.4	0.2	0.2	0.1
New England	2.5	3.3	0.2	0.2	0.1	0.2
Middle Atlantic	2.9	2.3	0.4	0.2	0.2	0.1
Midwest	3.4	3.2	0.3	0.3	0.2	0.2
East North Central	3.4	3.1	0.4	0.4	0.2	0.2
West North Central	3.2	3.5	0.1	0.1	0.0	0.1
South	3.5	3.5	$0.5^{\mathrm{a}}$	0.3	0.3	0.1
South Atlantic	3.7	3.9	0.6	0.3	0.4	0.2
East South Central	3.5	3.7	0.3	0.3	0.2	0.2
West South Central	3.3	2.7	0.4	0.2	0.0	0.1
West	3.9	3.9	0.1	0.2	0.1	0.1
Mountain	4.6	4.2	0.3	0.3	0.1	0.2
Pacific	3.7	3.8	0.1	0.1	0.0	0.1
COUNTY TYPE						
Large Metro	3.3	3.1	$0.4^{a}$	0.2	0.2	0.1
Small Metro	4.0	3.5	0.4	0.3	0.2	0.2
250K – 1 Mil. Pop.	3.9	3.7	0.3	0.4	0.2	0.2
< 250K Pop.	4.2	3.1	0.4	0.2	0.1	0.1
Nonmetro	$3.0^{a}$	3.9	0.3	0.3	0.1	0.2
Urbanized	3.0	3.6	0.2	0.2	0.1	0.1
Less Urbanized	3.2	4.1	0.3	0.2	0.0	0.1
Completely Rural	2.7	4.3	0.6	1.0	0.2	0.6

<sup>\*</sup>Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

<sup>&</sup>lt;sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>&</sup>lt;sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.