

Table 1.76A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2012 and 2013

Geographic Characteristic	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	21,095	21,068	1,693	1,533	493	496
GEOGRAPHIC DIVISION						
Northeast	3,892	4,135	266	247	43 ^a	97
New England	1,200	1,162	73	64	11	20
Middle Atlantic	2,692	2,973	193	183	32 ^a	77
Midwest	4,373	4,141	297	234	100	81
East North Central	3,182	2,902	221	181	71	71
West North Central	1,191	1,239	76	53	29	10
South	6,961	6,707	602	530	190	130
South Atlantic	4,039	3,748	350	320	100	71
East South Central	1,018	1,015	77	82	23	28
West South Central	1,905	1,945	175	128	68	31
West	5,867	6,085	528	522	160	187
Mountain	2,043	2,017	133	189	33	60
Pacific	3,824	4,068	395	332	126	127
COUNTY TYPE						
Large Metro	11,149	11,790	987	873	328	280
Small Metro	7,091	6,312	447	433	107	136
250K – 1 Mil. Pop.	4,719	4,526	313	294	67	76
< 250K Pop.	2,371 ^a	1,786	134	138	40	61
Nonmetro	2,855	2,966	259	227	58	80
Urbanized	1,520	1,513	157	121	21	48
Less Urbanized	1,115	1,229	74	89	25	20
Completely Rural	220	224	28	17	12	11

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.76B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2012 and 2013

Geographic Characteristic	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	8.1	8.0	0.7	0.6	0.2	0.2
GEOGRAPHIC DIVISION						
Northeast	8.3	8.7	0.6	0.5	0.1 ^a	0.2
New England	9.7	9.3	0.6	0.5	0.1	0.2
Middle Atlantic	7.7	8.5	0.6	0.5	0.1 ^a	0.2
Midwest	7.8	7.4	0.5	0.4	0.2	0.1
East North Central	8.2	7.4	0.6	0.5	0.2	0.2
West North Central	7.0	7.2	0.4	0.3	0.2	0.1
South	7.2	6.9	0.6	0.5	0.2	0.1
South Atlantic	7.9	7.3	0.7	0.6	0.2	0.1
East South Central	6.6	6.5	0.5	0.5	0.1	0.2
West South Central	6.3	6.4	0.6	0.4	0.2	0.1
West	9.7	9.9	0.9	0.9	0.3	0.3
Mountain	11.1	10.8	0.7	1.0	0.2	0.3
Pacific	9.1	9.6	0.9	0.8	0.3	0.3
COUNTY TYPE						
Large Metro	8.0	8.4	0.7	0.6	0.2	0.2
Small Metro	9.0 ^a	8.0	0.6	0.5	0.1	0.2
250K – 1 Mil. Pop.	8.9	8.5	0.6	0.6	0.1	0.1
< 250K Pop.	9.3 ^b	7.0	0.5	0.5	0.2	0.2
Nonmetro	6.8	7.0	0.6	0.5	0.1	0.2
Urbanized	8.2	8.0	0.8	0.6	0.1	0.3
Less Urbanized	5.9	6.5	0.4	0.5	0.1	0.1
Completely Rural	4.6	5.0	0.6	0.4	0.3	0.2

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.77A Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2012 and 2013

Geographic Characteristic	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	1,613 ^b	1,327	647 ^b	480	188 ^b	121
GEOGRAPHIC DIVISION						
Northeast	261 ^b	163	87	71	15	18
New England	58	36	22	14	3	5
Middle Atlantic	203 ^b	127	65	58	11	13
Midwest	347 ^b	251	139 ^a	98	42	25
East North Central	246 ^b	176	103	77	29	19
West North Central	101	75	35	21	13	6
South	596	573	253	212	83	62
South Atlantic	301	291	129	109	31	36
East South Central	101	91	39	29	16 ^a	4
West South Central	194	191	85	74	36	23
West	409	339	169 ^b	99	48 ^b	16
Mountain	126	110	46	40	15	7
Pacific	283	229	123 ^b	59	33 ^a	8
COUNTY TYPE						
Large Metro	865 ^b	699	355 ^b	256	106 ^b	52
Small Metro	483	429	187	150	48	45
250K – 1 Mil. Pop.	318	275	127	99	33	26
< 250K Pop.	164	154	60	51	16	19
Nonmetro	265 ^b	199	105 ^a	74	34	24
Urbanized	119 ^a	78	44	28	11	8
Less Urbanized	120	103	48	41	17	16
Completely Rural	27	18	13	5	6	0

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.77B Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2012 and 2013

Geographic Characteristic	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	6.5 ^b	5.3	2.6 ^b	1.9	0.8 ^b	0.5
GEOGRAPHIC DIVISION						
Northeast	6.1 ^b	3.9	2.1	1.7	0.3	0.4
New England	5.2	3.3	2.0	1.2	0.3	0.4
Middle Atlantic	6.5 ^b	4.1	2.1	1.9	0.4	0.4
Midwest	6.4 ^b	4.7	2.6 ^a	1.8	0.8	0.5
East North Central	6.5 ^b	4.7	2.7	2.1	0.8	0.5
West North Central	6.2	4.6	2.2	1.3	0.8	0.3
South	6.4	6.1	2.7	2.3	0.9	0.7
South Atlantic	6.5	6.2	2.8	2.3	0.7	0.8
East South Central	6.8	6.2	2.7	1.9	1.1 ^a	0.2
West South Central	6.1	5.9	2.7	2.3	1.1	0.7
West	6.8	5.7	2.8 ^b	1.7	0.8 ^b	0.3
Mountain	6.8	5.9	2.5	2.1	0.8	0.4
Pacific	6.9	5.6	3.0 ^b	1.4	0.8 ^a	0.2
COUNTY TYPE						
Large Metro	6.4 ^b	5.2	2.6 ^b	1.9	0.8 ^b	0.4
Small Metro	6.5	5.8	2.5	2.0	0.7	0.6
250K – 1 Mil. Pop.	6.2	5.4	2.5	1.9	0.6	0.5
< 250K Pop.	7.1	6.6	2.6	2.2	0.7	0.8
Nonmetro	6.8 ^b	5.1	2.7 ^a	1.9	0.9	0.6
Urbanized	6.5 ^a	4.7	2.4	1.7	0.6	0.5
Less Urbanized	7.1	5.8	2.9	2.3	1.0	0.9
Completely Rural	6.4	4.1	3.1	1.1	1.4	0.1

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.