

**Table 1.78A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	54,389	53,172	16,666 <sup>a</sup>	15,348	6,831	6,484
<b>GEOGRAPHIC DIVISION</b>						
Northeast	9,116	8,415	2,738	2,424	1,055	915
New England	2,599	2,486	686	641	277	197
Middle Atlantic	6,517	5,929	2,052	1,783	778	719
Midwest	11,045	10,627	3,372	3,002	1,302	1,275
East North Central	8,059 <sup>a</sup>	7,199	2,484	2,163	1,039	921
West North Central	2,986	3,428	887	839	263	354
South	19,039	19,315	6,075	5,874	2,661	2,528
South Atlantic	9,991	10,455	2,911	3,197	1,249	1,263
East South Central	2,954	2,908	1,058	906	492	456
West South Central	6,094	5,952	2,106	1,770	919	810
West	15,189	14,814	4,481	4,048	1,812	1,765
Mountain	4,630	4,450	1,354	1,272	540	464
Pacific	10,559	10,364	3,127	2,776	1,273	1,301
<b>COUNTY TYPE</b>						
Large Metro	30,081	28,830	9,421 <sup>a</sup>	8,342	3,871	3,526
Small Metro	16,870	16,499	4,990	4,722	2,121	2,104
250K – 1 Mil. Pop.	11,547	11,325	3,377	3,047	1,493	1,391
< 250K Pop.	5,324	5,174	1,612	1,675	628	713
Nonmetro	7,438	7,844	2,256	2,284	839	853
Urbanized	3,645	3,881	1,062	1,127	394	397
Less Urbanized	3,084	3,286	1,010	975	378	383
Completely Rural	709	677	184	182	66	74

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.78B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	20.9	20.3	6.4 <sup>a</sup>	5.8	2.6	2.5
<b>GEOGRAPHIC DIVISION</b>						
Northeast	19.3	17.8	5.8	5.1	2.2	1.9
New England	20.9	19.9	5.5	5.1	2.2	1.6
Middle Atlantic	18.7	17.0	5.9	5.1	2.2	2.1
Midwest	19.8	18.9	6.0	5.3	2.3	2.3
East North Central	20.8 <sup>b</sup>	18.5	6.4	5.5	2.7	2.4
West North Central	17.5	19.9	5.2	4.9	1.5	2.1
South	19.8	19.8	6.3	6.0	2.8	2.6
South Atlantic	19.6	20.3	5.7	6.2	2.5	2.5
East South Central	19.2	18.8	6.9	5.8	3.2	2.9
West South Central	20.2	19.5	7.0 <sup>a</sup>	5.8	3.1	2.6
West	25.1	24.2	7.4	6.6	3.0	2.9
Mountain	25.1	23.8	7.3	6.8	2.9	2.5
Pacific	25.0	24.3	7.4	6.5	3.0	3.1
<b>COUNTY TYPE</b>						
Large Metro	21.6	20.4	6.8 <sup>b</sup>	5.9	2.8	2.5
Small Metro	21.5	20.9	6.4	6.0	2.7	2.7
250K – 1 Mil. Pop.	21.8	21.2	6.4	5.7	2.8	2.6
< 250K Pop.	20.8	20.3	6.3	6.6	2.5	2.8
Nonmetro	17.6	18.5	5.3	5.4	2.0	2.0
Urbanized	19.7	20.4	5.7	5.9	2.1	2.1
Less Urbanized	16.2	17.3	5.3	5.1	2.0	2.0
Completely Rural	14.9	15.2	3.9	4.1	1.4	1.7

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.79A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	2,482 <sup>b</sup>	2,188	1,642 <sup>b</sup>	1,441	693 <sup>b</sup>	549
<b>GEOGRAPHIC DIVISION</b>						
Northeast	324	272	215	192	84	64
New England	74	70	55	51	23	18
Middle Atlantic	250	202	160	141	61	46
Midwest	510 <sup>a</sup>	441	356 <sup>a</sup>	285	147	118
East North Central	374 <sup>b</sup>	304	256 <sup>a</sup>	195	107	81
West North Central	136	137	100	90	40	37
South	984 <sup>a</sup>	864	638	553	287 <sup>a</sup>	213
South Atlantic	455	401	296	264	136	96
East South Central	164	141	109	91	45	30
West South Central	366	323	233	198	105	87
West	664	611	433	411	175	154
Mountain	229	197	166	137	67 <sup>a</sup>	40
Pacific	435	414	267	274	108	114
<b>COUNTY TYPE</b>						
Large Metro	1,248	1,106	841	752	351	280
Small Metro	800	717	528	478	226	183
250K – 1 Mil. Pop.	537	472	364	308	157 <sup>a</sup>	116
< 250K Pop.	263	245	164	171	69	67
Nonmetro	433 <sup>a</sup>	364	272 <sup>b</sup>	211	116	86
Urbanized	217 <sup>a</sup>	163	123	103	54	40
Less Urbanized	175	164	114	87	48	41
Completely Rural	41	37	35 <sup>a</sup>	21	14	5

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.79B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	10.0 <sup>b</sup>	8.8	6.6 <sup>b</sup>	5.8	2.8 <sup>b</sup>	2.2
<b>GEOGRAPHIC DIVISION</b>						
Northeast	7.6	6.5	5.1	4.6	2.0	1.5
New England	6.7	6.4	4.9	4.6	2.1	1.6
Middle Atlantic	8.0	6.5	5.1	4.6	1.9	1.5
Midwest	9.4 <sup>a</sup>	8.2	6.6 <sup>a</sup>	5.3	2.7	2.2
East North Central	9.9 <sup>a</sup>	8.1	6.8 <sup>a</sup>	5.2	2.8	2.1
West North Central	8.3	8.3	6.1	5.5	2.4	2.3
South	10.6 <sup>a</sup>	9.2	6.9 <sup>a</sup>	5.9	3.1 <sup>a</sup>	2.3
South Atlantic	9.8	8.6	6.4	5.7	2.9	2.1
East South Central	11.1	9.6	7.4	6.2	3.1	2.1
West South Central	11.5	10.0	7.3	6.1	3.3	2.7
West	11.1	10.3	7.3	6.9	2.9	2.6
Mountain	12.3	10.5	8.9	7.3	3.6 <sup>a</sup>	2.1
Pacific	10.6	10.2	6.5	6.7	2.6	2.8
<b>COUNTY TYPE</b>						
Large Metro	9.2 <sup>a</sup>	8.2	6.2	5.5	2.6	2.1
Small Metro	10.8	9.6	7.1	6.4	3.1	2.5
250K – 1 Mil. Pop.	10.5	9.2	7.1	6.0	3.1 <sup>a</sup>	2.3
< 250K Pop.	11.4	10.5	7.1	7.3	3.0	2.9
Nonmetro	11.1 <sup>a</sup>	9.4	6.9 <sup>a</sup>	5.4	3.0	2.2
Urbanized	12.0	9.8	6.8	6.2	3.0	2.4
Less Urbanized	10.3	9.2	6.7 <sup>a</sup>	4.9	2.9	2.3
Completely Rural	10.0	8.4	8.4 <sup>a</sup>	4.7	3.3 <sup>a</sup>	1.1

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.80A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	9,709 <sup>a</sup>	9,239	4,729 <sup>b</sup>	4,245	1,832	1,661
<b>GEOGRAPHIC DIVISION</b>						
Northeast	1,637	1,522	876	780	362	279
New England	480	414	238	253	101	72
Middle Atlantic	1,157	1,108	638 <sup>a</sup>	527	261	208
Midwest	2,072	1,970	969	900	338	380
East North Central	1,448	1,328	687	636	265	263
West North Central	624	642	283	264	72 <sup>a</sup>	118
South	3,541	3,355	1,693	1,523	694	615
South Atlantic	1,782	1,648	876	788	351	293
East South Central	540	549	257	242	124	117
West South Central	1,220	1,158	561	493	219	205
West	2,459	2,392	1,190	1,042	438	387
Mountain	777	746	342	303	133	118
Pacific	1,682	1,645	848	738	305	268
<b>COUNTY TYPE</b>						
Large Metro	5,352	4,978	2,687 <sup>b</sup>	2,283	1,056	895
Small Metro	3,074	2,874	1,490	1,353	554	522
250K – 1 Mil. Pop.	1,940	1,808	932	861	357	330
< 250K Pop.	1,134	1,066	558	492	197	192
Nonmetro	1,283	1,387	552	609	223	244
Urbanized	618	691	242	312	91	127
Less Urbanized	554	591	259	254	115	112
Completely Rural	111	105	51	43	16	6

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.80B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	28.1 <sup>a</sup>	26.6	13.7 <sup>b</sup>	12.2	5.3	4.8
<b>GEOGRAPHIC DIVISION</b>						
Northeast	26.6	24.8	14.2	12.7	5.9	4.5
New England	29.6	25.4	14.7	15.5	6.2	4.4
Middle Atlantic	25.5	24.5	14.1 <sup>a</sup>	11.7	5.8	4.6
Midwest	28.1	26.6	13.2	12.1	4.6	5.1
East North Central	28.5	26.0	13.5	12.5	5.2	5.1
West North Central	27.3	27.9	12.4	11.5	3.2 <sup>a</sup>	5.1
South	27.8	26.1	13.3 <sup>a</sup>	11.8	5.4	4.8
South Atlantic	27.3	25.1	13.4	12.0	5.4	4.5
East South Central	26.7	26.9	12.7	11.9	6.1	5.7
West South Central	29.0	27.2	13.3	11.6	5.2	4.8
West	29.6	28.6	14.3	12.4	5.3	4.6
Mountain	31.1	29.5	13.7	12.0	5.4	4.7
Pacific	28.9	28.2	14.6	12.6	5.2	4.6
<b>COUNTY TYPE</b>						
Large Metro	28.3	26.6	14.2 <sup>b</sup>	12.2	5.6	4.8
Small Metro	28.4 <sup>a</sup>	26.4	13.8	12.4	5.1	4.8
250K – 1 Mil. Pop.	28.2	25.9	13.5	12.3	5.2	4.7
< 250K Pop.	28.9	27.5	14.2	12.7	5.0	4.9
Nonmetro	26.5	26.8	11.4	11.8	4.6	4.7
Urbanized	27.8	28.0	10.9	12.6	4.1	5.1
Less Urbanized	25.4	26.6	11.9	11.4	5.3	5.0
Completely Rural	24.9	22.1	11.4	9.0	3.6	1.2

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.81A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	42,198	41,745	10,296	9,662	4,306	4,274
<b>GEOGRAPHIC DIVISION</b>						
Northeast	7,155	6,622	1,647	1,452	609	572
New England	2,045	2,002	393	337	152	108
Middle Atlantic	5,110	4,620	1,254	1,115	457	465
Midwest	8,463	8,217	2,046	1,817	818	777
East North Central	6,236 <sup>a</sup>	5,567	1,542	1,332	667	578
West North Central	2,227	2,650	504	485	151	199
South	14,514	15,095	3,744	3,798	1,679	1,700
South Atlantic	7,755	8,406	1,739	2,146	762	874
East South Central	2,251	2,219	693	573	322	309
West South Central	4,509	4,471	1,313	1,079	595	518
West	12,067	11,812	2,858	2,595	1,199	1,225
Mountain	3,625	3,507	846	832	339	306
Pacific	8,442	8,304	2,012	1,763	860	919
<b>COUNTY TYPE</b>						
Large Metro	23,480	22,746	5,892	5,307	2,465	2,352
Small Metro	12,997	12,907	2,972	2,891	1,341	1,399
250K – 1 Mil. Pop.	9,070	9,044	2,082	1,879	979	945
< 250K Pop.	3,927	3,863	890	1,013	362	454
Nonmetro	5,722	6,093	1,433	1,464	500	523
Urbanized	2,809	3,026	697	711	249	230
Less Urbanized	2,356	2,531	637	634	215	230
Completely Rural	557	535	99	118	36	63

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.81B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	21.0	20.6	5.1	4.8	2.1	2.1
<b>GEOGRAPHIC DIVISION</b>						
Northeast	19.5	17.9	4.5	3.9	1.7	1.5
New England	21.1	20.5	4.1	3.5	1.6	1.1
Middle Atlantic	18.9	16.9	4.6	4.1	1.7	1.7
Midwest	19.6	18.9	4.7	4.2	1.9	1.8
East North Central	20.8 <sup>a</sup>	18.5	5.1	4.4	2.2	1.9
West North Central	16.9	19.9	3.8	3.6	1.1	1.5
South	19.5	20.0	5.0	5.0	2.3	2.3
South Atlantic	19.5	20.9	4.4	5.3	1.9	2.2
East South Central	18.9	18.5	5.8	4.8	2.7	2.6
West South Central	19.8	19.4	5.8	4.7	2.6	2.2
West	26.1	25.2	6.2	5.5	2.6	2.6
Mountain	25.8	24.5	6.0	5.8	2.4	2.1
Pacific	26.2	25.4	6.2	5.4	2.7	2.8
<b>COUNTY TYPE</b>						
Large Metro	22.0	20.9	5.5	4.9	2.3	2.2
Small Metro	21.6	21.3	4.9	4.8	2.2	2.3
250K – 1 Mil. Pop.	22.2	21.9	5.1	4.6	2.4	2.3
< 250K Pop.	20.3	20.0	4.6	5.2	1.9	2.4
Nonmetro	17.1	18.2	4.3	4.4	1.5	1.6
Urbanized	19.4	20.4	4.8	4.8	1.7	1.5
Less Urbanized	15.6	16.9	4.2	4.2	1.4	1.5
Completely Rural	14.3	15.1	2.5	3.3	0.9	1.8

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.82A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	51,908	50,984	15,025	13,907	6,138	5,935
<b>GEOGRAPHIC DIVISION</b>						
Northeast	8,792	8,144	2,523	2,232	971	851
New England	2,525	2,416	631	591	254	179
Middle Atlantic	6,267	5,728	1,892	1,642	718	672
Midwest	10,535	10,186	3,016	2,717	1,156	1,157
East North Central	7,685 <sup>a</sup>	6,895	2,229	1,968	932	840
West North Central	2,850	3,292	787	749	224	317
South	18,055	18,451	5,438	5,321	2,374	2,316
South Atlantic	9,536	10,054	2,615	2,933	1,113	1,167
East South Central	2,791	2,767	950	815	447	426
West South Central	5,728	5,629	1,873	1,572	814	723
West	14,525	14,203	4,048	3,637	1,637	1,612
Mountain	4,401	4,254	1,188	1,135	472	425
Pacific	10,124	9,949	2,860	2,502	1,165	1,187
<b>COUNTY TYPE</b>						
Large Metro	28,833	27,723	8,579 <sup>a</sup>	7,590	3,520	3,247
Small Metro	16,070	15,781	4,461	4,244	1,895	1,921
250K – 1 Mil. Pop.	11,010	10,853	3,013	2,739	1,336	1,275
< 250K Pop.	5,061	4,929	1,448	1,505	559	646
Nonmetro	7,005	7,479	1,984	2,073	723	768
Urbanized	3,427	3,718	939	1,024	340	357
Less Urbanized	2,909	3,122	896	888	330	342
Completely Rural	668	640	150	161	53	69

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.82B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	22.1	21.5	6.4 <sup>a</sup>	5.9	2.6	2.5
<b>GEOGRAPHIC DIVISION</b>						
Northeast	20.5	18.9	5.9	5.2	2.3	2.0
New England	22.3	21.2	5.6	5.2	2.2	1.6
Middle Atlantic	19.8	18.0	6.0	5.2	2.3	2.1
Midwest	20.9	20.0	6.0	5.3	2.3	2.3
East North Central	21.9 <sup>a</sup>	19.6	6.4	5.6	2.7	2.4
West North Central	18.4	21.1	5.1	4.8	1.4	2.0
South	20.7	20.9	6.2	6.0	2.7	2.6
South Atlantic	20.6	21.5	5.7	6.3	2.4	2.5
East South Central	20.0	19.7	6.8	5.8	3.2	3.0
West South Central	21.3	20.6	7.0	5.8	3.0	2.6
West	26.6	25.7	7.4	6.6	3.0	2.9
Mountain	26.6	25.3	7.2	6.7	2.9	2.5
Pacific	26.6	25.8	7.5	6.5	3.1	3.1
<b>COUNTY TYPE</b>						
Large Metro	22.9	21.7	6.8 <sup>a</sup>	6.0	2.8	2.5
Small Metro	22.6	22.1	6.3	5.9	2.7	2.7
250K – 1 Mil. Pop.	23.0	22.5	6.3	5.7	2.8	2.6
< 250K Pop.	21.7	21.3	6.2	6.5	2.4	2.8
Nonmetro	18.3	19.4	5.2	5.4	1.9	2.0
Urbanized	20.6	21.4	5.6	5.9	2.0	2.1
Less Urbanized	16.8	18.1	5.2	5.2	1.9	2.0
Completely Rural	15.3	15.9	3.4	4.0	1.2	1.7

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.