

Table 1.11C Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	956	1,015	648	677	492	509
12	25	25	20	20	14	13
13	31	31	26	24	20	14
14	41	46	33	37	23	26
15	46	45	39	39	25	26
16	55	54	49	46	36	35
17	61	57	53	49	40	34
18	104	86	95	78	65	56
19	98	85	82	72	67	55
20	105	96	92	77	64	61
21	96	93	79	74	58	57
22	94	92	77	68	58	55
23	92	97	69	72	46	52
24	82	91	60	67	47	51
25	86	89	60	67	45	46
26-29	348	369	217	238	162	185
30-34	372	404	211	257	163	206
35-39	370	333	181	177	134	137
40-44	346	379	187	184	131	130
45-49	378	359	177	168	137	122
50-54	550	617	244	296	201	225
55-59	519	511	228	189	174	142
60-64	445	447	124	138	97	93
65 or Older	385	443	151	201	109	131

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.11D Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	0.37	0.39	0.25	0.26	0.19	0.19
12	0.58	0.65	0.47	0.53	0.33	0.35
13	0.72	0.68	0.62	0.54	0.49	0.34
14	0.86	0.92	0.72	0.79	0.52	0.58
15	0.92	0.93	0.81	0.86	0.56	0.61
16	1.00	1.01	0.96	0.93	0.76	0.74
17	1.07	1.06	1.01	1.01	0.83	0.77
18	1.18	1.25	1.20	1.19	0.98	1.00
19	1.40	1.31	1.35	1.26	1.23	1.14
20	1.28	1.33	1.34	1.31	1.15	1.17
21	1.25	1.23	1.37	1.25	1.14	1.12
22	1.22	1.25	1.28	1.22	1.08	1.07
23	1.26	1.23	1.25	1.27	0.94	1.07
24	1.21	1.21	1.16	1.21	1.02	1.03
25	1.32	1.29	1.24	1.27	0.99	1.03
26-29	1.23	1.32	1.08	1.14	0.87	0.97
30-34	1.15	1.21	0.91	1.05	0.76	0.91
35-39	1.26	1.19	0.85	0.87	0.66	0.68
40-44	1.15	1.16	0.85	0.80	0.62	0.58
45-49	1.16	1.18	0.77	0.79	0.62	0.59
50-54	1.47	1.49	0.99	1.17	0.84	0.92
55-59	1.68	1.61	1.05	0.89	0.82	0.68
60-64	1.74	1.79	0.73	0.75	0.57	0.52
65 or Older	0.92	1.02	0.36	0.46	0.26	0.30

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.11P Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2013 Versus 2012

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	0.2653		0.7799		0.4809	
12	0.8945		0.9362		0.7758	
13	0.0867		0.0048		0.0030	
14	0.5215		0.7609		0.8606	
15	0.2654		0.6256		0.5443	
16	0.4140		0.4111		0.7337	
17	0.8177		0.8421		0.0884	
18	0.3241		0.5244		0.2715	
19	0.4964		0.9850		0.3509	
20	0.4094		0.1348		0.5546	
21	0.6423		0.5156		0.0800	
22	0.2271		0.1155		0.6565	
23	0.6155		0.5738		0.4151	
24	0.5308		0.7704		0.2019	
25	0.7978		0.6973		0.4942	
26-29	0.9705		0.3373		0.3043	
30-34	0.3150		0.2032		0.2195	
35-39	0.2014		0.4859		0.6012	
40-44	0.6800		0.7053		0.6422	
45-49	0.6329		0.6744		0.6534	
50-54	0.2936		0.8099		0.6099	
55-59	0.9413		0.1028		0.3900	
60-64	0.3665		0.6862		0.7131	
65 or Older	0.0014		0.1848		0.5927	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.12C Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	1,005	1,017	556	577	431	442
12	11	12	10	12	3	7
13	20	18	18	17	13	10
14	29	32	25	29	18	20
15	38	38	34	34	24	24
16	50	49	45	43	31	33
17	57	51	49	46	36	33
18	97	81	90	72	61	52
19	90	80	77	68	62	53
20	101	91	84	73	61	56
21	89	90	73	69	53	54
22	90	86	72	63	54	52
23	86	88	65	67	42	49
24	80	87	57	63	44	48
25	81	87	54	64	42	43
26-29	330	353	187	217	144	175
30-34	347	383	183	227	145	181
35-39	347	313	153	139	115	116
40-44	317	347	139	149	105	112
45-49	365	339	149	133	111	102
50-54	535	604	199	272	166	202
55-59	498	494	186	162	151	122
60-64	440	424	115	111	89	77
65 or Older	338	425	108	146	99	110

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.12D Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	0.39	0.39	0.21	0.22	0.17	0.17
12	0.26	0.33	0.24	0.31	0.08	0.20
13	0.49	0.42	0.43	0.39	0.32	0.23
14	0.64	0.68	0.57	0.65	0.41	0.45
15	0.80	0.82	0.73	0.76	0.54	0.55
16	0.96	0.96	0.91	0.88	0.67	0.71
17	1.06	1.03	0.98	0.95	0.78	0.74
18	1.23	1.21	1.24	1.14	0.97	0.95
19	1.40	1.31	1.31	1.25	1.15	1.11
20	1.30	1.35	1.30	1.28	1.11	1.11
21	1.27	1.29	1.29	1.22	1.06	1.08
22	1.27	1.25	1.25	1.17	1.05	1.02
23	1.24	1.23	1.21	1.24	0.88	1.01
24	1.25	1.24	1.15	1.19	0.98	0.99
25	1.34	1.31	1.18	1.24	0.95	0.98
26-29	1.26	1.33	0.97	1.07	0.79	0.91
30-34	1.17	1.20	0.84	0.96	0.69	0.83
35-39	1.29	1.21	0.74	0.70	0.57	0.59
40-44	1.16	1.18	0.66	0.66	0.51	0.50
45-49	1.20	1.17	0.67	0.64	0.51	0.50
50-54	1.52	1.54	0.83	1.09	0.69	0.84
55-59	1.67	1.68	0.89	0.77	0.72	0.59
60-64	1.82	1.74	0.68	0.61	0.53	0.43
65 or Older	0.81	0.98	0.26	0.34	0.24	0.25

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.12P Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	0.0739		0.1471		0.1983	
12	0.2384		0.2171		0.1096	
13	0.0924		0.1565		0.0761	
14	0.7995		0.9263		0.8508	
15	0.7865		0.9402		0.3146	
16	0.4584		0.8462		0.2145	
17	0.6809		0.6621		0.3971	
18	0.5254		0.4980		0.4517	
19	0.8453		0.9834		0.4706	
20	0.3925		0.2899		0.4427	
21	0.1967		0.2786		0.0509	
22	0.3627		0.2596		0.9801	
23	0.9280		0.6658		0.4481	
24	0.8088		0.3118		0.3786	
25	0.9498		0.0936		0.3162	
26-29	0.6314		0.0350		0.1249	
30-34	0.1330		0.0414		0.3959	
35-39	0.2581		0.7628		0.6686	
40-44	0.5297		0.7867		0.4494	
45-49	0.7989		0.4642		1.0000	
50-54	0.1646		0.2474		0.7253	
55-59	0.8873		0.4830		0.6969	
60-64	0.3933		0.7257		0.3279	
65 or Older	0.0002		0.5346		0.9307	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.13C Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	703	693	217	189	142	133
12	1	2	1	2	1	*
13	4	3	4	2	*	1
14	6	5	5	4	1	4
15	9	6	6	5	4	3
16	11	11	10	8	2	4
17	17	14	14	10	5	6
18	31	21	24	17	15	8
19	35	30	24	23	10	10
20	43	37	31	26	13	12
21	39	36	23	26	9	12
22	43	47	33	31	17	15
23	45	48	25	29	14	11
24	41	48	21	27	8	13
25	42	43	24	24	10	14
26-29	178	184	77	81	40	44
30-34	201	229	87	95	64	73
35-39	200	162	68	61	48	41
40-44	199	199	59	43	31	24
45-49	245	220	75	47	58	32
50-54	377	436	56	65	38	53
55-59	312	271	79	36	54	22
60-64	206	229	33	43	44	40
65 or Older	156	216	*	21	*	21

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.13D Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	0.27	0.26	0.08	0.07	0.05	0.05
12	0.03	0.06	0.03	0.06	0.02	*
13	0.10	0.06	0.10	0.05	*	0.03
14	0.14	0.11	0.11	0.10	0.03	0.10
15	0.20	0.14	0.14	0.12	0.10	0.06
16	0.27	0.26	0.24	0.19	0.05	0.10
17	0.38	0.34	0.32	0.25	0.11	0.15
18	0.63	0.44	0.50	0.36	0.33	0.17
19	0.80	0.71	0.56	0.54	0.23	0.24
20	0.91	0.79	0.67	0.57	0.30	0.26
21	0.84	0.79	0.52	0.59	0.21	0.28
22	0.89	0.96	0.70	0.67	0.37	0.34
23	0.94	0.99	0.55	0.64	0.32	0.24
24	0.90	0.98	0.50	0.61	0.20	0.29
25	0.95	0.99	0.59	0.57	0.25	0.34
26-29	0.89	0.91	0.44	0.46	0.23	0.25
30-34	0.90	1.01	0.43	0.46	0.32	0.36
35-39	0.93	0.82	0.35	0.32	0.24	0.21
40-44	0.90	0.84	0.29	0.20	0.16	0.11
45-49	0.99	0.94	0.34	0.23	0.27	0.16
50-54	1.35	1.56	0.24	0.28	0.16	0.23
55-59	1.36	1.21	0.38	0.18	0.26	0.11
60-64	1.12	1.21	0.20	0.24	0.26	0.23
65 or Older	0.37	0.50	*	0.05	*	0.05

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.13P Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2013 Versus 2012

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	0.6902		0.0673		0.5526	
12	0.8067		0.6504		*	
13	0.6130		0.4204		*	
14	0.1518		0.3794		0.1779	
15	0.2661		0.7400		0.3984	
16	0.8621		0.3957		0.2747	
17	0.2200		0.1457		0.4631	
18	0.0098		0.0673		0.2273	
19	0.2135		0.3816		0.6904	
20	0.3350		0.1198		0.9852	
21	0.2047		0.6738		0.8648	
22	0.9104		0.9908		0.9814	
23	0.6999		0.4940		0.4574	
24	0.6414		0.1270		0.3335	
25	0.7643		0.6529		0.6344	
26-29	0.4084		0.9678		0.8160	
30-34	0.7356		0.4452		0.7057	
35-39	0.0652		0.5042		0.6046	
40-44	0.1345		0.0138		0.1456	
45-49	0.1990		0.0427		0.1332	
50-54	0.1748		0.4433		0.2613	
55-59	0.7313		0.0988		0.2400	
60-64	0.1198		0.6684		0.9428	
65 or Older	0.0439		*		*	

*Low precision; no estimate reported.

NOTE: Cocaine use includes the use of crack.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.14C Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	346	364	111	77	94	64
12	*	*	*	*	*	*
13	0	1	0	1	*	1
14	1	1	1	*	*	*
15	4	3	3	2	*	*
16	7	3	5	2	1	*
17	6	3	5	2	*	1
18	10	9	6	3	4	1
19	10	8	7	5	2	2
20	15	11	8	3	4	*
21	12	11	4	4	2	3
22	16	10	6	4	3	3
23	17	18	5	5	3	1
24	14	18	6	3	3	2
25	17	19	6	12	3	11
26-29	86	93	25	17	8	10
30-34	114	102	52	30	46	25
35-39	99	90	32	19	28	16
40-44	101	94	24	17	12	16
45-49	129	113	51	14	42	11
50-54	177	223	42	43	29	40
55-59	130	163	47	31	46	14
60-64	93	70	19	20	13	20
65 or Older	48	88	*	21	*	21

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.14D Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	0.13	0.14	0.04	0.03	0.04	0.02
12	*	*	*	*	*	*
13	0.01	0.03	0.01	0.03	*	0.03
14	0.03	0.02	0.02	*	*	*
15	0.11	0.07	0.06	0.05	*	*
16	0.15	0.07	0.13	0.05	0.02	*
17	0.13	0.08	0.12	0.04	*	0.02
18	0.20	0.19	0.14	0.07	0.09	0.03
19	0.25	0.19	0.17	0.11	0.04	0.04
20	0.35	0.25	0.19	0.07	0.08	*
21	0.27	0.25	0.10	0.09	0.05	0.07
22	0.35	0.23	0.13	0.09	0.08	0.07
23	0.37	0.41	0.11	0.12	0.07	0.02
24	0.34	0.41	0.14	0.08	0.08	0.05
25	0.42	0.47	0.14	0.29	0.07	0.26
26-29	0.49	0.52	0.14	0.10	0.05	0.06
30-34	0.55	0.49	0.26	0.15	0.23	0.13
35-39	0.50	0.48	0.16	0.10	0.15	0.08
40-44	0.49	0.43	0.12	0.08	0.06	0.08
45-49	0.59	0.55	0.23	0.07	0.20	0.05
50-54	0.73	0.93	0.18	0.18	0.12	0.17
55-59	0.63	0.76	0.23	0.15	0.22	0.07
60-64	0.55	0.39	0.12	0.11	0.07	0.11
65 or Older	0.11	0.20	*	0.05	*	0.05

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.14P Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	0.6470		0.0260		0.5388	
12	*		*		*	
13	0.3534		0.2689		*	
14	0.4305		*		*	
15	0.5724		0.8137		*	
16	0.4060		0.4719		*	
17	0.6770		0.5385		*	
18	0.4343		0.2527		0.2291	
19	0.1265		0.2429		0.8747	
20	0.1457		0.0133		*	
21	0.6494		0.8957		0.6087	
22	0.0477		0.4029		0.9075	
23	0.9033		0.8211		0.4693	
24	0.8320		0.1222		0.3457	
25	0.7148		0.3314		0.2265	
26-29	0.4753		0.3121		0.8645	
30-34	0.0891		0.6235		0.8061	
35-39	0.3847		0.3195		0.3184	
40-44	0.0626		0.0991		0.7087	
45-49	0.3543		0.0141		0.0870	
50-54	0.2354		0.9378		0.3429	
55-59	0.1518		0.8573		0.4569	
60-64	0.2797		0.8666		0.6506	
65 or Older	0.1034		*		*	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.15C Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	674	707	175	204	90	132
12	4	7	3	6	3	2
13	8	8	7	5	4	3
14	14	10	11	8	6	6
15	15	12	11	9	6	5
16	21	19	19	15	10	8
17	26	23	19	19	10	12
18	42	37	31	30	22	22
19	47	42	36	33	18	19
20	57	46	44	32	15	14
21	49	52	27	38	13	14
22	55	55	29	35	16	16
23	52	58	26	33	15	14
24	47	55	21	29	8	17
25	54	51	26	25	9	14
26-29	204	215	74	72	32	41
30-34	224	256	69	84	45	35
35-39	216	198	57	58	26	50
40-44	183	215	29	50	10	15
45-49	210	189	38	26	35	7
50-54	294	368	26	43	26	42
55-59	292	269	11	8	4	*
60-64	210	208	44	13	*	*
65 or Older	197	228	46	87	13	76

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.15D Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	0.26	0.27	0.07	0.08	0.03	0.05
12	0.10	0.20	0.08	0.16	0.08	0.06
13	0.19	0.18	0.17	0.12	0.10	0.08
14	0.32	0.23	0.26	0.19	0.15	0.14
15	0.36	0.28	0.26	0.22	0.15	0.13
16	0.49	0.43	0.44	0.35	0.24	0.19
17	0.57	0.54	0.42	0.46	0.23	0.29
18	0.82	0.75	0.64	0.62	0.45	0.44
19	0.96	0.90	0.80	0.75	0.41	0.45
20	1.10	0.95	0.91	0.70	0.33	0.32
21	0.97	1.05	0.58	0.82	0.30	0.32
22	1.04	1.10	0.63	0.75	0.36	0.35
23	1.03	1.13	0.57	0.70	0.33	0.32
24	1.00	1.08	0.50	0.65	0.21	0.39
25	1.16	1.08	0.62	0.58	0.22	0.35
26-29	1.04	1.05	0.43	0.41	0.19	0.23
30-34	0.97	1.06	0.35	0.41	0.23	0.18
35-39	1.00	0.96	0.29	0.30	0.13	0.26
40-44	0.81	0.89	0.14	0.23	0.05	0.07
45-49	0.88	0.83	0.17	0.13	0.16	0.04
50-54	1.14	1.37	0.11	0.19	0.11	0.18
55-59	1.30	1.23	0.05	0.04	0.02	*
60-64	1.13	1.08	0.26	0.07	*	*
65 or Older	0.47	0.52	0.11	0.20	0.03	0.18

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.15P Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	0.1318		0.7534		0.2272	
12	0.1744		0.2450		0.5050	
13	0.8208		0.4504		0.3094	
14	0.0268		0.0715		0.9936	
15	0.1383		0.0696		0.5456	
16	0.3875		0.1564		0.2623	
17	0.2860		0.5745		0.1547	
18	0.1810		0.3130		0.6272	
19	0.8433		0.1837		0.7193	
20	0.6038		0.2395		0.5182	
21	0.4799		0.0346		0.8690	
22	0.9198		0.1868		0.5557	
23	0.2637		0.0495		0.7254	
24	0.4024		0.1609		0.0282	
25	0.1924		0.5267		0.5829	
26-29	0.9454		0.4607		0.4716	
30-34	0.5816		0.4223		0.9534	
35-39	0.7815		0.7476		0.2704	
40-44	0.1242		0.2531		0.9010	
45-49	0.3508		0.2733		0.3073	
50-54	0.0236		0.2926		0.3851	
55-59	0.9718		0.5189		*	
60-64	0.5044		0.2136		*	
65 or Older	0.1394		0.3264		0.2828	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.16C Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	489	519	120	105	65	64
12	18	16	11	11	7	7
13	20	19	14	11	10	6
14	25	28	18	14	8	7
15	26	24	14	14	5	8
16	23	20	15	11	7	4
17	23	20	13	13	9	4
18	37	28	21	16	13	7
19	32	27	13	13	5	7
20	36	32	14	17	8	10
21	31	32	18	19	13	5
22	33	37	14	11	6	7
23	30	33	8	12	3	4
24	30	34	4	12	1	4
25	36	25	10	11	4	7
26-29	136	136	40	30	15	13
30-34	179	185	23	35	10	28
35-39	151	162	35	23	14	18
40-44	162	160	60	27	5	23
45-49	153	148	45	30	27	29
50-54	202	293	25	27	21	24
55-59	210	160	37	53	37	14
60-64	116	142	9	7	*	7
65 or Older	144	145	*	3	*	3

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.16D Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	0.19	0.20	0.05	0.04	0.02	0.02
12	0.42	0.43	0.27	0.30	0.17	0.19
13	0.48	0.42	0.35	0.27	0.24	0.14
14	0.57	0.62	0.41	0.30	0.19	0.15
15	0.58	0.56	0.33	0.35	0.12	0.19
16	0.53	0.46	0.35	0.26	0.17	0.09
17	0.51	0.47	0.30	0.31	0.21	0.10
18	0.69	0.58	0.43	0.33	0.28	0.14
19	0.70	0.62	0.30	0.32	0.12	0.17
20	0.78	0.70	0.32	0.39	0.18	0.23
21	0.67	0.73	0.39	0.44	0.29	0.12
22	0.72	0.77	0.31	0.25	0.13	0.15
23	0.64	0.71	0.17	0.28	0.07	0.09
24	0.71	0.73	0.11	0.26	0.02	0.08
25	0.84	0.60	0.25	0.27	0.09	0.16
26-29	0.74	0.73	0.23	0.18	0.09	0.08
30-34	0.83	0.82	0.11	0.17	0.05	0.14
35-39	0.74	0.80	0.18	0.12	0.07	0.09
40-44	0.77	0.71	0.29	0.12	0.02	0.11
45-49	0.70	0.69	0.21	0.15	0.13	0.14
50-54	0.82	1.17	0.11	0.12	0.09	0.10
55-59	0.98	0.75	0.18	0.26	0.18	0.07
60-64	0.68	0.77	0.06	0.04	*	0.04
65 or Older	0.34	0.33	*	0.01	*	0.01

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.16P Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2013 Versus 2012

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	0.7554		0.2732		0.9873	
12	0.3818		0.8504		0.8567	
13	0.1729		0.0583		0.0263	
14	0.0667		0.0080		0.0457	
15	0.3354		0.9905		0.1314	
16	0.0052		0.0014		0.0663	
17	0.0845		0.3906		0.0694	
18	0.0588		0.1851		0.4094	
19	0.4631		0.9839		0.9400	
20	0.7360		0.5272		0.7166	
21	0.9051		0.9627		0.1923	
22	0.5947		0.9031		0.6324	
23	0.6472		0.2054		0.6633	
24	0.7110		0.0193		0.1025	
25	0.0005		0.2920		0.7279	
26-29	0.6298		0.8681		0.3862	
30-34	0.9356		0.2590		0.1433	
35-39	0.8969		0.2317		0.8018	
40-44	0.5822		0.7885		0.0571	
45-49	0.2867		0.5029		0.6982	
50-54	0.0040		0.7648		0.8667	
55-59	0.0534		0.3824		0.6551	
60-64	0.3745		0.8143		*	
65 or Older	0.6383		*		*	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.17C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	765	785	406	418	265	256
12	18	16	16	13	12	10
13	19	21	17	14	12	10
14	24	29	18	24	13	16
15	30	26	21	21	13	12
16	34	32	30	25	19	17
17	39	35	32	28	19	15
18	66	52	50	39	22	23
19	57	53	42	42	23	22
20	74	59	56	44	31	28
21	63	60	45	42	24	25
22	67	64	52	45	32	26
23	62	69	40	42	26	26
24	56	62	35	39	23	26
25	62	57	35	36	25	20
26-29	240	246	144	147	85	95
30-34	254	289	135	166	91	121
35-39	253	229	120	114	80	72
40-44	236	237	136	111	79	81
45-49	257	204	104	100	69	61
50-54	368	398	164	149	117	96
55-59	339	283	151	111	93	69
60-64	232	231	63	76	53	36
65 or Older	260	329	113	146	45	83

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.17D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	0.29	0.30	0.16	0.16	0.10	0.10
12	0.43	0.42	0.38	0.34	0.29	0.26
13	0.48	0.49	0.41	0.32	0.31	0.23
14	0.54	0.62	0.42	0.52	0.30	0.36
15	0.66	0.60	0.49	0.49	0.30	0.28
16	0.74	0.70	0.66	0.56	0.44	0.39
17	0.82	0.76	0.70	0.65	0.44	0.36
18	1.01	0.99	0.81	0.78	0.45	0.48
19	1.09	1.05	0.86	0.90	0.52	0.51
20	1.29	1.11	1.09	0.91	0.68	0.62
21	1.18	1.13	0.92	0.89	0.52	0.58
22	1.18	1.22	1.05	0.93	0.68	0.56
23	1.18	1.27	0.84	0.86	0.57	0.58
24	1.12	1.16	0.79	0.83	0.56	0.58
25	1.26	1.20	0.81	0.79	0.59	0.48
26-29	1.10	1.16	0.79	0.81	0.48	0.54
30-34	1.02	1.15	0.63	0.75	0.45	0.57
35-39	1.09	1.05	0.59	0.59	0.40	0.37
40-44	1.01	0.95	0.65	0.51	0.39	0.38
45-49	1.03	0.88	0.47	0.49	0.32	0.30
50-54	1.32	1.43	0.68	0.62	0.50	0.41
55-59	1.44	1.26	0.72	0.54	0.45	0.33
60-64	1.22	1.22	0.37	0.43	0.31	0.20
65 or Older	0.62	0.76	0.27	0.33	0.11	0.19

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.17P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2013 Versus 2012

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	0.1119		0.0121		0.2760	
12	0.3469		0.2313		0.4465	
13	0.2752		0.0081		0.1487	
14	0.5828		0.4378		0.6696	
15	0.0050		0.0915		0.1740	
16	0.3598		0.4404		0.4046	
17	0.0274		0.1601		0.0163	
18	0.0426		0.0468		0.1634	
19	0.1131		0.2874		0.1265	
20	0.2001		0.0519		0.6957	
21	0.8053		0.3561		0.3482	
22	0.1105		0.0272		0.1096	
23	0.8872		0.8228		0.8199	
24	0.5808		0.2397		0.8700	
25	0.2609		0.1708		0.1466	
26-29	0.4503		0.6885		0.9141	
30-34	0.7788		0.5137		0.3871	
35-39	0.7234		0.5714		0.6141	
40-44	0.6088		0.1412		0.7236	
45-49	0.0149		0.8604		0.9709	
50-54	0.4609		0.1480		0.6137	
55-59	0.0431		0.1139		0.2405	
60-64	0.9305		0.9604		0.0788	
65 or Older	0.0319		0.2777		0.1186	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.18C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	622	640	355	343	211	206
12	17	14	15	11	11	8
13	18	20	15	13	11	9
14	21	28	17	23	12	16
15	27	24	20	19	12	11
16	33	29	28	23	19	15
17	34	32	27	24	15	12
18	57	46	40	34	19	18
19	48	48	33	36	20	17
20	66	54	48	38	27	26
21	53	53	31	34	18	20
22	61	57	44	39	27	21
23	52	56	34	35	22	24
24	49	53	31	33	18	22
25	54	49	31	32	22	17
26-29	214	206	127	109	72	69
30-34	213	256	122	138	80	104
35-39	225	195	108	96	71	60
40-44	202	201	122	99	64	68
45-49	204	165	92	80	62	53
50-54	256	269	126	134	86	75
55-59	249	218	131	95	55	43
60-64	149	168	65	60	52	31
65 or Older	159	208	104	111	42	80

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.18D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2012 and 2013

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	0.24	0.24	0.14	0.13	0.08	0.08
12	0.42	0.35	0.37	0.28	0.28	0.21
13	0.44	0.46	0.38	0.30	0.27	0.22
14	0.49	0.60	0.39	0.50	0.28	0.35
15	0.61	0.56	0.46	0.45	0.28	0.25
16	0.71	0.64	0.62	0.53	0.43	0.34
17	0.71	0.71	0.60	0.57	0.34	0.29
18	0.95	0.89	0.69	0.68	0.40	0.38
19	1.02	1.00	0.73	0.79	0.46	0.41
20	1.23	1.07	0.97	0.81	0.61	0.57
21	1.06	1.08	0.67	0.76	0.41	0.46
22	1.13	1.13	0.91	0.82	0.59	0.46
23	1.04	1.12	0.73	0.74	0.50	0.53
24	1.03	1.05	0.71	0.72	0.44	0.49
25	1.18	1.05	0.72	0.71	0.55	0.41
26-29	1.04	1.02	0.70	0.62	0.41	0.40
30-34	0.92	1.05	0.58	0.64	0.39	0.50
35-39	1.01	0.92	0.54	0.49	0.36	0.32
40-44	0.90	0.84	0.58	0.45	0.32	0.32
45-49	0.87	0.76	0.42	0.39	0.29	0.26
50-54	1.01	1.09	0.53	0.56	0.37	0.32
55-59	1.11	1.00	0.63	0.46	0.26	0.21
60-64	0.83	0.91	0.38	0.34	0.31	0.17
65 or Older	0.38	0.48	0.25	0.25	0.10	0.19

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.18P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2013 Versus 2012

Age Category	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOTAL	0.0293		0.0019		0.2010	
12	0.0721		0.0372		0.0993	
13	0.3169		0.0261		0.1681	
14	0.2639		0.3009		0.8590	
15	0.0206		0.1498		0.1526	
16	0.1735		0.2801		0.1875	
17	0.0330		0.2285		0.0661	
18	0.0054		0.0176		0.0321	
19	0.3513		0.3910		0.0841	
20	0.2652		0.0557		0.6559	
21	0.8981		0.1359		0.6246	
22	0.1605		0.0440		0.1068	
23	0.6198		0.4531		0.6290	
24	0.2046		0.1293		0.9937	
25	0.0634		0.1244		0.1101	
26-29	0.0936		0.2234		0.4393	
30-34	0.4333		0.6088		0.5869	
35-39	0.3965		0.4059		0.6376	
40-44	0.6794		0.1876		0.9353	
45-49	0.1030		0.6786		0.8215	
50-54	0.9098		0.4397		0.8235	
55-59	0.0334		0.1665		0.6692	
60-64	0.6129		0.5829		0.2445	
65 or Older	0.0257		0.5908		0.1041	

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.