

**Table 1.74C Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	346	364	111	77	94	64
<b>GEOGRAPHIC DIVISION</b>						
Northeast	143	134	38	25	31	23
New England	46	62	10	16	8	16
Middle Atlantic	135	117	37	20	30	17
Midwest	136	161	44	47	33	40
East North Central	104	127	43	46	32	38
West North Central	87	99	10	12	5	11
South	225	222	95	49	81	35
South Atlantic	177	179	87	39	79	27
East South Central	82	91	16	20	15	18
West South Central	111	99	34	21	11	14
West	180	200	25	30	12	28
Mountain	90	91	19	21	12	21
Pacific	157	180	17	21	4	19
<b>COUNTY TYPE</b>						
Large Metro	273	274	92	51	79	48
Small Metro	205	204	61	47	47	33
250K – 1 Mil. Pop.	169	176	52	44	46	30
< 250K Pop.	116	101	32	17	9	15
Nonmetro	109	158	31	34	18	26
Urbanized	71	85	16	13	14	9
Less Urbanized	77	120	22	21	6	18
Completely Rural	39	57	16	23	9	17

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.74D Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.13	0.14	0.04	0.03	0.04	0.02
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.30	0.28	0.08	0.05	0.07	0.05
New England	0.37	0.49	0.08	0.13	0.07	0.12
Middle Atlantic	0.39	0.34	0.11	0.06	0.09	0.05
Midwest	0.24	0.29	0.08	0.08	0.06	0.07
East North Central	0.27	0.33	0.11	0.12	0.08	0.10
West North Central	0.51	0.57	0.06	0.07	0.03	0.07
South	0.23	0.23	0.10	0.05	0.08	0.04
South Atlantic	0.35	0.35	0.17	0.08	0.16	0.05
East South Central	0.53	0.59	0.10	0.13	0.10	0.11
West South Central	0.37	0.32	0.11	0.07	0.04	0.05
West	0.30	0.33	0.04	0.05	0.02	0.05
Mountain	0.49	0.48	0.10	0.11	0.06	0.11
Pacific	0.37	0.42	0.04	0.05	0.01	0.04
<b>COUNTY TYPE</b>						
Large Metro	0.19	0.19	0.07	0.04	0.06	0.03
Small Metro	0.25	0.25	0.08	0.06	0.06	0.04
250K – 1 Mil. Pop.	0.31	0.32	0.10	0.08	0.09	0.06
< 250K Pop.	0.42	0.37	0.12	0.07	0.04	0.06
Nonmetro	0.25	0.36	0.07	0.08	0.04	0.06
Urbanized	0.37	0.43	0.09	0.07	0.08	0.05
Less Urbanized	0.38	0.62	0.11	0.11	0.03	0.10
Completely Rural	0.78	1.17	0.34	0.49	0.20	0.36

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.74P Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2013 Versus 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.6470		0.0260		0.5388	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.5866		0.0861		0.6015	
New England	0.2015		0.9149		0.5196	
Middle Atlantic	0.2605		0.0547		0.3605	
Midwest	0.6991		0.9258		0.7905	
East North Central	0.4126		0.8658		0.9554	
West North Central	0.6282		0.7697		0.3804	
South	0.9237		0.0300		0.2705	
South Atlantic	0.6569		0.0835		0.2113	
East South Central	0.8062		0.9115		0.8606	
West South Central	0.2435		0.1202		0.5315	
West	0.9592		0.7100		0.2530	
Mountain	0.5897		0.8764		0.6725	
Pacific	0.8100		0.4877		0.1962	
<b>COUNTY TYPE</b>						
Large Metro	0.5323		0.0145		0.3326	
Small Metro	0.1860		0.5764		0.9841	
250K – 1 Mil. Pop.	0.7347		0.7508		0.9623	
< 250K Pop.	0.0558		0.0667		0.8403	
Nonmetro	0.0455		0.9309		0.5446	
Urbanized	0.2701		0.9259		0.3870	
Less Urbanized	0.1836		0.6817		0.4317	
Completely Rural	0.1743		0.3175		0.3099	

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.