

Table 1.76C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2012 and 2013

| Geographic Characteristic | Lifetime (2012) | Lifetime (2013) | Past Year (2012) | Past Year (2013) | Past Month (2012) | Past Month (2013) |
|----------------------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 489 | 519 | 120 | 105 | 65 | 64 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 202 | 240 | 35 | 32 | 10 | 21 |
| New England | 94 | 109 | 16 | 17 | 5 | 7 |
| Middle Atlantic | 177 | 215 | 31 | 27 | 9 | 20 |
| Midwest | 175 | 188 | 29 | 31 | 17 | 26 |
| East North Central | 147 | 141 | 27 | 29 | 15 | 26 |
| West North Central | 97 | 126 | 12 | 11 | 9 | 4 |
| South | 304 | 319 | 76 | 65 | 35 | 25 |
| South Atlantic | 251 | 257 | 69 | 59 | 30 | 19 |
| East South Central | 104 | 104 | 14 | 20 | 7 | 14 |
| West South Central | 145 | 162 | 29 | 18 | 17 | 7 |
| West | 282 | 301 | 77 | 68 | 49 | 48 |
| Mountain | 141 | 153 | 28 | 38 | 10 | 23 |
| Pacific | 242 | 261 | 72 | 56 | 48 | 42 |
| COUNTY TYPE | | | | | | |
| Large Metro | 414 | 458 | 91 | 78 | 61 | 52 |
| Small Metro | 322 | 293 | 49 | 58 | 16 | 23 |
| 250K – 1 Mil. Pop. | 258 | 267 | 44 | 54 | 12 | 15 |
| < 250K Pop. | 198 | 149 | 21 | 23 | 12 | 18 |
| Nonmetro | 161 | 199 | 63 | 37 | 12 | 27 |
| Urbanized | 135 | 151 | 61 | 30 | 5 | 24 |
| Less Urbanized | 92 | 121 | 12 | 18 | 7 | 7 |
| Completely Rural | 50 | 53 | 9 | 10 | 7 | 10 |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.76D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2012 and 2013

| Geographic Characteristic | Lifetime (2012) | Lifetime (2013) | Past Year (2012) | Past Year (2013) | Past Month (2012) | Past Month (2013) |
|----------------------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.19 | 0.20 | 0.05 | 0.04 | 0.02 | 0.02 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.43 | 0.51 | 0.07 | 0.07 | 0.02 | 0.04 |
| New England | 0.76 | 0.87 | 0.13 | 0.13 | 0.04 | 0.06 |
| Middle Atlantic | 0.51 | 0.61 | 0.09 | 0.08 | 0.03 | 0.06 |
| Midwest | 0.31 | 0.34 | 0.05 | 0.06 | 0.03 | 0.05 |
| East North Central | 0.38 | 0.36 | 0.07 | 0.07 | 0.04 | 0.07 |
| West North Central | 0.57 | 0.73 | 0.07 | 0.06 | 0.05 | 0.03 |
| South | 0.32 | 0.33 | 0.08 | 0.07 | 0.04 | 0.03 |
| South Atlantic | 0.49 | 0.50 | 0.14 | 0.12 | 0.06 | 0.04 |
| East South Central | 0.68 | 0.67 | 0.09 | 0.13 | 0.04 | 0.09 |
| West South Central | 0.48 | 0.53 | 0.10 | 0.06 | 0.06 | 0.02 |
| West | 0.47 | 0.49 | 0.13 | 0.11 | 0.08 | 0.08 |
| Mountain | 0.77 | 0.82 | 0.15 | 0.20 | 0.05 | 0.12 |
| Pacific | 0.57 | 0.61 | 0.17 | 0.13 | 0.11 | 0.10 |
| COUNTY TYPE | | | | | | |
| Large Metro | 0.27 | 0.30 | 0.06 | 0.06 | 0.04 | 0.04 |
| Small Metro | 0.36 | 0.33 | 0.06 | 0.07 | 0.02 | 0.03 |
| 250K – 1 Mil. Pop. | 0.43 | 0.44 | 0.08 | 0.10 | 0.02 | 0.03 |
| < 250K Pop. | 0.67 | 0.53 | 0.08 | 0.09 | 0.05 | 0.07 |
| Nonmetro | 0.36 | 0.44 | 0.15 | 0.09 | 0.03 | 0.06 |
| Urbanized | 0.67 | 0.71 | 0.33 | 0.16 | 0.03 | 0.13 |
| Less Urbanized | 0.46 | 0.62 | 0.06 | 0.10 | 0.04 | 0.04 |
| Completely Rural | 0.96 | 1.08 | 0.19 | 0.22 | 0.15 | 0.21 |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.76P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2013 Versus 2012

| Geographic Characteristic | Lifetime (2012) | Lifetime (2013) | Past Year (2012) | Past Year (2013) | Past Month (2012) | Past Month (2013) |
|----------------------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.7554 | | 0.2732 | | 0.9873 | |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.4577 | | 0.6760 | | 0.0217 | |
| New England | 0.7567 | | 0.6904 | | 0.3163 | |
| Middle Atlantic | 0.3102 | | 0.8020 | | 0.0376 | |
| Midwest | 0.3122 | | 0.1490 | | 0.5363 | |
| East North Central | 0.1521 | | 0.3214 | | 0.9877 | |
| West North Central | 0.8018 | | 0.1621 | | 0.0561 | |
| South | 0.4312 | | 0.4331 | | 0.1529 | |
| South Atlantic | 0.3180 | | 0.7048 | | 0.4071 | |
| East South Central | 0.9459 | | 0.8311 | | 0.7561 | |
| West South Central | 0.9604 | | 0.1737 | | 0.0508 | |
| West | 0.7101 | | 0.9041 | | 0.7109 | |
| Mountain | 0.7788 | | 0.2543 | | 0.2886 | |
| Pacific | 0.5572 | | 0.4665 | | 0.9851 | |
| COUNTY TYPE | | | | | | |
| Large Metro | 0.3544 | | 0.2850 | | 0.5248 | |
| Small Metro | 0.0319 | | 0.8348 | | 0.3171 | |
| 250K – 1 Mil. Pop. | 0.4738 | | 0.7629 | | 0.6726 | |
| < 250K Pop. | 0.0038 | | 0.8770 | | 0.3450 | |
| Nonmetro | 0.6768 | | 0.6463 | | 0.4627 | |
| Urbanized | 0.7727 | | 0.5576 | | 0.2671 | |
| Less Urbanized | 0.4325 | | 0.4893 | | 0.6548 | |
| Completely Rural | 0.7823 | | 0.4954 | | 0.9337 | |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.77C Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2012 and 2013

| Geographic Characteristic | Lifetime (2012) | Lifetime (2013) | Past Year (2012) | Past Year (2013) | Past Month (2012) | Past Month (2013) |
|----------------------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 51 | 52 | 34 | 29 | 19 | 15 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 20 | 15 | 11 | 11 | 4 | 5 |
| New England | 9 | 6 | 5 | 3 | 2 | 2 |
| Middle Atlantic | 18 | 13 | 10 | 10 | 3 | 5 |
| Midwest | 20 | 18 | 13 | 11 | 7 | 5 |
| East North Central | 17 | 15 | 11 | 11 | 5 | 5 |
| West North Central | 11 | 11 | 7 | 4 | 5 | 3 |
| South | 33 | 38 | 21 | 20 | 13 | 12 |
| South Atlantic | 24 | 32 | 15 | 16 | 7 | 10 |
| East South Central | 12 | 11 | 9 | 6 | 6 | 2 |
| West South Central | 20 | 17 | 12 | 11 | 9 | 6 |
| West | 31 | 26 | 20 | 14 | 12 | 5 |
| Mountain | 13 | 13 | 7 | 7 | 5 | 3 |
| Pacific | 28 | 23 | 18 | 12 | 10 | 4 |
| COUNTY TYPE | | | | | | |
| Large Metro | 46 | 39 | 29 | 23 | 16 | 11 |
| Small Metro | 28 | 34 | 16 | 16 | 8 | 9 |
| 250K – 1 Mil. Pop. | 22 | 29 | 13 | 13 | 7 | 6 |
| < 250K Pop. | 18 | 17 | 10 | 10 | 5 | 6 |
| Nonmetro | 20 | 17 | 12 | 11 | 7 | 7 |
| Urbanized | 14 | 10 | 9 | 5 | 4 | 3 |
| Less Urbanized | 13 | 13 | 7 | 9 | 5 | 6 |
| Completely Rural | 6 | 5 | 4 | 2 | 3 | 0 |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.77D Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2012 and 2013

| Geographic Characteristic | Lifetime (2012) | Lifetime (2013) | Past Year (2012) | Past Year (2013) | Past Month (2012) | Past Month (2013) |
|----------------------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.21 | 0.21 | 0.14 | 0.12 | 0.08 | 0.06 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.48 | 0.35 | 0.25 | 0.25 | 0.09 | 0.13 |
| New England | 0.82 | 0.51 | 0.42 | 0.32 | 0.16 | 0.20 |
| Middle Atlantic | 0.59 | 0.43 | 0.31 | 0.32 | 0.11 | 0.16 |
| Midwest | 0.37 | 0.33 | 0.24 | 0.21 | 0.13 | 0.10 |
| East North Central | 0.45 | 0.41 | 0.30 | 0.28 | 0.14 | 0.12 |
| West North Central | 0.68 | 0.65 | 0.42 | 0.26 | 0.29 | 0.16 |
| South | 0.35 | 0.40 | 0.23 | 0.21 | 0.14 | 0.13 |
| South Atlantic | 0.51 | 0.68 | 0.32 | 0.34 | 0.16 | 0.21 |
| East South Central | 0.82 | 0.76 | 0.59 | 0.41 | 0.37 | 0.14 |
| West South Central | 0.61 | 0.54 | 0.39 | 0.33 | 0.28 | 0.19 |
| West | 0.51 | 0.45 | 0.33 | 0.23 | 0.19 | 0.08 |
| Mountain | 0.73 | 0.71 | 0.40 | 0.39 | 0.26 | 0.16 |
| Pacific | 0.68 | 0.56 | 0.45 | 0.29 | 0.25 | 0.09 |
| COUNTY TYPE | | | | | | |
| Large Metro | 0.31 | 0.27 | 0.21 | 0.17 | 0.12 | 0.08 |
| Small Metro | 0.35 | 0.42 | 0.21 | 0.21 | 0.11 | 0.11 |
| 250K – 1 Mil. Pop. | 0.40 | 0.53 | 0.24 | 0.25 | 0.13 | 0.12 |
| < 250K Pop. | 0.65 | 0.67 | 0.40 | 0.39 | 0.20 | 0.26 |
| Nonmetro | 0.47 | 0.42 | 0.30 | 0.27 | 0.18 | 0.18 |
| Urbanized | 0.65 | 0.59 | 0.47 | 0.31 | 0.19 | 0.17 |
| Less Urbanized | 0.76 | 0.69 | 0.43 | 0.51 | 0.31 | 0.36 |
| Completely Rural | 1.54 | 1.06 | 1.03 | 0.54 | 0.79 | 0.04 |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.77P Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2013 Versus 2012

| Geographic Characteristic | Lifetime (2012) | Lifetime (2013) | Past Year (2012) | Past Year (2013) | Past Month (2012) | Past Month (2013) |
|----------------------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 0.0001 | | 0.0001 | | 0.0048 | |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 0.0003 | | 0.3460 | | 0.6203 | |
| New England | 0.0588 | | 0.2000 | | 0.6148 | |
| Middle Atlantic | 0.0021 | | 0.6379 | | 0.7576 | |
| Midwest | 0.0005 | | 0.0203 | | 0.0575 | |
| East North Central | 0.0028 | | 0.0952 | | 0.1800 | |
| West North Central | 0.0775 | | 0.0707 | | 0.1713 | |
| South | 0.5917 | | 0.1345 | | 0.2257 | |
| South Atlantic | 0.7676 | | 0.3026 | | 0.7121 | |
| East South Central | 0.5490 | | 0.3257 | | 0.0426 | |
| West South Central | 0.8419 | | 0.4964 | | 0.2141 | |
| West | 0.0833 | | 0.0048 | | 0.0095 | |
| Mountain | 0.3799 | | 0.5457 | | 0.1728 | |
| Pacific | 0.1376 | | 0.0045 | | 0.0255 | |
| COUNTY TYPE | | | | | | |
| Large Metro | 0.0029 | | 0.0053 | | 0.0048 | |
| Small Metro | 0.1697 | | 0.0834 | | 0.7410 | |
| 250K – 1 Mil. Pop. | 0.1997 | | 0.1140 | | 0.4373 | |
| < 250K Pop. | 0.5745 | | 0.4613 | | 0.6541 | |
| Nonmetro | 0.0058 | | 0.0489 | | 0.3314 | |
| Urbanized | 0.0370 | | 0.2020 | | 0.6648 | |
| Less Urbanized | 0.1637 | | 0.3656 | | 0.7956 | |
| Completely Rural | 0.1986 | | 0.0927 | | 0.0836 | |

*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.