

**Table 1.78C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	765	785	406	418	265	256
<b>GEOGRAPHIC DIVISION</b>						
Northeast	297	273	151	138	96	77
New England	117	142	64	65	41	35
Middle Atlantic	272	232	137	122	86	69
Midwest	282	304	157	132	85	81
East North Central	242	252	135	111	72	70
West North Central	152	178	80	69	43	41
South	464	505	250	280	168	176
South Atlantic	346	409	179	232	113	141
East South Central	172	159	97	83	64	64
West South Central	250	233	138	117	102	84
West	431	431	222	238	152	144
Mountain	200	220	95	127	59	64
Pacific	380	374	203	202	141	128
<b>COUNTY TYPE</b>						
Large Metro	771	750	347	326	222	198
Small Metro	513	524	212	241	131	154
250K – 1 Mil. Pop.	440	427	171	183	114	128
< 250K Pop.	278	322	134	164	66	89
Nonmetro	304	362	161	176	73	75
Urbanized	220	280	119	135	50	45
Less Urbanized	196	233	105	111	48	59
Completely Rural	116	91	34	39	23	23

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.78D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.29	0.30	0.16	0.16	0.10	0.10
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.63	0.58	0.32	0.29	0.20	0.16
New England	0.94	1.14	0.52	0.52	0.33	0.28
Middle Atlantic	0.78	0.66	0.39	0.35	0.25	0.20
Midwest	0.50	0.54	0.28	0.23	0.15	0.14
East North Central	0.62	0.65	0.35	0.29	0.19	0.18
West North Central	0.89	1.03	0.47	0.40	0.25	0.24
South	0.48	0.52	0.26	0.29	0.17	0.18
South Atlantic	0.68	0.79	0.35	0.45	0.22	0.27
East South Central	1.12	1.03	0.63	0.53	0.41	0.41
West South Central	0.83	0.76	0.46	0.38	0.34	0.28
West	0.71	0.70	0.37	0.39	0.25	0.23
Mountain	1.09	1.18	0.51	0.68	0.32	0.34
Pacific	0.90	0.88	0.48	0.47	0.33	0.30
<b>COUNTY TYPE</b>						
Large Metro	0.44	0.43	0.23	0.22	0.16	0.14
Small Metro	0.49	0.53	0.25	0.29	0.16	0.19
250K – 1 Mil. Pop.	0.62	0.65	0.31	0.33	0.21	0.24
< 250K Pop.	0.78	0.96	0.45	0.58	0.24	0.32
Nonmetro	0.59	0.70	0.35	0.39	0.17	0.17
Urbanized	1.00	1.14	0.61	0.64	0.27	0.22
Less Urbanized	0.78	0.98	0.48	0.56	0.24	0.30
Completely Rural	1.87	1.57	0.66	0.77	0.46	0.49

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.78P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2013 Versus 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.1119		0.0121		0.2760	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.0649		0.1145		0.2379	
New England	0.4720		0.5894		0.1535	
Middle Atlantic	0.0856		0.1344		0.5599	
Midwest	0.2473		0.0569		0.7647	
East North Central	0.0083		0.0619		0.2144	
West North Central	0.0770		0.5809		0.1294	
South	0.9422		0.4685		0.5129	
South Atlantic	0.5245		0.3922		0.9946	
East South Central	0.7706		0.2036		0.6771	
West South Central	0.5017		0.0396		0.3426	
West	0.3476		0.1340		0.7507	
Mountain	0.3705		0.5171		0.3010	
Pacific	0.5555		0.1759		0.9356	
<b>COUNTY TYPE</b>						
Large Metro	0.0512		0.0065		0.1814	
Small Metro	0.4126		0.3177		0.8866	
250K – 1 Mil. Pop.	0.4753		0.1184		0.4937	
< 250K Pop.	0.6812		0.6929		0.3596	
Nonmetro	0.3212		0.9418		0.9153	
Urbanized	0.6388		0.8430		0.8971	
Less Urbanized	0.3744		0.8099		0.9487	
Completely Rural	0.8947		0.8325		0.6827	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.79C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	64	63	53	50	37	32
<b>GEOGRAPHIC DIVISION</b>						
Northeast	22	19	19	17	11	9
New England	11	8	9	7	5	4
Middle Atlantic	19	17	17	15	9	8
Midwest	24	24	21	19	13	13
East North Central	20	19	17	15	11	10
West North Central	13	14	12	11	8	8
South	43	41	34	31	26	20
South Atlantic	30	28	24	22	19	14
East South Central	14	13	12	11	8	7
West South Central	26	25	21	18	15	13
West	38	37	31	30	19	19
Mountain	17	18	16	14	9	7
Pacific	34	33	26	26	17	18
<b>COUNTY TYPE</b>						
Large Metro	56	52	47	41	32	25
Small Metro	36	40	28	31	18	19
250K – 1 Mil. Pop.	31	33	24	25	15	16
< 250K Pop.	22	23	16	17	10	10
Nonmetro	26	26	19	17	12	11
Urbanized	20	17	14	13	9	9
Less Urbanized	15	21	11	10	6	7
Completely Rural	8	8	7	6	5	3

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.79D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.26	0.25	0.21	0.20	0.15	0.13
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.52	0.45	0.46	0.40	0.26	0.22
New England	1.01	0.76	0.85	0.67	0.49	0.37
Middle Atlantic	0.61	0.54	0.54	0.49	0.30	0.26
Midwest	0.44	0.44	0.38	0.35	0.24	0.24
East North Central	0.54	0.49	0.46	0.40	0.28	0.26
West North Central	0.80	0.84	0.72	0.67	0.48	0.48
South	0.46	0.44	0.37	0.33	0.28	0.21
South Atlantic	0.65	0.61	0.52	0.48	0.41	0.29
East South Central	0.98	0.91	0.81	0.77	0.57	0.45
West South Central	0.81	0.78	0.65	0.55	0.49	0.40
West	0.63	0.63	0.52	0.50	0.33	0.32
Mountain	0.90	0.93	0.86	0.74	0.48	0.39
Pacific	0.82	0.81	0.64	0.65	0.42	0.43
<b>COUNTY TYPE</b>						
Large Metro	0.37	0.34	0.32	0.28	0.22	0.18
Small Metro	0.44	0.47	0.36	0.37	0.23	0.24
250K – 1 Mil. Pop.	0.53	0.56	0.44	0.44	0.27	0.30
< 250K Pop.	0.82	0.86	0.63	0.68	0.43	0.44
Nonmetro	0.59	0.63	0.46	0.42	0.29	0.28
Urbanized	0.91	0.94	0.70	0.69	0.46	0.49
Less Urbanized	0.82	1.06	0.64	0.56	0.38	0.37
Completely Rural	1.71	1.80	1.56	1.37	1.11	0.64

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.79P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2013 Versus 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.0012		0.0061		0.0023	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.0922		0.4409		0.1903	
New England	0.8135		0.7892		0.4243	
Middle Atlantic	0.0720		0.4619		0.2894	
Midwest	0.0400		0.0164		0.1137	
East North Central	0.0106		0.0133		0.0775	
West North Central	0.9676		0.5457		0.8175	
South	0.0344		0.0458		0.0178	
South Atlantic	0.1641		0.2797		0.0633	
East South Central	0.2394		0.2451		0.1354	
West South Central	0.2016		0.1602		0.3160	
West	0.3370		0.6339		0.4637	
Mountain	0.1649		0.1482		0.0156	
Pacific	0.7234		0.7862		0.7672	
<b>COUNTY TYPE</b>						
Large Metro	0.0332		0.1171		0.0693	
Small Metro	0.0758		0.1775		0.0739	
250K – 1 Mil. Pop.	0.0935		0.0768		0.0385	
< 250K Pop.	0.4572		0.8308		0.8605	
Nonmetro	0.0489		0.0138		0.0648	
Urbanized	0.0993		0.5646		0.4281	
Less Urbanized	0.4142		0.0323		0.2827	
Completely Rural	0.4197		0.0206		0.0467	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.80C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	151	149	111	110	74	69
<b>GEOGRAPHIC DIVISION</b>						
Northeast	62	60	47	43	38	24
New England	33	29	24	26	21	13
Middle Atlantic	53	52	40	33	32	20
Midwest	59	54	45	42	27	26
East North Central	46	46	34	34	24	21
West North Central	39	28	30	24	12	16
South	97	92	66	70	43	43
South Atlantic	70	73	52	54	33	31
East South Central	33	33	21	23	15	17
West South Central	55	46	33	38	22	24
West	85	82	66	59	39	40
Mountain	36	36	25	27	15	17
Pacific	77	74	61	52	36	36
<b>COUNTY TYPE</b>						
Large Metro	170	156	111	97	66	57
Small Metro	130	107	82	73	39	38
250K – 1 Mil. Pop.	87	83	56	56	29	29
< 250K Pop.	96	73	61	47	25	25
Nonmetro	57	70	35	50	21	27
Urbanized	43	55	23	39	13	20
Less Urbanized	38	46	24	29	16	19
Completely Rural	17	17	11	11	6	3

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.80D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.44	0.43	0.32	0.32	0.21	0.20
<b>GEOGRAPHIC DIVISION</b>						
Northeast	1.02	0.97	0.76	0.70	0.62	0.39
New England	2.03	1.79	1.48	1.61	1.28	0.77
Middle Atlantic	1.18	1.15	0.89	0.74	0.71	0.45
Midwest	0.81	0.74	0.61	0.57	0.36	0.35
East North Central	0.91	0.91	0.67	0.67	0.47	0.40
West North Central	1.72	1.24	1.31	1.05	0.53	0.69
South	0.76	0.72	0.52	0.54	0.33	0.34
South Atlantic	1.07	1.12	0.80	0.83	0.50	0.48
East South Central	1.65	1.63	1.05	1.12	0.75	0.82
West South Central	1.31	1.08	0.78	0.88	0.53	0.57
West	1.02	0.98	0.79	0.70	0.46	0.47
Mountain	1.43	1.40	1.00	1.07	0.61	0.67
Pacific	1.32	1.26	1.04	0.90	0.62	0.61
<b>COUNTY TYPE</b>						
Large Metro	0.65	0.61	0.47	0.43	0.33	0.29
Small Metro	0.74	0.72	0.56	0.54	0.32	0.32
250K – 1 Mil. Pop.	0.84	0.92	0.66	0.67	0.40	0.39
< 250K Pop.	1.42	1.10	1.07	0.88	0.54	0.54
Nonmetro	0.91	0.92	0.64	0.77	0.42	0.48
Urbanized	1.42	1.21	0.89	1.14	0.56	0.71
Less Urbanized	1.34	1.53	0.96	1.09	0.70	0.77
Completely Rural	2.78	2.68	2.18	1.92	1.25	0.61

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.80P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2013 Versus 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.0122		0.0011		0.0771	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.1808		0.1325		0.0739	
New England	0.1609		0.7219		0.2620	
Middle Atlantic	0.5120		0.0280		0.1629	
Midwest	0.1704		0.2352		0.2787	
East North Central	0.0584		0.2679		0.8894	
West North Central	0.7788		0.6065		0.0253	
South	0.0890		0.0409		0.1709	
South Atlantic	0.1291		0.1850		0.1928	
East South Central	0.9289		0.5613		0.7139	
West South Central	0.3089		0.1347		0.6391	
West	0.4664		0.0743		0.3092	
Mountain	0.4282		0.2877		0.4687	
Pacific	0.6796		0.1479		0.4414	
<b>COUNTY TYPE</b>						
Large Metro	0.0523		0.0015		0.0675	
Small Metro	0.0480		0.0729		0.4806	
250K – 1 Mil. Pop.	0.0572		0.1797		0.4206	
< 250K Pop.	0.4100		0.2369		0.9048	
Nonmetro	0.7747		0.6868		0.8310	
Urbanized	0.9457		0.2508		0.2561	
Less Urbanized	0.5638		0.7725		0.8109	
Completely Rural	0.4717		0.4059		0.1006	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.81C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	740	751	380	386	247	244
<b>GEOGRAPHIC DIVISION</b>						
Northeast	288	265	137	127	87	71
New England	116	138	57	53	36	31
Middle Atlantic	263	226	124	115	79	64
Midwest	268	297	148	119	78	76
East North Central	231	245	128	103	66	66
West North Central	142	171	71	59	37	37
South	444	492	236	266	161	169
South Atlantic	331	393	169	221	108	137
East South Central	167	155	92	77	60	60
West South Central	236	230	132	108	98	79
West	423	406	209	216	143	139
Mountain	194	211	88	116	54	61
Pacific	374	352	190	183	132	125
<b>COUNTY TYPE</b>						
Large Metro	730	701	317	293	207	187
Small Metro	464	490	185	223	121	149
250K – 1 Mil. Pop.	408	394	152	163	105	121
< 250K Pop.	234	295	110	155	59	87
Nonmetro	281	341	150	166	65	68
Urbanized	201	266	111	127	45	37
Less Urbanized	180	212	96	105	41	54
Completely Rural	104	78	29	35	20	21

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.81D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.37	0.37	0.19	0.19	0.12	0.12
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.78	0.72	0.37	0.34	0.24	0.19
New England	1.19	1.41	0.59	0.55	0.37	0.32
Middle Atlantic	0.97	0.83	0.46	0.42	0.29	0.24
Midwest	0.62	0.68	0.34	0.27	0.18	0.18
East North Central	0.77	0.81	0.43	0.34	0.22	0.22
West North Central	1.07	1.29	0.54	0.44	0.28	0.28
South	0.60	0.65	0.32	0.35	0.22	0.22
South Atlantic	0.83	0.98	0.43	0.55	0.27	0.34
East South Central	1.40	1.29	0.77	0.64	0.51	0.50
West South Central	1.04	1.00	0.58	0.47	0.43	0.34
West	0.91	0.87	0.45	0.46	0.31	0.30
Mountain	1.38	1.47	0.63	0.81	0.38	0.43
Pacific	1.16	1.08	0.59	0.56	0.41	0.38
<b>COUNTY TYPE</b>						
Large Metro	0.56	0.53	0.28	0.26	0.19	0.17
Small Metro	0.61	0.66	0.30	0.35	0.20	0.24
250K – 1 Mil. Pop.	0.77	0.80	0.36	0.39	0.25	0.29
< 250K Pop.	0.97	1.21	0.53	0.74	0.30	0.42
Nonmetro	0.72	0.85	0.43	0.48	0.19	0.20
Urbanized	1.21	1.41	0.74	0.80	0.32	0.25
Less Urbanized	0.97	1.19	0.59	0.68	0.26	0.36
Completely Rural	2.14	1.75	0.73	0.93	0.51	0.58

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.81P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2013 Versus 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.3796		0.1677		0.8243	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.1393		0.2708		0.7096	
New England	0.7420		0.4264		0.3391	
Middle Atlantic	0.1338		0.3907		0.9643	
Midwest	0.4570		0.2025		0.6632	
East North Central	0.0332		0.1933		0.3145	
West North Central	0.0756		0.7881		0.3732	
South	0.5666		0.9925		0.9954	
South Atlantic	0.2925		0.1722		0.5567	
East South Central	0.8293		0.2909		0.8550	
West South Central	0.7518		0.1379		0.4801	
West	0.4483		0.3010		0.9653	
Mountain	0.5106		0.8460		0.6066	
Pacific	0.6073		0.2835		0.8058	
<b>COUNTY TYPE</b>						
Large Metro	0.1538		0.0875		0.5673	
Small Metro	0.7761		0.7316		0.7816	
250K – 1 Mil. Pop.	0.8038		0.2953		0.7855	
< 250K Pop.	0.8765		0.4632		0.3284	
Nonmetro	0.2828		0.8711		0.7860	
Urbanized	0.6156		0.9730		0.6382	
Less Urbanized	0.3841		0.9822		0.7915	
Completely Rural	0.7362		0.4953		0.2679	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.82C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	759	778	398	415	263	255
<b>GEOGRAPHIC DIVISION</b>						
Northeast	293	273	149	136	95	76
New England	114	142	64	65	41	35
Middle Atlantic	269	231	135	119	85	68
Midwest	280	302	155	130	84	81
East North Central	241	249	133	110	72	69
West North Central	149	178	78	69	41	41
South	458	502	244	280	166	176
South Atlantic	343	406	177	233	111	141
East South Central	171	158	95	81	63	64
West South Central	245	230	134	117	101	84
West	430	422	219	234	152	143
Mountain	198	217	93	125	58	65
Pacific	380	366	200	198	140	127
<b>COUNTY TYPE</b>						
Large Metro	761	743	338	322	218	198
Small Metro	505	514	209	236	130	153
250K – 1 Mil. Pop.	432	417	168	178	113	127
< 250K Pop.	272	316	130	162	64	88
Nonmetro	298	359	158	174	71	73
Urbanized	215	279	116	133	48	42
Less Urbanized	194	228	104	111	48	58
Completely Rural	112	87	31	36	21	21

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.82D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.32	0.33	0.17	0.17	0.11	0.11
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.68	0.63	0.35	0.31	0.22	0.18
New England	1.01	1.25	0.57	0.57	0.36	0.30
Middle Atlantic	0.85	0.73	0.43	0.38	0.27	0.21
Midwest	0.55	0.59	0.31	0.26	0.17	0.16
East North Central	0.69	0.71	0.38	0.31	0.21	0.20
West North Central	0.96	1.14	0.50	0.44	0.26	0.26
South	0.53	0.57	0.28	0.32	0.19	0.20
South Atlantic	0.74	0.87	0.38	0.50	0.24	0.30
East South Central	1.23	1.12	0.68	0.58	0.45	0.45
West South Central	0.91	0.84	0.50	0.43	0.37	0.31
West	0.79	0.76	0.40	0.42	0.28	0.26
Mountain	1.20	1.29	0.56	0.74	0.35	0.39
Pacific	1.00	0.95	0.53	0.52	0.37	0.33
<b>COUNTY TYPE</b>						
Large Metro	0.48	0.47	0.25	0.24	0.17	0.15
Small Metro	0.54	0.58	0.28	0.31	0.18	0.21
250K – 1 Mil. Pop.	0.68	0.71	0.34	0.36	0.23	0.26
< 250K Pop.	0.85	1.04	0.49	0.63	0.26	0.35
Nonmetro	0.64	0.77	0.38	0.43	0.18	0.19
Urbanized	1.08	1.24	0.66	0.70	0.29	0.23
Less Urbanized	0.86	1.08	0.53	0.61	0.26	0.33
Completely Rural	1.98	1.66	0.68	0.82	0.46	0.51

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.82P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2013 Versus 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.1738		0.0281		0.4773	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.0801		0.1319		0.2995	
New England	0.4740		0.6070		0.1782	
Middle Atlantic	0.1069		0.1535		0.6435	
Midwest	0.3109		0.1145		0.9587	
East North Central	0.0129		0.1193		0.3249	
West North Central	0.0771		0.6554		0.1099	
South	0.8051		0.6194		0.7215	
South Atlantic	0.4585		0.3291		0.8271	
East South Central	0.8453		0.2534		0.7962	
West South Central	0.5871		0.0599		0.4231	
West	0.3756		0.1446		0.8251	
Mountain	0.4238		0.6431		0.4972	
Pacific	0.5577		0.1581		0.9680	
<b>COUNTY TYPE</b>						
Large Metro	0.0675		0.0105		0.2679	
Small Metro	0.5042		0.4057		0.9318	
250K – 1 Mil. Pop.	0.5598		0.1828		0.6590	
< 250K Pop.	0.7351		0.7090		0.3387	
Nonmetro	0.2460		0.7357		0.6781	
Urbanized	0.5896		0.7888		0.9659	
Less Urbanized	0.3190		0.9876		0.8465	
Completely Rural	0.8070		0.5823		0.4438	

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.