

**Table 1.83C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	622	640	355	343	211	206
<b>GEOGRAPHIC DIVISION</b>						
Northeast	241	204	126	101	82	67
New England	103	115	51	51	29	26
Middle Atlantic	216	168	115	88	77	62
Midwest	245	226	145	112	77	68
East North Central	206	177	126	94	64	61
West North Central	133	138	68	55	41	31
South	377	401	215	234	132	148
South Atlantic	272	320	156	199	94	118
East South Central	150	130	88	78	55	60
West South Central	195	195	112	98	73	65
West	342	375	188	191	110	112
Mountain	157	189	82	113	49	45
Pacific	304	326	170	154	100	103
<b>COUNTY TYPE</b>						
Large Metro	573	570	297	258	174	166
Small Metro	389	376	176	191	99	122
250K – 1 Mil. Pop.	326	313	139	152	81	108
< 250K Pop.	223	221	107	113	56	60
Nonmetro	230	263	146	161	63	57
Urbanized	177	193	111	129	34	38
Less Urbanized	143	176	92	91	47	39
Completely Rural	74	71	29	34	23	21

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.83D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.24	0.24	0.14	0.13	0.08	0.08
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.51	0.43	0.27	0.21	0.17	0.14
New England	0.83	0.92	0.41	0.41	0.23	0.20
Middle Atlantic	0.62	0.48	0.33	0.25	0.22	0.18
Midwest	0.44	0.40	0.26	0.20	0.14	0.12
East North Central	0.53	0.45	0.32	0.24	0.16	0.16
West North Central	0.78	0.80	0.40	0.32	0.24	0.18
South	0.39	0.41	0.22	0.24	0.14	0.15
South Atlantic	0.53	0.62	0.31	0.39	0.18	0.23
East South Central	0.97	0.84	0.57	0.50	0.35	0.39
West South Central	0.65	0.64	0.37	0.32	0.24	0.21
West	0.57	0.61	0.31	0.31	0.18	0.18
Mountain	0.85	1.01	0.44	0.60	0.26	0.24
Pacific	0.72	0.77	0.40	0.36	0.24	0.24
<b>COUNTY TYPE</b>						
Large Metro	0.35	0.35	0.20	0.17	0.12	0.12
Small Metro	0.41	0.42	0.21	0.23	0.12	0.15
250K – 1 Mil. Pop.	0.52	0.53	0.25	0.27	0.15	0.20
< 250K Pop.	0.66	0.72	0.36	0.41	0.21	0.23
Nonmetro	0.47	0.54	0.33	0.36	0.14	0.13
Urbanized	0.83	0.84	0.57	0.63	0.18	0.20
Less Urbanized	0.59	0.78	0.44	0.46	0.24	0.20
Completely Rural	1.30	1.31	0.56	0.70	0.45	0.47

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.83P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2013 Versus 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.0293		0.0019		0.2010	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.0023		0.0243		0.2239	
New England	0.1230		0.3392		0.0325	
Middle Atlantic	0.0079		0.0417		0.6264	
Midwest	0.0810		0.0123		0.1302	
East North Central	0.0059		0.0079		0.0377	
West North Central	0.4695		0.7234		0.5956	
South	0.9131		0.2542		0.4969	
South Atlantic	0.4227		0.9104		0.6104	
East South Central	0.2039		0.3441		0.8969	
West South Central	0.5525		0.1290		0.4683	
West	0.5073		0.1927		0.8174	
Mountain	0.7347		0.6995		0.2337	
Pacific	0.3552		0.0829		0.4465	
<b>COUNTY TYPE</b>						
Large Metro	0.0302		0.0015		0.2111	
Small Metro	0.1908		0.1336		0.9873	
250K – 1 Mil. Pop.	0.3258		0.1536		0.9390	
< 250K Pop.	0.3792		0.5297		0.9151	
Nonmetro	0.5980		0.9131		0.3211	
Urbanized	0.9072		0.7726		0.7888	
Less Urbanized	0.3893		0.7267		0.1690	
Completely Rural	0.9477		0.7896		0.6776	

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.84C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	58	58	47	46	32	29
<b>GEOGRAPHIC DIVISION</b>						
Northeast	18	16	16	13	10	8
New England	9	7	7	6	5	4
Middle Atlantic	16	15	14	11	8	7
Midwest	22	22	19	18	12	11
East North Central	19	18	16	14	10	9
West North Central	12	12	11	10	8	7
South	38	38	29	29	21	18
South Atlantic	27	26	20	20	14	12
East South Central	14	12	12	11	8	6
West South Central	23	25	17	17	12	11
West	34	34	30	27	19	17
Mountain	16	15	15	12	9	6
Pacific	31	30	25	24	17	16
<b>COUNTY TYPE</b>						
Large Metro	50	46	41	37	27	22
Small Metro	33	36	26	27	17	17
250K – 1 Mil. Pop.	28	30	22	22	13	14
< 250K Pop.	20	20	15	15	10	9
Nonmetro	24	23	16	16	11	11
Urbanized	18	15	12	12	8	8
Less Urbanized	15	18	10	10	6	7
Completely Rural	6	8	6	5	4	2

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.84D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.23	0.23	0.19	0.18	0.13	0.11
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.43	0.39	0.37	0.31	0.23	0.19
New England	0.79	0.66	0.64	0.58	0.48	0.37
Middle Atlantic	0.51	0.47	0.45	0.37	0.26	0.22
Midwest	0.41	0.41	0.34	0.33	0.23	0.20
East North Central	0.49	0.48	0.41	0.38	0.26	0.23
West North Central	0.74	0.72	0.65	0.62	0.49	0.40
South	0.41	0.40	0.31	0.31	0.23	0.19
South Atlantic	0.59	0.55	0.43	0.44	0.31	0.26
East South Central	0.93	0.82	0.80	0.73	0.53	0.41
West South Central	0.73	0.76	0.53	0.54	0.39	0.36
West	0.58	0.57	0.50	0.45	0.31	0.29
Mountain	0.84	0.82	0.82	0.62	0.46	0.32
Pacific	0.75	0.74	0.62	0.59	0.40	0.40
<b>COUNTY TYPE</b>						
Large Metro	0.33	0.31	0.28	0.25	0.19	0.16
Small Metro	0.41	0.44	0.33	0.33	0.22	0.22
250K – 1 Mil. Pop.	0.49	0.53	0.40	0.39	0.25	0.27
< 250K Pop.	0.77	0.78	0.58	0.62	0.41	0.39
Nonmetro	0.54	0.57	0.39	0.39	0.27	0.27
Urbanized	0.81	0.87	0.60	0.64	0.42	0.47
Less Urbanized	0.79	0.92	0.55	0.53	0.37	0.36
Completely Rural	1.47	1.64	1.28	1.14	0.89	0.36

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.84P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:  
P Values from Tests of Differences of Percentages, 2013 Versus 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.0014		0.0076		0.0017	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.0648		0.1792		0.1770	
New England	0.3157		0.4625		0.4689	
Middle Atlantic	0.1165		0.2567		0.2536	
Midwest	0.0427		0.0273		0.0497	
East North Central	0.0198		0.0187		0.0532	
West North Central	0.8285		0.6610		0.4962	
South	0.0551		0.1533		0.0342	
South Atlantic	0.2845		0.5040		0.1796	
East South Central	0.0707		0.1568		0.1654	
West South Central	0.3463		0.4495		0.2373	
West	0.2942		0.4216		0.3191	
Mountain	0.1653		0.0884		0.0030	
Pacific	0.6499		0.9881		0.8163	
<b>COUNTY TYPE</b>						
Large Metro	0.0262		0.1085		0.0668	
Small Metro	0.0619		0.1070		0.0320	
250K – 1 Mil. Pop.	0.0676		0.0374		0.0168	
< 250K Pop.	0.4969		0.8439		0.6964	
Nonmetro	0.1447		0.0726		0.1179	
Urbanized	0.3425		0.9946		0.5043	
Less Urbanized	0.3683		0.0573		0.3536	
Completely Rural	0.6522		0.0823		0.0739	

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.85C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:  
Standard Errors of Numbers in Thousands, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	138	133	97	97	65	59
<b>GEOGRAPHIC DIVISION</b>						
Northeast	55	53	38	38	30	22
New England	28	28	14	25	11	10
Middle Atlantic	48	45	36	29	28	20
Midwest	54	52	37	36	24	20
East North Central	44	42	31	28	21	15
West North Central	33	29	20	23	10	13
South	81	82	57	61	40	34
South Atlantic	59	64	43	47	31	24
East South Central	31	30	20	20	14	14
West South Central	45	43	30	34	20	19
West	84	75	61	54	34	36
Mountain	32	32	22	25	13	14
Pacific	77	67	57	48	32	34
<b>COUNTY TYPE</b>						
Large Metro	139	127	88	82	56	49
Small Metro	105	91	63	60	30	31
250K – 1 Mil. Pop.	75	73	48	49	23	25
< 250K Pop.	73	60	41	35	19	19
Nonmetro	51	61	30	41	19	20
Urbanized	37	48	20	34	11	16
Less Urbanized	35	40	21	22	15	13
Completely Rural	16	13	10	9	5	3

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.85D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.40	0.38	0.28	0.28	0.19	0.17
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.90	0.86	0.62	0.62	0.49	0.36
New England	1.76	1.73	0.85	1.54	0.66	0.64
Middle Atlantic	1.06	0.99	0.78	0.64	0.62	0.44
Midwest	0.73	0.70	0.50	0.48	0.32	0.27
East North Central	0.86	0.82	0.61	0.54	0.42	0.30
West North Central	1.44	1.26	0.86	0.99	0.44	0.55
South	0.64	0.64	0.45	0.47	0.31	0.26
South Atlantic	0.90	0.98	0.66	0.71	0.48	0.37
East South Central	1.55	1.49	0.99	0.98	0.67	0.69
West South Central	1.07	1.02	0.71	0.79	0.48	0.44
West	1.01	0.89	0.73	0.64	0.41	0.43
Mountain	1.30	1.28	0.90	0.97	0.52	0.56
Pacific	1.32	1.15	0.98	0.82	0.55	0.57
<b>COUNTY TYPE</b>						
Large Metro	0.60	0.55	0.41	0.39	0.29	0.25
Small Metro	0.64	0.64	0.47	0.48	0.27	0.27
250K – 1 Mil. Pop.	0.80	0.84	0.60	0.62	0.33	0.35
< 250K Pop.	1.09	1.04	0.80	0.73	0.47	0.44
Nonmetro	0.86	0.86	0.57	0.67	0.38	0.37
Urbanized	1.29	1.09	0.79	1.05	0.49	0.57
Less Urbanized	1.29	1.42	0.83	0.87	0.64	0.54
Completely Rural	2.66	2.21	2.05	1.56	1.12	0.60

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.



**Table 1.85P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:  
P Values from Tests of Differences of Percentages, 2013 Versus 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.0033		0.0006		0.0342	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.0460		0.1254		0.1085	
New England	0.0214		0.8738		0.0378	
Middle Atlantic	0.4219		0.0858		0.3963	
Midwest	0.4346		0.1123		0.4374	
East North Central	0.0314		0.0236		0.0352	
West North Central	0.0921		0.6820		0.0490	
South	0.0423		0.0244		0.1351	
South Atlantic	0.0968		0.1171		0.1133	
East South Central	0.5947		0.2209		0.5718	
West South Central	0.3253		0.2663		0.8517	
West	0.2853		0.1314		0.6458	
Mountain	0.1236		0.1963		0.2482	
Pacific	0.6560		0.3138		0.9883	
<b>COUNTY TYPE</b>						
Large Metro	0.0125		0.0023		0.0438	
Small Metro	0.0887		0.0851		0.6674	
250K – 1 Mil. Pop.	0.1950		0.2574		0.7539	
< 250K Pop.	0.2764		0.1478		0.7462	
Nonmetro	0.8974		0.6576		0.3840	
Urbanized	0.9758		0.4252		0.7478	
Less Urbanized	0.7533		0.1801		0.2486	
Completely Rural	0.1464		0.3938		0.2412	

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.86C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	589	616	332	315	199	196
<b>GEOGRAPHIC DIVISION</b>						
Northeast	226	199	114	92	75	62
New England	103	112	49	43	27	23
Middle Atlantic	200	163	103	82	69	58
Midwest	231	218	138	98	71	64
East North Central	196	169	121	84	59	58
West North Central	123	133	60	48	36	29
South	360	396	205	219	127	141
South Atlantic	262	313	149	190	90	114
East South Central	143	125	84	71	52	56
West South Central	186	194	107	87	70	62
West	322	358	175	178	105	108
Mountain	145	184	73	104	46	40
Pacific	287	309	158	143	95	100
<b>COUNTY TYPE</b>						
Large Metro	535	541	274	231	164	156
Small Metro	348	349	158	174	91	118
250K – 1 Mil. Pop.	297	284	123	136	74	102
< 250K Pop.	191	203	97	105	52	59
Nonmetro	211	245	138	154	54	51
Urbanized	161	180	105	123	29	32
Less Urbanized	126	159	85	88	40	36
Completely Rural	64	64	24	32	20	20

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.86D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.29	0.30	0.17	0.16	0.10	0.10
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.61	0.54	0.31	0.25	0.20	0.17
New England	1.07	1.15	0.50	0.44	0.28	0.23
Middle Atlantic	0.74	0.60	0.38	0.30	0.26	0.21
Midwest	0.54	0.50	0.32	0.23	0.16	0.15
East North Central	0.65	0.56	0.40	0.28	0.20	0.19
West North Central	0.93	1.00	0.46	0.36	0.27	0.22
South	0.48	0.53	0.28	0.29	0.17	0.19
South Atlantic	0.66	0.78	0.37	0.47	0.23	0.28
East South Central	1.20	1.05	0.70	0.59	0.44	0.47
West South Central	0.82	0.84	0.47	0.38	0.31	0.27
West	0.70	0.76	0.38	0.38	0.23	0.23
Mountain	1.03	1.29	0.52	0.73	0.33	0.28
Pacific	0.89	0.95	0.49	0.44	0.29	0.31
<b>COUNTY TYPE</b>						
Large Metro	0.44	0.44	0.25	0.20	0.15	0.14
Small Metro	0.50	0.52	0.26	0.28	0.15	0.19
250K – 1 Mil. Pop.	0.63	0.65	0.30	0.33	0.18	0.25
< 250K Pop.	0.82	0.92	0.47	0.51	0.26	0.30
Nonmetro	0.56	0.65	0.40	0.45	0.16	0.15
Urbanized	1.00	1.04	0.70	0.79	0.20	0.22
Less Urbanized	0.70	0.94	0.53	0.57	0.26	0.24
Completely Rural	1.48	1.56	0.61	0.86	0.51	0.56

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.86P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2013 Versus 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.1951		0.0556		0.8288	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.0121		0.0853		0.5949	
New England	0.4530		0.3946		0.1870	
Middle Atlantic	0.0131		0.1346		0.9417	
Midwest	0.1493		0.0634		0.3123	
East North Central	0.0357		0.0645		0.2281	
West North Central	0.6870		0.6385		0.9475	
South	0.6295		0.7039		1.0000	
South Atlantic	0.2061		0.6920		0.9462	
East South Central	0.3056		0.6074		0.6698	
West South Central	0.8168		0.2728		0.6283	
West	0.7308		0.4455		0.5817	
Mountain	0.4088		0.2697		0.7371	
Pacific	0.4157		0.1348		0.4659	
<b>COUNTY TYPE</b>						
Large Metro	0.1537		0.0284		0.7149	
Small Metro	0.4425		0.4254		0.6724	
250K – 1 Mil. Pop.	0.5738		0.3952		0.7430	
< 250K Pop.	0.5805		0.8212		0.7826	
Nonmetro	0.5329		0.7408		0.5460	
Urbanized	0.8706		0.9144		0.7426	
Less Urbanized	0.3647		0.9060		0.2975	
Completely Rural	0.6942		0.4711		0.3528	

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.87C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	614	636	349	339	210	205
<b>GEOGRAPHIC DIVISION</b>						
Northeast	237	203	124	100	82	66
New England	102	115	52	51	29	25
Middle Atlantic	213	167	113	86	76	61
Midwest	242	223	142	109	76	67
East North Central	204	174	124	92	63	60
West North Central	129	139	64	56	38	31
South	372	401	210	233	130	148
South Atlantic	270	319	155	199	93	119
East South Central	149	128	86	76	54	59
West South Central	192	195	109	97	72	65
West	339	369	185	187	110	111
Mountain	156	188	80	111	48	45
Pacific	301	319	167	150	100	101
<b>COUNTY TYPE</b>						
Large Metro	563	564	290	253	172	164
Small Metro	381	370	172	187	98	122
250K – 1 Mil. Pop.	319	306	136	149	80	108
< 250K Pop.	218	218	103	110	55	59
Nonmetro	226	260	144	159	61	55
Urbanized	172	191	109	128	32	35
Less Urbanized	141	173	91	91	46	39
Completely Rural	72	67	27	33	21	20

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.87D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2012 and 2013**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.26	0.27	0.15	0.14	0.09	0.09
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.55	0.47	0.29	0.23	0.19	0.15
New England	0.90	1.01	0.46	0.45	0.26	0.22
Middle Atlantic	0.67	0.52	0.36	0.27	0.24	0.19
Midwest	0.48	0.44	0.28	0.21	0.15	0.13
East North Central	0.58	0.49	0.35	0.26	0.18	0.17
West North Central	0.83	0.89	0.41	0.36	0.25	0.20
South	0.43	0.45	0.24	0.26	0.15	0.17
South Atlantic	0.59	0.68	0.34	0.43	0.20	0.25
East South Central	1.07	0.92	0.62	0.54	0.39	0.42
West South Central	0.71	0.71	0.40	0.35	0.27	0.24
West	0.62	0.67	0.34	0.34	0.20	0.20
Mountain	0.94	1.12	0.48	0.66	0.29	0.27
Pacific	0.79	0.83	0.44	0.39	0.26	0.26
<b>COUNTY TYPE</b>						
Large Metro	0.38	0.38	0.22	0.19	0.14	0.13
Small Metro	0.45	0.46	0.23	0.25	0.14	0.17
250K – 1 Mil. Pop.	0.57	0.58	0.28	0.30	0.16	0.22
< 250K Pop.	0.73	0.79	0.39	0.44	0.23	0.25
Nonmetro	0.51	0.59	0.36	0.39	0.15	0.14
Urbanized	0.90	0.91	0.62	0.68	0.19	0.20
Less Urbanized	0.65	0.86	0.48	0.51	0.26	0.22
Completely Rural	1.41	1.39	0.57	0.75	0.46	0.49

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 1.87P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2013 Versus 2012**

<b>Geographic Characteristic</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOTAL</b>	0.0552		0.0054		0.4024	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.0032		0.0352		0.2849	
New England	0.1393		0.3955		0.0443	
Middle Atlantic	0.0101		0.0542		0.7107	
Midwest	0.1130		0.0259		0.2301	
East North Central	0.0099		0.0176		0.0727	
West North Central	0.4533		0.7787		0.4836	
South	0.9432		0.3360		0.6867	
South Atlantic	0.3694		0.9704		0.7223	
East South Central	0.2654		0.4355		0.7790	
West South Central	0.6368		0.1578		0.6002	
West	0.5600		0.2299		0.6917	
Mountain	0.6348		0.5132		0.5069	
Pacific	0.3669		0.0766		0.4695	
<b>COUNTY TYPE</b>						
Large Metro	0.0442		0.0027		0.3201	
Small Metro	0.2603		0.1953		0.7731	
250K – 1 Mil. Pop.	0.4151		0.2560		0.8157	
< 250K Pop.	0.4183		0.4996		0.8628	
Nonmetro	0.4986		0.7663		0.4543	
Urbanized	0.9369		0.7633		0.9193	
Less Urbanized	0.3297		0.8813		0.2034	
Completely Rural	0.8875		0.5762		0.4765	

\*Low precision; no estimate reported.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.