

Table 1.88C Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2012 and 2013

Drug or Method of Administration	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
HALLUCINOGENS	674	707	39	35	131	133	641	678
PCP (Angel Dust, Phencyclidine)	311	346	13	8	30	30	309	342
LSD (Acid)	561	580	21	19	87	81	538	567
Peyote	338	328	10	9	31	33	331	327
Mescaline	362	378	7	6	30	35	359	375
Psilocybin (Mushrooms)	524	561	24	22	99	100	502	542
Ecstasy (MDMA)	405	464	30	24	117	114	374	442
INHALANTS	489	519	51	52	84	85	471	505
Amyl Nitrite, Poppers, Locker Room Odorizers, or Rush	311	364	21	18	40	40	306	357
Correction Fluid, Degreaser, or Cleaning Fluid	162	100	26	26	28	26	157	91
Gasoline or Lighter Fluid	184	164	31	28	40	29	174	156
Glue, Shoe Polish, or Toluene	182	175	31	31	32	30	175	168
Halothane, Ether, or Other Anesthetics	89	85	8	10	15	17	88	84
Lacquer Thinner or Other Paint Solvents	132	117	21	18	21	21	130	114
Lighter Gases (Butane, Propane)	91	80	18	15	16	17	89	77
Nitrous Oxide or Whippits	365	409	18	17	71	70	349	399
Spray Paints	123	121	34	24	28	28	115	115
Other Aerosol Sprays	154	130	24	21	33	39	149	121
NEEDLE USE^{1,2,3}	225	265	10	10	38	40	222	261
Heroin Needle Use ^{1,2}	162	211	7	6	28	33	161	209
Cocaine Needle Use ^{1,2}	160	193	3	1	20	21	158	192
Stimulant Needle Use ^{1,2,3}	143	199	4	5	22	26	139	196
Methamphetamine Needle Use ^{1,2,3}	134	185	3	5	21	22	131	182
HEROIN	245	281	12	9	43	44	241	278
Smoke Heroin ²	153	152	9	5	30	32	148	150
Sniff or Snort Heroin ²	203	215	5	5	35	35	203	212

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

³ Estimates of Stimulant Needle Use, Methamphetamine Needle Use, and the Needle Use summary variable include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.88D Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Standard Errors of Percentages, 2012 and 2013

Drug or Method of Administration	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
HALLUCINOGENS	0.26	0.27	0.15	0.14	0.38	0.38	0.32	0.33
PCP (Angel Dust, Phencyclidine)	0.12	0.13	0.05	0.03	0.09	0.09	0.15	0.17
LSD (Acid)	0.22	0.22	0.09	0.08	0.25	0.23	0.27	0.28
Peyote	0.13	0.13	0.04	0.03	0.09	0.09	0.17	0.16
Mescaline	0.14	0.14	0.03	0.02	0.09	0.10	0.18	0.18
Psilocybin (Mushrooms)	0.20	0.21	0.10	0.09	0.29	0.29	0.25	0.27
Ecstasy (MDMA)	0.16	0.18	0.12	0.10	0.34	0.33	0.19	0.22
INHALANTS	0.19	0.20	0.21	0.21	0.24	0.24	0.23	0.25
Amyl Nitrite, Poppers, Locker Room Odorizers, or Rush	0.12	0.14	0.08	0.07	0.12	0.12	0.15	0.18
Correction Fluid, Degreaser, or Cleaning Fluid	0.06	0.04	0.10	0.11	0.08	0.08	0.08	0.05
Gasoline or Lighter Fluid	0.07	0.06	0.13	0.11	0.12	0.08	0.09	0.08
Glue, Shoe Polish, or Toluene	0.07	0.07	0.13	0.12	0.09	0.09	0.09	0.08
Halothane, Ether, or Other Anesthetics	0.03	0.03	0.03	0.04	0.04	0.05	0.04	0.04
Lacquer Thinner or Other Paint Solvents	0.05	0.04	0.08	0.07	0.06	0.06	0.07	0.06
Lighter Gases (Butane, Propane)	0.04	0.03	0.07	0.06	0.05	0.05	0.04	0.04
Nitrous Oxide or Whippits	0.14	0.16	0.07	0.07	0.21	0.20	0.17	0.20
Spray Paints	0.05	0.05	0.14	0.10	0.08	0.08	0.06	0.06
Other Aerosol Sprays	0.06	0.05	0.10	0.08	0.10	0.11	0.07	0.06
NEEDLE USE^{1,2,3}	0.09	0.10	0.04	0.04	0.11	0.11	0.11	0.13
Heroin Needle Use ^{1,2}	0.06	0.08	0.03	0.02	0.08	0.09	0.08	0.10
Cocaine Needle Use ^{1,2}	0.06	0.07	0.01	0.00	0.06	0.06	0.08	0.09
Stimulant Needle Use ^{1,2,3}	0.05	0.08	0.02	0.02	0.06	0.07	0.07	0.10
Methamphetamine Needle Use ^{1,2,3}	0.05	0.07	0.01	0.02	0.06	0.06	0.07	0.09
HEROIN	0.09	0.11	0.05	0.04	0.12	0.13	0.12	0.14
Smoke Heroin ²	0.06	0.06	0.04	0.02	0.09	0.09	0.07	0.07
Sniff or Snort Heroin ²	0.08	0.08	0.02	0.02	0.10	0.10	0.10	0.10

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

³ Estimates of Stimulant Needle Use, Methamphetamine Needle Use, and the Needle Use summary variable include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.88P Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Drug or Method of Administration	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
HALLUCINOGENS	0.1318		0.0231		0.9354		0.1001	
PCP (Angel Dust, Phencyclidine)	0.7531		0.0008		0.6347		0.8517	
LSD (Acid)	0.3144		0.3394		0.1296		0.4213	
Peyote	0.6359		0.4603		0.7695		0.6123	
Mescaline	0.7687		0.6085		0.7711		0.7588	
Psilocybin (Mushrooms)	0.0277		0.4732		0.4598		0.0350	
Ecstasy (MDMA)	0.0142		0.0024		0.7920		0.0052	
INHALANTS	0.7554		0.0001		0.0046		0.5668	
Amyl Nitrite, Poppers, Locker Room Odorizers, or Rush	0.2034		0.0534		0.7237		0.1755	
Correction Fluid, Degreaser, or Cleaning Fluid	0.0532		0.1438		0.5439		0.1076	
Gasoline or Lighter Fluid	0.0045		0.0017		0.0069		0.0626	
Glue, Shoe Polish, or Toluene	0.3043		0.0107		0.5409		0.6184	
Halothane, Ether, or Other Anesthetics	0.6157		0.6285		0.5659		0.6483	
Lacquer Thinner or Other Paint Solvents	0.2327		0.0052		0.5023		0.4954	
Lighter Gases (Butane, Propane)	0.8674		0.0458		0.7249		0.8738	
Nitrous Oxide or Whippits	0.8217		0.1846		0.0021		0.4024	
Spray Paints	0.5304		0.0000		0.7799		0.6171	
Other Aerosol Sprays	0.4444		0.0206		0.7855		0.7513	
NEEDLE USE^{1,2,3}	0.2843		0.9420		0.9355		0.2929	
Heroin Needle Use ^{1,2}	0.0775		0.8556		0.5008		0.0951	
Cocaine Needle Use ^{1,2}	0.3290		0.2797		0.9538		0.3292	
Stimulant Needle Use ^{1,2,3}	0.0357		0.9465		0.3147		0.0479	
Methamphetamine Needle Use ^{1,2,3}	0.2062		0.6530		0.6648		0.2289	
HEROIN	0.5592		0.1537		0.4432		0.4421	
Smoke Heroin ²	0.5583		0.2071		0.5444		0.6800	
Sniff or Snort Heroin ²	0.8671		0.6165		0.5147		0.9584	

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

³ Estimates of Stimulant Needle Use, Methamphetamine Needle Use, and the Needle Use summary variable include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.89C Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2012 and 2013

Pain Reliever	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
Darvocet [®] , Darvon [®] , or Tylenol [®] with Codeine	443	453	39	40	88	79	432	436
Percocet [®] , Percodan [®] , or Tylox [®]	390	408	25	21	90	93	375	393
Vicodin [®] , Lortab [®] , or Lorcet [®]	512	481	40	36	118	118	479	461
Codeine	264	293	25	24	74	76	250	284
Demerol [®]	144	196	9	8	27	25	139	195
Dilaudid [®]	162	144	4	5	34	31	158	140
Fioricet [®]	60	94	3	2	12	14	59	93
Fiorinal [®]	51	85	5	4	11	10	49	84
Hydrocodone	358	359	26	29	94	89	333	344
Methadone	142	159	14	11	45	45	134	153
Morphine	176	200	17	17	55	57	165	192
OxyContin [®]	245	280	23	22	77	74	228	268
Phenaphen [®] with Codeine	72	81	10	7	21	23	67	78
Propoxyphene	50	42	9	4	14	13	47	40
SK-65 [®]	12	17	4	7	9	7	*	14
Stadol [®]	33	43	3	2	9	4	32	43
Talacen [®]	18	17	4	4	8	7	16	15
Talwin [®]	51	60	6	3	9	3	50	60
Talwin [®] NX	39	18	2	3	9	9	38	15
Tramadol	157	192	12	11	47	52	146	186
Ultram [®]	105	145	9	6	24	29	101	142
SELECTED GROUPS OF DRUGS¹								
Propoxyphene or Codeine Products ²	463	479	42	43	102	95	447	460
Oxycodone Products ³	409	443	30	28	102	100	395	427
Hydrocodone Products ⁴	535	508	44	41	123	123	498	488
Tramadol Products ⁵	170	210	13	12	49	52	159	204

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Pain Relievers module but fall into this category.

² Includes Darvocet[®], Darvon[®] or Tylenol[®] with Codeine, codeine, Phenaphen[®] with Codeine, propoxyphene, and SK-65[®].

³ Includes Percocet[®], Percodan[®] or Tylox[®], and OxyContin[®].

⁴ Includes Vicodin[®], Lortab[®], or Lorcet[®], and hydrocodone.

⁵ Includes tramadol and Ultram[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.89D Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Standard Errors of Percentages, 2012 and 2013

Pain Reliever	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
Darvocet [®] , Darvon [®] , or Tylenol [®] with Codeine	0.17	0.17	0.16	0.16	0.25	0.23	0.22	0.21
Percocet [®] , Percodan [®] , or Tylox [®]	0.15	0.16	0.10	0.09	0.26	0.27	0.19	0.19
Vicodin [®] , Lortab [®] , or Lorcet [®]	0.20	0.18	0.16	0.14	0.34	0.34	0.24	0.23
Codeine	0.10	0.11	0.10	0.09	0.21	0.22	0.12	0.14
Demerol [®]	0.06	0.07	0.03	0.03	0.08	0.07	0.07	0.10
Dilaudid [®]	0.06	0.06	0.02	0.02	0.10	0.09	0.08	0.07
Fioricet [®]	0.02	0.04	0.01	0.01	0.03	0.04	0.03	0.05
Fiorinal [®]	0.02	0.03	0.02	0.02	0.03	0.03	0.02	0.04
Hydrocodone	0.14	0.14	0.10	0.12	0.27	0.26	0.17	0.17
Methadone	0.05	0.06	0.06	0.04	0.13	0.13	0.07	0.08
Morphine	0.07	0.08	0.07	0.07	0.16	0.16	0.08	0.09
OxyContin [®]	0.09	0.11	0.09	0.09	0.22	0.21	0.11	0.13
Phenaphen [®] with Codeine	0.03	0.03	0.04	0.03	0.06	0.07	0.03	0.04
Propoxyphene	0.02	0.02	0.04	0.02	0.04	0.04	0.02	0.02
SK-65 [®]	0.00	0.01	0.02	0.03	0.03	0.02	*	0.01
Stadol [®]	0.01	0.02	0.01	0.01	0.02	0.01	0.02	0.02
Talacen [®]	0.01	0.01	0.02	0.02	0.02	0.02	0.01	0.01
Talwin [®]	0.02	0.02	0.02	0.01	0.03	0.01	0.02	0.03
Talwin [®] NX	0.01	0.01	0.01	0.01	0.02	0.03	0.02	0.01
Tramadol	0.06	0.07	0.05	0.04	0.14	0.15	0.07	0.09
Ultram [®]	0.04	0.06	0.03	0.03	0.07	0.08	0.05	0.07
SELECTED GROUPS OF DRUGS¹								
Propoxyphene or Codeine Products ²	0.18	0.18	0.17	0.17	0.30	0.27	0.22	0.23
Oxycodone Products ³	0.16	0.17	0.12	0.11	0.29	0.29	0.20	0.21
Hydrocodone Products ⁴	0.21	0.19	0.18	0.17	0.36	0.35	0.25	0.24
Tramadol Products ⁵	0.07	0.08	0.05	0.05	0.14	0.15	0.08	0.10

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Pain Relievers module but fall into this category.

² Includes Darvocet[®], Darvon[®] or Tylenol[®] with Codeine, codeine, Phenaphen[®] with Codeine, propoxyphene, and SK-65[®].

³ Includes Percocet[®], Percodan[®] or Tylox[®], and OxyContin[®].

⁴ Includes Vicodin[®], Lortab[®], or Lorcet[®], and hydrocodone.

⁵ Includes tramadol and Ultram[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.89P Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Pain Reliever	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
Darvocet [®] , Darvon [®] , or Tylenol [®] with Codeine	0.0384		0.0459		0.0024		0.1615	
Percocet [®] , Percodan [®] , or Tylox [®]	0.5703		0.0262		0.1233		0.9453	
Vicodin [®] , Lortab [®] , or Lorcet [®]	0.0165		0.0043		0.0001		0.1983	
Codeine	0.3374		0.0618		0.6954		0.2918	
Demerol [®]	0.1004		0.3174		0.0938		0.0486	
Dilaudid [®]	0.8254		0.4910		0.8919		0.7745	
Fioricet [®]	0.0654		0.1083		0.7059		0.0631	
Fiorinal [®]	0.1315		0.4045		0.9027		0.1161	
Hydrocodone	0.8908		0.9429		0.0411		0.4828	
Methadone	0.9741		0.0638		0.3386		0.6257	
Morphine	0.5113		0.3222		0.1379		0.2142	
OxyContin [®]	0.3663		0.1443		0.2325		0.1421	
Phenaphen [®] with Codeine	0.2748		0.1575		0.2768		0.3202	
Propoxyphene	0.4896		0.0755		0.9187		0.6689	
SK-65 [®]	0.7515		0.6914		0.7744		*	
Stadol [®]	0.2654		0.3906		0.5164		0.1856	
Talacen [®]	0.7618		0.4160		0.9194		0.8196	
Talwin [®]	0.9515		0.0523		0.1192		0.6778	
Talwin [®] NX	0.0838		0.9742		0.9329		0.0531	
Tramadol	0.0671		0.9295		0.3449		0.1003	
Ultram [®]	0.0596		0.6120		0.3616		0.0808	
SELECTED GROUPS OF DRUGS¹								
Propoxyphene or Codeine Products ²	0.1162		0.0400		0.1714		0.2501	
Oxycodone Products ³	0.7689		0.0439		0.0898		0.7885	
Hydrocodone Products ⁴	0.0373		0.0109		0.0006		0.2748	
Tramadol Products ⁵	0.0510		0.8121		0.3682		0.0769	

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Pain Relievers module but fall into this category.

² Includes Darvocet[®], Darvon[®] or Tylenol[®] with Codeine, codeine, Phenaphen[®] with Codeine, propoxyphene, and SK-65[®].

³ Includes Percocet[®], Percodan[®] or Tylox[®], and OxyContin[®].

⁴ Includes Vicodin[®], Lortab[®], or Lorcet[®], and hydrocodone.

⁵ Includes tramadol and Ultram[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.90C Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2012 and 2013

Tranquilizer	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
Klonopin [®] or Clonazepam	240	293	18	16	65	68	228	284
Xanax [®] , Alprazolam, Ativan [®] , or Lorazepam	423	454	32	28	104	102	393	435
Valium [®] or Diazepam	435	443	22	14	74	70	424	433
Atarax [®]	54	64	5	4	13	10	53	63
BuSpar [®]	81	73	6	4	21	23	78	69
Equanil [®]	18	31	4	2	9	9	15	30
Flexeril [®]	184	189	7	8	38	37	182	186
Librium [®]	111	106	4	4	14	13	110	105
Limbitrol [®]	17	13	4	4	9	6	13	11
Meprobamate	43	46	5	3	10	6	42	45
Miltown [®]	55	23	3	3	10	8	54	22
Rohypnol [®]	52	71	8	4	16	15	49	70
Serax [®]	16	18	4	2	11	5	10	17
Soma [®]	201	196	11	12	51	46	190	187
Tranxene [®]	38	20	2	2	10	7	37	19
Vistaril [®]	48	56	5	4	14	17	45	53
SELECTED GROUPS OF DRUGS¹								
Benzodiazepines ²	533	564	37	30	107	107	510	546
Meprobamate Products ³	61	57	6	3	12	10	59	56
Muscle Relaxants ⁴	249	245	12	14	58	53	239	237

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Tranquilizers module but fall into this category.

² Includes Klonopin[®] or clonazepam, Xanax[®], alprazolam, Ativan[®] or lorazepam, Valium[®] or diazepam, Librium[®], Limbitrol[®], Rohypnol[®], Serax[®], and Tranxene[®].

³ Includes Equanil[®], meprobamate, and Miltown[®].

⁴ Includes Flexeril[®] and Soma[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.90D Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Standard Errors of Percentages, 2012 and 2013

Tranquilizer	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
Klonopin [®] or Clonazepam	0.09	0.11	0.07	0.06	0.19	0.20	0.11	0.14
Xanax [®] , Alprazolam, Ativan [®] , or Lorazepam	0.16	0.17	0.13	0.11	0.30	0.29	0.20	0.21
Valium [®] or Diazepam	0.17	0.17	0.09	0.06	0.21	0.20	0.21	0.21
Atarax [®]	0.02	0.02	0.02	0.02	0.04	0.03	0.03	0.03
BuSpar [®]	0.03	0.03	0.02	0.02	0.06	0.07	0.04	0.03
Equanil [®]	0.01	0.01	0.02	0.01	0.03	0.03	0.01	0.01
Flexeril [®]	0.07	0.07	0.03	0.03	0.11	0.11	0.09	0.09
Librium [®]	0.04	0.04	0.02	0.02	0.04	0.04	0.05	0.05
Limbitrol [®]	0.01	0.00	0.02	0.02	0.03	0.02	0.01	0.01
Meprobamate	0.02	0.02	0.02	0.01	0.03	0.02	0.02	0.02
Miltown [®]	0.02	0.01	0.01	0.01	0.03	0.02	0.03	0.01
Rohypnol [®]	0.02	0.03	0.03	0.02	0.05	0.04	0.02	0.03
Serax [®]	0.01	0.01	0.02	0.01	0.03	0.02	0.00	0.01
Soma [®]	0.08	0.07	0.04	0.05	0.15	0.13	0.09	0.09
Tranxene [®]	0.01	0.01	0.01	0.01	0.03	0.02	0.02	0.01
Vistaril [®]	0.02	0.02	0.02	0.02	0.04	0.05	0.02	0.03
SELECTED GROUPS OF DRUGS¹								
Benzodiazepines ²	0.21	0.22	0.15	0.12	0.31	0.31	0.25	0.27
Meprobamate Products ³	0.02	0.02	0.02	0.01	0.04	0.03	0.03	0.03
Muscle Relaxants ⁴	0.10	0.09	0.05	0.06	0.17	0.15	0.12	0.12

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Tranquilizers module but fall into this category.

² Includes Klonopin[®] or clonazepam, Xanax[®], alprazolam, Ativan[®] or lorazepam, Valium[®] or diazepam, Librium[®], Limbitrol[®], Rohypnol[®], Serax[®], and Tranxene[®].

³ Includes Equanil[®], meprobamate, and Miltown[®].

⁴ Includes Flexeril[®] and Soma[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.90P Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Tranquilizer	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
Klonopin [®] or Clonazepam	0.4898		0.0984		0.4512		0.3093	
Xanax [®] , Alprazolam, Ativan [®] , or Lorazepam	0.7199		0.0082		0.4290		0.9964	
Valium [®] or Diazepam	0.2574		0.0005		0.1183		0.4458	
Atarax [®]	0.9709		0.7280		0.2061		0.7559	
BuSpar [®]	0.1256		0.4027		0.6828		0.1530	
Equanil [®]	0.3796		0.3263		0.6296		0.3742	
Flexeril [®]	0.2515		0.4629		0.4142		0.3297	
Librium [®]	0.6908		0.8189		0.9741		0.6945	
Limbitrol [®]	0.2634		0.8326		0.4678		0.4014	
Meprobamate	0.8790		0.3055		0.3305		0.8978	
Miltown [®]	0.1516		0.6113		0.6167		0.1833	
Rohypnol [®]	0.3024		0.0870		0.8399		0.1971	
Serax [®]	0.7235		0.2429		0.3818		0.2192	
Soma [®]	0.1459		0.6802		0.0505		0.2868	
Tranxene [®]	0.3451		0.8564		0.8061		0.3679	
Vistaril [®]	0.3911		0.4818		0.8436		0.3652	
SELECTED GROUPS OF DRUGS¹								
Benzodiazepines ²	0.8075		0.0013		0.1237		0.8040	
Meprobamate Products ³	0.6678		0.1704		0.8079		0.7779	
Muscle Relaxants ⁴	0.1570		0.8167		0.0255		0.3355	

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Tranquilizers module but fall into this category.

² Includes Klonopin[®] or clonazepam, Xanax[®], alprazolam, Ativan[®] or lorazepam, Valium[®] or diazepam, Librium[®], Limbitrol[®], Rohypnol[®], Serax[®], and Tranxene[®].

³ Includes Equanil[®], meprobamate, and Miltown[®].

⁴ Includes Flexeril[®] and Soma[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.91C Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2012 and 2013

Stimulant	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
Methamphetamine, Desoxyn [®] , or Methedrine ¹	426	393	15	14	62	57	418	386
Prescription Diet Pills ²	343	326	14	12	45	41	338	326
Ritalin [®] or Methylphenidate	239	265	21	19	72	74	219	255
Cylert [®]	33	50	3	4	8	5	32	50
Dexedrine [®]	172	189	5	4	17	16	171	188
Dextroamphetamine	93	128	5	10	23	23	90	125
Didrex [®]	31	22	2	2	4	9	30	20
Eskatrol [®]	61	28	2	1	6	5	61	28
Ionamin [®]	95	69	2	6	2	4	95	69
Mazanor [®]	13	8	5	2	7	2	10	8
Obedrin-LA [®]	11	6	4	2	*	5	10	*
Plegine [®]	13	12	2	3	6	2	12	11
Preludin [®]	70	51	4	*	8	7	70	50
Sanorex [®]	12	*	6	1	3	2	10	*
Tenuate [®]	52	34	1	2	4	5	52	34
SELECTED GROUPS OF DRUGS³								
Amphetamine, Dextroamphetamine, and Phentermine Products ⁴	379	363	21	20	67	71	367	357
Mazindol Products ⁵	18	9	7	3	7	3	15	8
Methylphenidate or Dexmethylphenidate Products ⁶	239	267	21	20	73	74	220	257

*Low precision; no estimate reported.

¹ Also known as crank, crystal, ice, or speed. Estimates of Use of Methamphetamine, Desoxyn[®], or Methedrine include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Respondents were asked about their use of prescription diet pills and were given the following as examples: amphetamines, Benzedrine[®], Biphentamine[®], Fastin[®], or phentermine. However, respondents were not given an exhaustive list of examples of prescription diet pills.

³ Includes other-specify drug responses that are not asked about explicitly in the Stimulants module but fall into this category.

⁴ Includes prescription diet pills, Dexedrine[®], dextroamphetamine, and Ionamin[®].

⁵ Includes Mazanor[®] and Sanorex[®].

⁶ Includes Ritalin[®], other products containing methylphenidate, and products containing dexmethylphenidate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.91D Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Standard Errors of Percentages, 2012 and 2013

Stimulant	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
Methamphetamine, Desoxyn [®] , or Methedrine ¹	0.16	0.15	0.06	0.06	0.18	0.17	0.21	0.19
Prescription Diet Pills ²	0.13	0.12	0.06	0.05	0.13	0.12	0.17	0.16
Ritalin [®] or Methylphenidate	0.09	0.10	0.08	0.08	0.21	0.21	0.11	0.13
Cylert [®]	0.01	0.02	0.01	0.01	0.02	0.01	0.02	0.02
Dexedrine [®]	0.07	0.07	0.02	0.02	0.05	0.05	0.09	0.09
Dextroamphetamine	0.04	0.05	0.02	0.04	0.07	0.07	0.04	0.06
Didrex [®]	0.01	0.01	0.01	0.01	0.01	0.03	0.02	0.01
Eskatrol [®]	0.02	0.01	0.01	0.01	0.02	0.02	0.03	0.01
Ionamin [®]	0.04	0.03	0.01	0.02	0.00	0.01	0.05	0.03
Mazanor [®]	0.01	0.00	0.02	0.01	0.02	0.00	0.01	0.00
Obedrin-LA [®]	0.00	0.00	0.01	0.01	*	0.01	0.00	*
Plegine [®]	0.01	0.00	0.01	0.01	0.02	0.00	0.01	0.01
Preludin [®]	0.03	0.02	0.01	*	0.02	0.02	0.03	0.02
Sanorex [®]	0.00	*	0.02	0.01	0.01	0.01	0.01	*
Tenuate [®]	0.02	0.01	0.00	0.01	0.01	0.01	0.03	0.02
SELECTED GROUPS OF DRUGS³								
Amphetamine, Dextroamphetamine, and Phentermine Products ⁴	0.15	0.14	0.08	0.08	0.19	0.20	0.18	0.18
Mazindol Products ⁵	0.01	0.00	0.03	0.01	0.02	0.01	0.01	0.00
Methylphenidate or Dexmethylphenidate Products ⁶	0.09	0.10	0.09	0.08	0.21	0.21	0.11	0.13

*Low precision; no estimate reported.

¹ Also known as crank, crystal, ice, or speed. Estimates of Use of Methamphetamine, Desoxyn[®], or Methedrine include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Respondents were asked about their use of prescription diet pills and were given the following as examples: amphetamines, Benzedrine[®], Biphetamine[®], Fastin[®], or phentermine. However, respondents were not given an exhaustive list of examples of prescription diet pills.

³ Includes other-specify drug responses that are not asked about explicitly in the Stimulants module but fall into this category.

⁴ Includes prescription diet pills, Dexedrine[®], dextroamphetamine, and Ionamin[®].

⁵ Includes Mazanor[®] and Sanorex[®].

⁶ Includes Ritalin[®], other products containing methylphenidate, and products containing dexmethylphenidate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.91P Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Stimulant	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
Methamphetamine, Desoxyn [®] , or Methedrine ¹	0.8425		0.2150		0.5278		0.9306	
Prescription Diet Pills ²	0.7750		0.1033		0.0194		0.9566	
Ritalin [®] or Methylphenidate	0.3545		0.3747		0.8109		0.3247	
Cylert [®]	0.5683		0.9679		0.4747		0.4916	
Dexedrine [®]	0.3752		0.5913		0.1933		0.3059	
Dextroamphetamine	0.1766		0.2198		0.3826		0.1380	
Didrex [®]	0.5721		0.9722		0.1630		0.3428	
Eskatrol [®]	0.3602		0.2041		0.9025		0.3745	
Ionamin [®]	0.9331		0.4588		0.0819		0.9780	
Mazanor [®]	0.1674		0.1834		0.2377		0.6134	
Obedrin-LA [®]	0.6717		0.6059		*		*	
Plegine [®]	0.9787		0.8404		0.5368		0.8209	
Preludin [®]	0.1881		*		0.6027		0.2305	
Sanorex [®]	*		0.0277		0.7627		*	
Tenuate [®]	0.9714		0.3759		0.7992		0.9608	
SELECTED GROUPS OF DRUGS³								
Amphetamine, Dextroamphetamine, and Phentermine Products ⁴	0.5130		0.2674		0.5134		0.4045	
Mazindol Products ⁵	0.0085		0.0172		0.2932		0.1155	
Methylphenidate or Dexmethylphenidate Products ⁶	0.3452		0.4869		0.7959		0.3264	

*Low precision; no estimate reported.

¹ Also known as crank, crystal, ice, or speed. Estimates of Use of Methamphetamine, Desoxyn[®], or Methedrine include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Respondents were asked about their use of prescription diet pills and were given the following as examples: amphetamines, Benzedrine[®], Biphetamine[®], Fastin[®], or phentermine. However, respondents were not given an exhaustive list of examples of prescription diet pills.

³ Includes other-specify drug responses that are not asked about explicitly in the Stimulants module but fall into this category.

⁴ Includes prescription diet pills, Dexedrine[®], dextroamphetamine, and Ionamin[®].

⁵ Includes Mazanor[®] and Sanorex[®].

⁶ Includes Ritalin[®], other products containing methylphenidate, and products containing dexmethylphenidate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.92C Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2012 and 2013

Sedative	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
Methaqualone, Sopor [®] , or Quaalude [®]	325	312	6	6	12	14	325	312
Barbiturates ¹	225	198	6	4	9	17	224	198
Restoril [®] or Temazepam	166	125	10	8	22	22	165	121
Amytal [®]	36	36	4	4	4	8	35	35
Butisol [®]	90	9	3	2	6	5	89	7
Chloral Hydrate	35	39	3	3	6	6	35	38
Dalmane [®]	48	64	3	2	9	8	47	64
Halcion [®]	84	93	4	3	7	9	84	92
Phenobarbital	117	103	6	5	8	17	117	101
Placidyl [®]	132	109	2	2	2	5	132	109
Tuinal [®]	112	121	2	2	4	4	111	121
SELECTED GROUPS OF DRUGS²								
Temazepam, Flurazepam, or Triazolam ³	189	162	11	9	23	24	186	158
Any Barbiturates ⁴	239	217	10	7	13	21	238	216

*Low precision; no estimate reported.

¹ Respondents were asked about their use of barbiturates and were given the following as examples: Nembutal[®], pentobarbital, Seconal[®], secobarbital, or butalbital. However, respondents were not given an exhaustive list of examples of barbiturates.

² Includes other-specify drug responses that are not asked about explicitly in the Sedatives module but fall into this category.

³ Includes Restoril[®], temazepam, Dalmane[®], or Halcion[®].

⁴ Includes barbiturates, Amytal[®], Butisol[®], phenobarbital, and Tuinal[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.92D Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Standard Errors of Percentages, 2012 and 2013

Sedative	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
Methaqualone, Sopor [®] , or Quaalude [®]	0.12	0.12	0.03	0.02	0.03	0.04	0.16	0.15
Barbiturates ¹	0.09	0.08	0.02	0.01	0.03	0.05	0.11	0.10
Restoril [®] or Temazepam	0.06	0.05	0.04	0.03	0.06	0.06	0.08	0.06
Amytal [®]	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.02
Butisol [®]	0.03	0.00	0.01	0.01	0.02	0.01	0.04	0.00
Chloral Hydrate	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.02
Dalmane [®]	0.02	0.02	0.01	0.01	0.02	0.02	0.02	0.03
Halcion [®]	0.03	0.04	0.01	0.01	0.02	0.03	0.04	0.05
Phenobarbital	0.05	0.04	0.03	0.02	0.02	0.05	0.06	0.05
Placidyl [®]	0.05	0.04	0.01	0.01	0.01	0.01	0.07	0.05
Tuinal [®]	0.04	0.05	0.01	0.01	0.01	0.01	0.06	0.06
SELECTED GROUPS OF DRUGS²								
Temazepam, Flurazepam, or Triazolam ³	0.07	0.06	0.04	0.04	0.07	0.07	0.09	0.08
Any Barbiturates ⁴	0.09	0.08	0.04	0.03	0.04	0.06	0.12	0.11

*Low precision; no estimate reported.

¹ Respondents were asked about their use of barbiturates and were given the following as examples: Nembutal[®], pentobarbital, Seconal[®], secobarbital, or butalbital. However, respondents were not given an exhaustive list of examples of barbiturates.

² Includes other-specify drug responses that are not asked about explicitly in the Sedatives module but fall into this category.

³ Includes Restoril[®], temazepam, Dalmane[®], or Halcion[®].

⁴ Includes barbiturates, Amytal[®], Butisol[®], phenobarbital, and Tuinal[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 1.92P Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Sedative	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
Methaqualone, Sopor [®] , or Quaalude [®]	0.3732		0.6902		0.2138		0.3405	
Barbiturates ¹	0.7179		0.0542		0.0444		0.6491	
Restoril [®] or Temazepam	0.0972		0.0310		0.9524		0.1181	
Amytal [®]	0.9466		0.5422		0.2012		0.9155	
Butisol [®]	0.1263		0.6105		0.4087		0.1497	
Chloral Hydrate	0.9152		0.6756		0.6710		0.8354	
Dalmane [®]	0.3406		0.2518		0.8397		0.2982	
Halcion [®]	0.8707		0.7430		0.5184		0.9086	
Phenobarbital	0.9727		0.4635		0.0277		0.8365	
Placidyl [®]	0.4665		0.6216		0.6564		0.4594	
Tuinal [®]	0.8586		0.4105		0.4731		0.8621	
SELECTED GROUPS OF DRUGS²								
Temazepam, Flurazepam, or Triazolam ³	0.2845		0.0495		0.8268		0.3131	
Any Barbiturates ⁴	0.5743		0.1056		0.0592		0.5093	

*Low precision; no estimate reported.

¹ Respondents were asked about their use of barbiturates and were given the following as examples: Nembutal[®], pentobarbital, Seconal[®], secobarbital, or butalbital. However, respondents were not given an exhaustive list of examples of barbiturates.

² Includes other-specify drug responses that are not asked about explicitly in the Sedatives module but fall into this category.

³ Includes Restoril[®], temazepam, Dalmane[®], or Halcion[®].

⁴ Includes barbiturates, Amytal[®], Butisol[®], phenobarbital, and Tuinal[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.