

**Table 2.1A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Numbers in Thousands, 2012 and 2013**

<b>Substance</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	173,654	175,260	82,829	80,846	69,497 <sup>a</sup>	66,879
Cigarettes	161,057	162,138	67,802	66,256	57,525	55,778
Smokeless Tobacco	45,936	46,139	12,040	11,866	9,036	8,813
Cigars	88,596	89,117	25,647	24,933	13,417	12,438
Pipe Tobacco <sup>2</sup>	34,604	34,173	--	--	2,530	2,271
<b>ALCOHOL</b>	214,013	213,794	173,573	173,966	135,502	136,868
Binge Alcohol Use <sup>3</sup>	--	--	--	--	59,685	60,064
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	17,021	16,541

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.1B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Percentages, 2012 and 2013**

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	66.8	66.8	31.9 <sup>a</sup>	30.8	26.7 <sup>b</sup>	25.5
Cigarettes	61.9	61.8	26.1	25.3	22.1 <sup>a</sup>	21.3
Smokeless Tobacco	17.7	17.6	4.6	4.5	3.5	3.4
Cigars	34.1	34.0	9.9	9.5	5.2 <sup>a</sup>	4.7
Pipe Tobacco <sup>2</sup>	13.3	13.0	--	--	1.0	0.9
<b>ALCOHOL</b>	82.3 <sup>a</sup>	81.5	66.7	66.3	52.1	52.2
Binge Alcohol Use <sup>3</sup>	--	--	--	--	23.0	22.9
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	6.5	6.3

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.2A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Numbers in Thousands, 2012 and 2013**

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	512	454	293	263	128	107
Cigarettes	383	317	222	184	93	71
Smokeless Tobacco	154	136	81	75	36	25
Cigars	136	130	81	83	31	18
Pipe Tobacco <sup>2</sup>	56	50	--	--	24	17
<b>ALCOHOL</b>	794	730	546	470	179	167
Binge Alcohol Use <sup>3</sup>	--	--	--	--	73	67
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	16	10

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.2B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Percentages, 2012 and 2013**

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	6.4	5.7	3.6	3.3	1.6	1.3
Cigarettes	4.8	4.0	2.8	2.3	1.2	0.9
Smokeless Tobacco	1.9	1.7	1.0	0.9	0.4	0.3
Cigars	1.7	1.6	1.0	1.0	0.4	0.2
Pipe Tobacco <sup>2</sup>	0.7	0.6	--	--	0.3	0.2
<b>ALCOHOL</b>	9.9	9.2	6.8	5.9	2.2	2.1
Binge Alcohol Use <sup>3</sup>	--	--	--	--	0.9	0.8
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.2	0.1

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.3A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Numbers in Thousands, 2012 and 2013**

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	1,677	1,577	1,140	1,067	529	544
Cigarettes	1,339	1,224	859	774	389	367
Smokeless Tobacco	466	456	280	319	143	154
Cigars	645 <sup>a</sup>	540	463 <sup>b</sup>	360	141	122
Pipe Tobacco <sup>2</sup>	165	165	--	--	41	47
<b>ALCOHOL</b>	2,658 <sup>a</sup>	2,427	2,116 <sup>a</sup>	1,896	936 <sup>a</sup>	806
Binge Alcohol Use <sup>3</sup>	--	--	--	--	454	385
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	50	57

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.3B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Percentages, 2012 and 2013**

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	20.0	18.5	13.6	12.5	6.3	6.4
Cigarettes	16.0 <sup>a</sup>	14.4	10.2	9.1	4.6	4.3
Smokeless Tobacco	5.6	5.4	3.3	3.7	1.7	1.8
Cigars	7.7 <sup>a</sup>	6.3	5.5 <sup>b</sup>	4.2	1.7	1.4
Pipe Tobacco <sup>2</sup>	2.0	1.9	--	--	0.5	0.6
<b>ALCOHOL</b>	31.7 <sup>b</sup>	28.5	25.2 <sup>b</sup>	22.3	11.1 <sup>a</sup>	9.5
Binge Alcohol Use <sup>3</sup>	--	--	--	--	5.4 <sup>a</sup>	4.5
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.6	0.7

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.4A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Numbers in Thousands, 2012 and 2013**

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	3,165 <sup>a</sup>	2,891	2,368 <sup>a</sup>	2,131	1,496 <sup>b</sup>	1,300
Cigarettes	2,616 <sup>b</sup>	2,362	1,868 <sup>b</sup>	1,603	1,152 <sup>b</sup>	959
Smokeless Tobacco	966	908	611	617	336	309
Cigars	1,608 <sup>b</sup>	1,400	1,139 <sup>a</sup>	981	473	436
Pipe Tobacco <sup>2</sup>	399 <sup>a</sup>	303	--	--	100	96
<b>ALCOHOL</b>	4,615	4,512	3,893	3,766	2,102 <sup>a</sup>	1,903
Binge Alcohol Use <sup>3</sup>	--	--	--	--	1,273 <sup>a</sup>	1,104
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	260	226

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.4B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Percentages, 2012 and 2013**

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	37.3 <sup>b</sup>	34.4	27.9 <sup>b</sup>	25.4	17.6 <sup>b</sup>	15.5
Cigarettes	30.8 <sup>b</sup>	28.1	22.0 <sup>b</sup>	19.1	13.6 <sup>b</sup>	11.4
Smokeless Tobacco	11.4	10.8	7.2	7.3	4.0	3.7
Cigars	19.0 <sup>b</sup>	16.7	13.4 <sup>a</sup>	11.7	5.6	5.2
Pipe Tobacco <sup>2</sup>	4.7 <sup>a</sup>	3.6	--	--	1.2	1.1
<b>ALCOHOL</b>	54.4	53.7	45.9	44.8	24.8 <sup>a</sup>	22.7
Binge Alcohol Use <sup>3</sup>	--	--	--	--	15.0 <sup>a</sup>	13.1
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	3.1	2.7

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.5A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Numbers in Thousands, 2012 and 2013**

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	7,865	7,478	6,248	5,952	4,634	4,479
Cigarettes	6,878	6,420	5,131	4,792	3,745	3,571
Smokeless Tobacco	2,313	2,394	1,255	1,393	710 <sup>a</sup>	849
Cigars	4,557	4,273	3,188	2,999	1,582	1,462
Pipe Tobacco <sup>2</sup>	1,182	1,204	--	--	276 <sup>b</sup>	400
<b>ALCOHOL</b>	9,849	9,581	8,817	8,559	6,077	5,774
Binge Alcohol Use <sup>3</sup>	--	--	--	--	4,053	3,843
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	1,325 <sup>a</sup>	1,120

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.5B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Percentages, 2012 and 2013**

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	59.3 <sup>a</sup>	56.7	47.1	45.1	34.9	33.9
Cigarettes	51.8 <sup>b</sup>	48.7	38.7 <sup>a</sup>	36.3	28.2	27.1
Smokeless Tobacco	17.4	18.1	9.5	10.6	5.4 <sup>a</sup>	6.4
Cigars	34.3	32.4	24.0	22.7	11.9	11.1
Pipe Tobacco <sup>2</sup>	8.9	9.1	--	--	2.1 <sup>b</sup>	3.0
<b>ALCOHOL</b>	74.2	72.6	66.4	64.9	45.8	43.8
Binge Alcohol Use <sup>3</sup>	--	--	--	--	30.5	29.1
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	10.0 <sup>a</sup>	8.5

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.6A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Numbers in Thousands, 2012 and 2013**

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	15,061	15,172	10,643	10,645	8,533	8,403
Cigarettes	13,714	13,721	9,063	8,958	7,263	7,085
Smokeless Tobacco	4,578	4,671	1,854	1,839	1,201	1,170
Cigars	8,933	9,106	4,332	4,215	2,134	2,010
Pipe Tobacco <sup>2</sup>	2,239	2,393	--	--	349	357
<b>ALCOHOL</b>	19,345	19,567	17,961	18,148	14,750	14,964
Binge Alcohol Use <sup>3</sup>	--	--	--	--	9,609	9,343
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	3,061	2,819

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.6B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Percentages, 2012 and 2013**

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	70.7	70.3	49.9	49.3	40.0	38.9
Cigarettes	64.3	63.6	42.5	41.5	34.1	32.8
Smokeless Tobacco	21.5	21.6	8.7	8.5	5.6	5.4
Cigars	41.9	42.2	20.3	19.5	10.0	9.3
Pipe Tobacco <sup>2</sup>	10.5	11.1	--	--	1.6	1.7
<b>ALCOHOL</b>	90.8	90.6	84.3	84.1	69.2	69.3
Binge Alcohol Use <sup>3</sup>	--	--	--	--	45.1 <sup>a</sup>	43.3
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	14.4 <sup>a</sup>	13.1

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.7A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Numbers in Thousands, 2012 and 2013**

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	27,131	27,926	16,479	17,361	13,824	14,316
Cigarettes	25,468	26,160	14,110	14,760	12,033	12,312
Smokeless Tobacco	8,157	7,987	2,290	2,374	1,722	1,736
Cigars	15,869	16,641	5,396	5,951	2,709	2,895
Pipe Tobacco <sup>2</sup>	3,370	3,304	--	--	402	349
<b>ALCOHOL</b>	33,463	33,703	29,722	30,065	23,792 <sup>a</sup>	24,686
Binge Alcohol Use <sup>3</sup>	--	--	--	--	13,126 <sup>a</sup>	13,983
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	3,465 <sup>a</sup>	4,058

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.7B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Percentages, 2012 and 2013**

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	73.5	74.8	44.6	46.5	37.5	38.3
Cigarettes	69.0	70.0	38.2	39.5	32.6	33.0
Smokeless Tobacco	22.1	21.4	6.2	6.4	4.7	4.6
Cigars	43.0	44.6	14.6	15.9	7.3	7.8
Pipe Tobacco <sup>2</sup>	9.1	8.8	--	--	1.1	0.9
<b>ALCOHOL</b>	90.7	90.2	80.5	80.5	64.5	66.1
Binge Alcohol Use <sup>3</sup>	--	--	--	--	35.6	37.4
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	9.4	10.9

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.8A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Numbers in Thousands, 2012 and 2013**

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	118,243	119,762	45,659 <sup>a</sup>	43,427	40,355 <sup>b</sup>	37,730
Cigarettes	110,657	111,934	36,549	35,185	32,849	31,412
Smokeless Tobacco	29,302	29,587	5,670	5,251	4,888	4,572
Cigars	56,847	57,027	11,047	10,345	6,346	5,494
Pipe Tobacco <sup>2</sup>	27,193	26,754	--	--	1,338	1,004
<b>ALCOHOL</b>	143,288	143,274	110,518	111,061	87,666	88,567
Binge Alcohol Use <sup>3</sup>	--	--	--	--	31,098	31,339
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	8,844	8,252

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.8B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Percentages, 2012 and 2013**

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	72.3	72.4	27.9 <sup>a</sup>	26.3	24.7 <sup>b</sup>	22.8
Cigarettes	67.6	67.7	22.3	21.3	20.1	19.0
Smokeless Tobacco	17.9	17.9	3.5	3.2	3.0	2.8
Cigars	34.7	34.5	6.8	6.3	3.9 <sup>a</sup>	3.3
Pipe Tobacco <sup>2</sup>	16.6	16.2	--	--	0.8	0.6
<b>ALCOHOL</b>	87.6	86.6	67.5	67.2	53.6	53.6
Binge Alcohol Use <sup>3</sup>	--	--	--	--	19.0	19.0
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	5.4	5.0

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.9A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Numbers in Thousands, 2012 and 2013**

<b>Substance</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	93,877	94,449	49,600 <sup>a</sup>	47,773	41,522 <sup>a</sup>	39,549
Cigarettes	84,479	84,457	36,941	35,858	31,021	30,041
Smokeless Tobacco	38,912	38,921	11,130	10,806	8,496	8,211
Cigars	65,531	64,644	19,909	18,848	10,756 <sup>a</sup>	9,755
Pipe Tobacco <sup>2</sup>	30,116	29,642	--	--	2,038	1,878
<b>ALCOHOL</b>	107,153	107,186	88,046	88,185	71,187	72,605
Binge Alcohol Use <sup>3</sup>	--	--	--	--	38,277	38,379
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	12,469	12,093

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.9B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Percentages, 2012 and 2013**

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	74.6	74.3	39.4 <sup>b</sup>	37.6	33.0 <sup>b</sup>	31.1
Cigarettes	67.1	66.4	29.3	28.2	24.6	23.6
Smokeless Tobacco	30.9	30.6	8.8	8.5	6.7	6.5
Cigars	52.1	50.9	15.8 <sup>a</sup>	14.8	8.5 <sup>a</sup>	7.7
Pipe Tobacco <sup>2</sup>	23.9	23.3	--	--	1.6	1.5
<b>ALCOHOL</b>	85.1	84.3	69.9	69.4	56.5	57.1
Binge Alcohol Use <sup>3</sup>	--	--	--	--	30.4	30.2
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	9.9	9.5

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.10A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Numbers in Thousands, 2012 and 2013**

<b>Substance</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	79,777	80,810	33,229	33,073	27,976	27,331
Cigarettes	76,578	77,682	30,861	30,398	26,504	25,737
Smokeless Tobacco	7,025	7,218	910	1,060	540	603
Cigars	23,065 <sup>a</sup>	24,473	5,738	6,085	2,661	2,683
Pipe Tobacco <sup>2</sup>	4,488	4,532	--	--	492	392
<b>ALCOHOL</b>	106,860	106,608	85,526	85,781	64,315	64,263
Binge Alcohol Use <sup>3</sup>	--	--	--	--	21,409	21,685
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	4,553	4,448

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 2.10B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Percentages, 2012 and 2013**

<b>Substance</b>	<b>Lifetime (2012)</b>	<b>Lifetime (2013)</b>	<b>Past Year (2012)</b>	<b>Past Year (2013)</b>	<b>Past Month (2012)</b>	<b>Past Month (2013)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	59.5	59.7	24.8	24.4	20.9	20.2
Cigarettes	57.1	57.4	23.0	22.5	19.8	19.0
Smokeless Tobacco	5.2	5.3	0.7	0.8	0.4	0.4
Cigars	17.2	18.1	4.3	4.5	2.0	2.0
Pipe Tobacco <sup>2</sup>	3.3	3.4	--	--	0.4	0.3
<b>ALCOHOL</b>	79.6	78.8	63.7	63.4	47.9	47.5
Binge Alcohol Use <sup>3</sup>	--	--	--	--	16.0	16.0
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	3.4	3.3

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.