

Table 2.1C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	939	942	933	875	882	835
Cigarettes	956	966	890	830	839	798
Smokeless Tobacco	705	727	325	334	298	300
Cigars	882	922	511	493	388	354
Pipe Tobacco ²	679	768	--	--	184	151
ALCOHOL	675	801	955	1,007	1,016	1,076
Binge Alcohol Use ³	--	--	--	--	797	804
Heavy Alcohol Use ³	--	--	--	--	448	438

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.1D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Percentages, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.36	0.36	0.36	0.33	0.34	0.32
Cigarettes	0.37	0.37	0.34	0.32	0.32	0.30
Smokeless Tobacco	0.27	0.28	0.13	0.13	0.11	0.11
Cigars	0.34	0.35	0.20	0.19	0.15	0.14
Pipe Tobacco ²	0.26	0.29	--	--	0.07	0.06
ALCOHOL	0.26	0.31	0.37	0.38	0.39	0.41
Binge Alcohol Use ³	--	--	--	--	0.31	0.31
Heavy Alcohol Use ³	--	--	--	--	0.17	0.17

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.1P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.9709		0.0278		0.0050	
Cigarettes	0.7810		0.0669		0.0411	
Smokeless Tobacco	0.8283		0.5328		0.4649	
Cigars	0.8196		0.1812		0.0351	
Pipe Tobacco ²	0.4672		--		0.2396	
ALCOHOL	0.0373		0.3796		0.9167	
Binge Alcohol Use ³	--		--		0.8875	
Heavy Alcohol Use ³	--		--		0.3092	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.2C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Numbers in Thousands, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	31	32	24	23	15	16
Cigarettes	26	26	20	20	12	13
Smokeless Tobacco	17	17	12	13	7	6
Cigars	16	15	11	12	7	5
Pipe Tobacco ²	10	10	--	--	7	6
ALCOHOL	42	41	36	31	20	17
Binge Alcohol Use ³	--	--	--	--	13	12
Heavy Alcohol Use ³	--	--	--	--	5	4

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.2D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Percentages, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.37	0.38	0.29	0.28	0.18	0.20
Cigarettes	0.32	0.32	0.25	0.24	0.16	0.16
Smokeless Tobacco	0.21	0.21	0.15	0.16	0.09	0.07
Cigars	0.20	0.18	0.14	0.15	0.08	0.07
Pipe Tobacco ²	0.12	0.13	--	--	0.08	0.08
ALCOHOL	0.49	0.47	0.42	0.37	0.24	0.22
Binge Alcohol Use ³	--	--	--	--	0.16	0.15
Heavy Alcohol Use ³	--	--	--	--	0.07	0.05

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.2P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.2153		0.4162		0.3622	
Cigarettes	0.0891		0.2206		0.2381	
Smokeless Tobacco	0.4754		0.7570		0.2535	
Cigars	0.8330		0.8867		0.1550	
Pipe Tobacco ²	0.7106		--		0.4106	
ALCOHOL	0.3071		0.1287		0.6888	
Binge Alcohol Use ³	--		--		0.7694	
Heavy Alcohol Use ³	--		--		0.3790	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.3C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Numbers in Thousands, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	56	59	45	47	30	33
Cigarettes	50	49	40	38	25	26
Smokeless Tobacco	30	32	22	27	16	17
Cigars	35	31	30	25	15	14
Pipe Tobacco ²	18	17	--	--	8	10
ALCOHOL	77	74	67	59	44	39
Binge Alcohol Use ³	--	--	--	--	29	27
Heavy Alcohol Use ³	--	--	--	--	8	9

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.3D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Percentages, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.58	0.59	0.50	0.51	0.35	0.37
Cigarettes	0.54	0.52	0.45	0.43	0.29	0.30
Smokeless Tobacco	0.35	0.36	0.27	0.30	0.19	0.20
Cigars	0.40	0.36	0.35	0.29	0.18	0.16
Pipe Tobacco ²	0.21	0.20	--	--	0.09	0.12
ALCOHOL	0.72	0.74	0.67	0.63	0.47	0.44
Binge Alcohol Use ³	--	--	--	--	0.33	0.31
Heavy Alcohol Use ³	--	--	--	--	0.10	0.11

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.3P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.0782		0.1348		0.8696	
Cigarettes	0.0301		0.0626		0.4298	
Smokeless Tobacco	0.6875		0.2900		0.7227	
Cigars	0.0121		0.0028		0.3100	
Pipe Tobacco ²	0.9284		--		0.6375	
ALCOHOL	0.0023		0.0013		0.0105	
Binge Alcohol Use ³	--		--		0.0487	
Heavy Alcohol Use ³	--		--		0.5845	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.4C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Numbers in Thousands, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	83	77	70	67	54	51
Cigarettes	77	68	62	54	49	43
Smokeless Tobacco	43	42	32	35	24	24
Cigars	56	55	46	44	28	29
Pipe Tobacco ²	32	22	--	--	14	13
ALCOHOL	102	101	92	93	67	65
Binge Alcohol Use ³	--	--	--	--	51	48
Heavy Alcohol Use ³	--	--	--	--	23	22

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.4D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Percentages, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.73	0.74	0.68	0.67	0.56	0.55
Cigarettes	0.72	0.69	0.63	0.58	0.51	0.48
Smokeless Tobacco	0.47	0.47	0.36	0.39	0.28	0.28
Cigars	0.58	0.58	0.50	0.49	0.32	0.33
Pipe Tobacco ²	0.36	0.26	--	--	0.16	0.15
ALCOHOL	0.78	0.79	0.77	0.80	0.64	0.66
Binge Alcohol Use ³	--	--	--	--	0.52	0.52
Heavy Alcohol Use ³	--	--	--	--	0.27	0.25

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.4P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.0050		0.0068		0.0061	
Cigarettes	0.0053		0.0005		0.0018	
Smokeless Tobacco	0.3698		0.7862		0.4693	
Cigars	0.0064		0.0139		0.3868	
Pipe Tobacco ²	0.0130		--		0.8639	
ALCOHOL	0.5294		0.3479		0.0230	
Binge Alcohol Use ³	--		--		0.0113	
Heavy Alcohol Use ³	--		--		0.3042	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.5C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Numbers in Thousands, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	220	182	187	155	152	126
Cigarettes	202	160	162	133	135	111
Smokeless Tobacco	88	85	64	64	44	48
Cigars	151	131	122	108	75	68
Pipe Tobacco ²	66	64	--	--	24	36
ALCOHOL	277	224	270	212	222	168
Binge Alcohol Use ³	--	--	--	--	176	136
Heavy Alcohol Use ³	--	--	--	--	85	63

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.5D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Percentages, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.80	0.77	0.78	0.75	0.75	0.74
Cigarettes	0.83	0.80	0.78	0.75	0.73	0.71
Smokeless Tobacco	0.59	0.56	0.45	0.44	0.32	0.34
Cigars	0.73	0.70	0.66	0.64	0.45	0.46
Pipe Tobacco ²	0.41	0.45	--	--	0.17	0.27
ALCOHOL	0.73	0.70	0.81	0.75	0.90	0.79
Binge Alcohol Use ³	--	--	--	--	0.87	0.76
Heavy Alcohol Use ³	--	--	--	--	0.54	0.42

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.5P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.0127		0.0622		0.3466	
Cigarettes	0.0029		0.0226		0.2427	
Smokeless Tobacco	0.3472		0.0664		0.0167	
Cigars	0.0540		0.1612		0.1897	
Pipe Tobacco ²	0.7073		--		0.0028	
ALCOHOL	0.1065		0.1495		0.0874	
Binge Alcohol Use ³	--		--		0.2125	
Heavy Alcohol Use ³	--		--		0.0261	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.6C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Numbers in Thousands, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	266	286	212	210	186	175
Cigarettes	244	257	188	184	165	157
Smokeless Tobacco	123	135	70	73	58	55
Cigars	197	211	122	123	79	81
Pipe Tobacco ²	90	88	--	--	31	36
ALCOHOL	313	346	301	331	269	294
Binge Alcohol Use ³	--	--	--	--	208	218
Heavy Alcohol Use ³	--	--	--	--	107	101

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.6D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Percentages, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.54	0.55	0.60	0.66	0.60	0.61
Cigarettes	0.58	0.56	0.59	0.63	0.58	0.59
Smokeless Tobacco	0.46	0.50	0.31	0.33	0.26	0.25
Cigars	0.60	0.60	0.45	0.49	0.33	0.36
Pipe Tobacco ²	0.37	0.36	--	--	0.14	0.16
ALCOHOL	0.35	0.37	0.44	0.48	0.56	0.60
Binge Alcohol Use ³	--	--	--	--	0.60	0.63
Heavy Alcohol Use ³	--	--	--	--	0.42	0.39

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.6P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.6127		0.4722		0.1883	
Cigarettes	0.3045		0.2213		0.1237	
Smokeless Tobacco	0.8095		0.6810		0.5635	
Cigars	0.7452		0.2252		0.1480	
Pipe Tobacco ²	0.2789		--		0.9305	
ALCOHOL	0.8034		0.7615		0.8918	
Binge Alcohol Use ³	--		--		0.0323	
Heavy Alcohol Use ³	--		--		0.0197	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.7C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Numbers in Thousands, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	289	302	300	334	293	311
Cigarettes	293	312	288	320	278	302
Smokeless Tobacco	261	253	143	143	129	122
Cigars	309	331	222	238	168	166
Pipe Tobacco ²	180	176	--	--	60	63
ALCOHOL	195	209	256	277	312	322
Binge Alcohol Use ³	--	--	--	--	313	318
Heavy Alcohol Use ³	--	--	--	--	188	221

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.7D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Percentages, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.78	0.81	0.81	0.89	0.79	0.83
Cigarettes	0.79	0.83	0.78	0.86	0.75	0.81
Smokeless Tobacco	0.71	0.68	0.39	0.38	0.35	0.33
Cigars	0.84	0.89	0.60	0.64	0.45	0.44
Pipe Tobacco ²	0.49	0.47	--	--	0.16	0.17
ALCOHOL	0.53	0.56	0.69	0.74	0.85	0.86
Binge Alcohol Use ³	--	--	--	--	0.85	0.85
Heavy Alcohol Use ³	--	--	--	--	0.51	0.59

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.7P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.2451		0.1314		0.4398	
Cigarettes	0.3635		0.2696		0.7389	
Smokeless Tobacco	0.4596		0.7737		0.9713	
Cigars	0.1718		0.1323		0.5106	
Pipe Tobacco ²	0.6673		--		0.5117	
ALCOHOL	0.5898		0.9842		0.1666	
Binge Alcohol Use ³	--		--		0.1053	
Heavy Alcohol Use ³	--		--		0.0522	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.8C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Numbers in Thousands, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	821	828	799	747	765	714
Cigarettes	852	854	755	693	721	681
Smokeless Tobacco	638	667	278	267	261	254
Cigars	788	832	424	411	324	291
Pipe Tobacco ²	644	726	--	--	165	127
ALCOHOL	610	684	874	890	899	956
Binge Alcohol Use ³	--	--	--	--	664	664
Heavy Alcohol Use ³	--	--	--	--	374	344

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.8D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Percentages, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.50	0.50	0.49	0.45	0.47	0.43
Cigarettes	0.52	0.52	0.46	0.42	0.44	0.41
Smokeless Tobacco	0.39	0.40	0.17	0.16	0.16	0.15
Cigars	0.48	0.50	0.26	0.25	0.20	0.18
Pipe Tobacco ²	0.39	0.44	--	--	0.10	0.08
ALCOHOL	0.37	0.41	0.53	0.54	0.55	0.58
Binge Alcohol Use ³	--	--	--	--	0.41	0.40
Heavy Alcohol Use ³	--	--	--	--	0.23	0.21

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.8P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.8251		0.0134		0.0029	
Cigarettes	0.9350		0.0830		0.0651	
Smokeless Tobacco	0.9757		0.2085		0.3080	
Cigars	0.7020		0.1714		0.0385	
Pipe Tobacco ²	0.4551		--		0.0990	
ALCOHOL	0.0910		0.5966		0.9792	
Binge Alcohol Use ³	--		--		0.9227	
Heavy Alcohol Use ³	--		--		0.1891	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.9C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Standard Errors of Numbers in Thousands, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	536	562	655	630	618	612
Cigarettes	577	613	598	571	556	549
Smokeless Tobacco	609	634	307	312	281	282
Cigars	667	704	451	416	351	312
Pipe Tobacco ²	615	683	--	--	170	146
ALCOHOL	380	458	605	641	675	693
Binge Alcohol Use ³	--	--	--	--	625	607
Heavy Alcohol Use ³	--	--	--	--	382	367

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.9D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Standard Errors of Percentages, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.43	0.44	0.52	0.50	0.49	0.48
Cigarettes	0.46	0.48	0.47	0.45	0.44	0.43
Smokeless Tobacco	0.48	0.50	0.24	0.25	0.22	0.22
Cigars	0.53	0.55	0.36	0.33	0.28	0.25
Pipe Tobacco ²	0.49	0.54	--	--	0.14	0.12
ALCOHOL	0.30	0.36	0.48	0.50	0.54	0.55
Binge Alcohol Use ³	--	--	--	--	0.50	0.48
Heavy Alcohol Use ³	--	--	--	--	0.30	0.29

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.9P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: *P* Values from Tests of Differences of Percentages, 2013 Versus 2012

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.6532		0.0090		0.0043	
Cigarettes	0.2996		0.0749		0.0911	
Smokeless Tobacco	0.6658		0.3187		0.3536	
Cigars	0.1005		0.0386		0.0163	
Pipe Tobacco ²	0.3996		--		0.4243	
ALCOHOL	0.0846		0.4155		0.4621	
Binge Alcohol Use ³	--		--		0.7559	
Heavy Alcohol Use ³	--		--		0.3506	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.10C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Standard Errors of Numbers in Thousands, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	683	698	578	568	554	524
Cigarettes	688	693	561	546	539	513
Smokeless Tobacco	296	283	109	108	103	94
Cigars	472	506	211	250	146	161
Pipe Tobacco ²	222	237	--	--	61	35
ALCOHOL	516	598	660	709	686	736
Binge Alcohol Use ³	--	--	--	--	427	463
Heavy Alcohol Use ³	--	--	--	--	203	232

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.10D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Standard Errors of Percentages, 2012 and 2013

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.51	0.52	0.43	0.42	0.41	0.39
Cigarettes	0.51	0.51	0.42	0.40	0.40	0.38
Smokeless Tobacco	0.22	0.21	0.08	0.08	0.08	0.07
Cigars	0.35	0.37	0.16	0.18	0.11	0.12
Pipe Tobacco ²	0.17	0.18	--	--	0.05	0.03
ALCOHOL	0.38	0.44	0.49	0.52	0.51	0.54
Binge Alcohol Use ³	--	--	--	--	0.32	0.34
Heavy Alcohol Use ³	--	--	--	--	0.15	0.17

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 2.10P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Substance	Lifetime (2012)	Lifetime (2013)	Past Year (2012)	Past Year (2013)	Past Month (2012)	Past Month (2013)
TOBACCO PRODUCTS¹	0.6909		0.5933		0.2473	
Cigarettes	0.6177		0.3535		0.1788	
Smokeless Tobacco	0.7396		0.3571		0.6792	
Cigars	0.0720		0.3404		1.0000	
Pipe Tobacco ²	0.9829		--		0.1490	
ALCOHOL	0.1398		0.6240		0.5396	
Binge Alcohol Use ³	--		--		0.8719	
Heavy Alcohol Use ³	--		--		0.6413	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.