

Table 6.11A Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2012 and 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLICIT DRUGS²	5,281	5,566	7,900	8,043	5,407	5,603	5,275	5,360
Marijuana and Hashish	4,476	4,723	6,656	6,905	4,252	4,572	3,471	3,611
Cocaine	678	738	561	414	199	253	212	145
Crack	169	111	146	86	14 ^a	105	115	75
Heroin	64	33	75	68	51	79	145	110
Hallucinogens	469	450	429	466	102	248	134	170
LSD	79	90	75	101	14	13	16	24
PCP	16	3	8	15	4	*	*	14
Ecstasy	287	290	229	238	68	89	44	43
Inhalants	91	106	195	111	68	99	139	180
Nonmedical Use of Psychotherapeutics ^{3,4}	1,473	1,427	2,001	1,921	1,456	1,225	1,901	1,911
Pain Relievers	1,007	968	1,499	1,351	923	824	1,434	1,378
OxyContin [®]	125	113	91	139	41	59	102	181
Tranquilizers	491	442	663	574	486 ^a	279	499	410
Stimulants ⁴	354	421	382	400	244	256	191	288
Methamphetamine ⁴	88	170	161	138	87	106	104	181
Sedatives	35	40	118 ^a	36	54	73	63	102
ILLICIT DRUGS OTHER THAN MARIJUANA²	2,196	2,140	2,737	2,586	1,717	1,686	2,233	2,253

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.11B Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2012 and 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLCIT DRUGS²	31.0	33.7	18.5	18.5	7.1	7.3	4.2	4.3
Marijuana and Hashish	26.3	28.6	15.6	15.9	5.6	6.0	2.8	2.9
Cocaine	4.0	4.5	1.3	1.0	0.3	0.3	0.2	0.1
Crack	1.0	0.7	0.3	0.2	0.0 ^a	0.1	0.1	0.1
Heroin	0.4	0.2	0.2	0.2	0.1	0.1	0.1	0.1
Hallucinogens	2.8	2.7	1.0	1.1	0.1	0.3	0.1	0.1
LSD	0.5	0.5	0.2	0.2	0.0	0.0	0.0	0.0
PCP	0.1	0.0	0.0	0.0	0.0	*	*	0.0
Ecstasy	1.7	1.8	0.5	0.5	0.1	0.1	0.0	0.0
Inhalants	0.5	0.6	0.5	0.3	0.1	0.1	0.1	0.1
Nonmedical Use of Psychotherapeutics ^{3,4}	8.7	8.6	4.7	4.4	1.9	1.6	1.5	1.5
Pain Relievers	5.9	5.8	3.5	3.1	1.2	1.1	1.2	1.1
OxyContin [®]	0.7	0.7	0.2	0.3	0.1	0.1	0.1	0.1
Tranquilizers	2.9	2.7	1.6	1.3	0.6 ^a	0.4	0.4	0.3
Stimulants ⁴	2.1	2.5	0.9	0.9	0.3	0.3	0.2	0.2
Methamphetamine ⁴	0.5	1.0	0.4	0.3	0.1	0.1	0.1	0.1
Sedatives	0.2	0.2	0.3 ^a	0.1	0.1	0.1	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	12.9	12.9	6.4	5.9	2.3	2.2	1.8	1.8

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.12A Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2012 and 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLICIT DRUGS²	221	182	621	589	406	345	1,128	1,081
Marijuana and Hashish	196	169	557	545	346	311	701	736
Cocaine	6	18	15	7	3	2	7	16
Crack	0	0	1	0	*	1	*	*
Heroin	*	*	*	5	*	2	*	6
Hallucinogens	46	29	53	60	11	13	50	52
LSD	7	15	9 ^a	24	5	1	6	12
PCP	*	1	1	4	2	1	2	2
Ecstasy	22	11	27	21	9	1	16	14
Inhalants	26	10	34	20	27 ^b	6	101	85
Nonmedical Use of Psychotherapeutics ^{3,4}	69	46	163 ^a	110	76	63	384	329
Pain Relievers	44	37	132 ^a	83	63	44	319	262
OxyContin [®]	5	5	30	14	4	3	9	7
Tranquilizers	27 ^a	9	42	35	11	13	60	33
Stimulants ⁴	21	11	44 ^a	19	12	12	40	42
Methamphetamine ⁴	6	5	12	5	3	1	9	11
Sedatives	3	3	8	4	0	2	15	17
ILLICIT DRUGS OTHER THAN MARIJUANA²	104	75	208 ^a	158	104	77	501	448

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.12B Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2012 and 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLICIT DRUGS²	67.9	62.3	42.1	46.6	28.6	26.1	5.2	4.9
Marijuana and Hashish	60.2	57.9	37.8 ^a	43.2	24.4	23.6	3.2	3.3
Cocaine	1.7 ^a	6.3	1.0	0.5	0.2	0.2	0.0	0.1
Crack	0.0	0.0	0.0	0.0	*	0.1	*	*
Heroin	*	*	*	0.4	*	0.1	*	0.0
Hallucinogens	14.2	10.0	3.6	4.8	0.8	1.0	0.2	0.2
LSD	2.1	5.0	0.6 ^a	1.9	0.3	0.1	0.0	0.1
PCP	*	0.2	0.0	0.3	0.2	0.0	0.0	0.0
Ecstasy	6.6	3.9	1.8	1.7	0.6	0.1	0.1	0.1
Inhalants	7.9	3.3	2.3	1.6	1.9 ^b	0.5	0.5	0.4
Nonmedical Use of Psychotherapeutics ^{3,4}	21.3	15.9	11.1	8.7	5.4	4.8	1.8	1.5
Pain Relievers	13.5	12.5	9.0	6.5	4.4	3.3	1.5	1.2
OxyContin [®]	1.7	1.8	2.0	1.1	0.3	0.2	0.0	0.0
Tranquilizers	8.4 ^a	3.1	2.8	2.8	0.8	1.0	0.3	0.1
Stimulants ⁴	6.4	3.7	3.0 ^a	1.5	0.8	0.9	0.2	0.2
Methamphetamine ⁴	1.8	1.7	0.8	0.4	0.2	0.1	0.0	0.1
Sedatives	0.8	0.9	0.5	0.3	0.0	0.1	0.1	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA²	31.9	25.5	14.1	12.5	7.3	5.8	2.3	2.0

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.13A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2012 and 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLCIT DRUGS²	2,061	1,917	2,744 ^a	3,093	1,306	1,283	1,270	1,196
Marijuana and Hashish	1,869	1,712	2,468 ^b	2,835	1,169	1,140	980	949
Cocaine	205	228	140	130	22	20	31	20
Crack	18	14	7	11	2	2	10	11
Heroin	22	14	36	16	15	22	65	40
Hallucinogens	272	249	229	250	46	47	52	71
LSD	39	40	60	47	9	9	10	9
PCP	6	2	4	*	2	*	*	2
Ecstasy	190	151	127	137	26	13	18	19
Inhalants	41	32	45	35	11	9	25	30
Nonmedical Use of Psychotherapeutics ^{3,4}	597	511	630	581	253	257	352	313
Pain Relievers	391	336	440	371	184	192	295	234
OxyContin [®]	48	49	37	38	26	14	34	6
Tranquilizers	167	146	194	166	95 ^a	51	81	71
Stimulants ⁴	186	212	140	139	36	59	37	52
Methamphetamine ⁴	36	57	31	33	17	18	13	13
Sedatives	19	11	21	13	6	3	6	8
ILLCIT DRUGS OTHER THAN MARIJUANA²	832	769	872	846	303	322	428	401

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.13B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2012 and 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLCIT DRUGS²	47.0	48.7	29.6 ^b	33.4	18.2	17.0	9.2	8.5
Marijuana and Hashish	42.6	43.5	26.6 ^b	30.7	16.3	15.1	7.1	6.8
Cocaine	4.7	5.8	1.5	1.4	0.3	0.3	0.2	0.1
Crack	0.4	0.4	0.1	0.1	0.0	0.0	0.1	0.1
Heroin	0.5	0.3	0.4	0.2	0.2	0.3	0.5	0.3
Hallucinogens	6.2	6.3	2.5	2.7	0.6	0.6	0.4	0.5
LSD	0.9	1.0	0.7	0.5	0.1	0.1	0.1	0.1
PCP	0.1	0.1	0.0	*	0.0	*	*	0.0
Ecstasy	4.3	3.8	1.4	1.5	0.4	0.2	0.1	0.1
Inhalants	0.9	0.8	0.5	0.4	0.2	0.1	0.2	0.2
Nonmedical Use of Psychotherapeutics ^{3,4}	13.6	13.0	6.8	6.3	3.5	3.4	2.6	2.2
Pain Relievers	8.9	8.5	4.7	4.0	2.6	2.5	2.1	1.7
OxyContin [®]	1.1	1.2	0.4	0.4	0.4	0.2	0.2	0.0
Tranquilizers	3.8	3.7	2.1	1.8	1.3 ^b	0.7	0.6	0.5
Stimulants ⁴	4.2	5.4	1.5	1.5	0.5	0.8	0.3	0.4
Methamphetamine ⁴	0.8	1.4	0.3	0.4	0.2	0.2	0.1	0.1
Sedatives	0.4	0.3	0.2	0.1	0.1	0.0	0.0	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	19.0	19.5	9.4	9.1	4.2	4.3	3.1	2.9

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.14A Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2012 and 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLICIT DRUGS²	2,999	3,467	4,534	4,361	3,695	3,975	2,877	3,083
Marijuana and Hashish	2,411	2,841	3,631	3,525	2,738	3,120	1,791	1,926
Cocaine	468	491	407	277	174	231	174	109
Crack	150	98	138	75	12 ^a	102	104	64
Heroin	42	19	39	48	35	55	80	63
Hallucinogens	151	172	147	155	45	188	32	47
LSD	33	35	6	30	*	*	*	*
PCP	11	*	4	11	*	*	*	11
Ecstasy	76	128	74	80	33	75	11	10
Inhalants	24	65	116	56	30	83	13	65
Nonmedical Use of Psychotherapeutics ^{3,4}	807	870	1,207	1,230	1,126	906	1,165	1,268
Pain Relievers	571	595	926	898	676	588	820	883
OxyContin [®]	72	59	25	87	10	43	59	167
Tranquilizers	296	287	427	373	380	214	358	306
Stimulants ⁴	148	199	197	242	196	185	114	193
Methamphetamine ⁴	46	108	118	101	67	87	81	157
Sedatives	13	25	90	19	48	69	41	78
ILLICIT DRUGS OTHER THAN MARIJUANA²	1,260	1,297	1,657	1,582	1,310	1,287	1,304	1,404

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.14B Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2012 and 2013

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLICIT DRUGS²	24.4	28.2	14.2	13.2	5.5	5.9	3.2	3.4
Marijuana and Hashish	19.6	23.1	11.4	10.7	4.1	4.6	2.0	2.2
Cocaine	3.8	4.0	1.3	0.8	0.3	0.3	0.2	0.1
Crack	1.2	0.8	0.4	0.2	0.0 ^a	0.2	0.1	0.1
Heroin	0.3	0.2	0.1	0.1	0.1	0.1	0.1	0.1
Hallucinogens	1.2	1.4	0.5	0.5	0.1	0.3	0.0	0.1
LSD	0.3	0.3	0.0	0.1	*	*	*	*
PCP	0.1	*	0.0	0.0	*	*	*	0.0
Ecstasy	0.6	1.0	0.2	0.2	0.0	0.1	0.0	0.0
Inhalants	0.2	0.5	0.4	0.2	0.0	0.1	0.0	0.1
Nonmedical Use of Psychotherapeutics ^{3,4}	6.6	7.1	3.8	3.7	1.7	1.3	1.3	1.4
Pain Relievers	4.6	4.8	2.9	2.7	1.0	0.9	0.9	1.0
OxyContin [®]	0.6	0.5	0.1	0.3	0.0	0.1	0.1	0.2
Tranquilizers	2.4	2.3	1.3	1.1	0.6	0.3	0.4	0.3
Stimulants ⁴	1.2	1.6	0.6	0.7	0.3	0.3	0.1	0.2
Methamphetamine ⁴	0.4	0.9	0.4	0.3	0.1	0.1	0.1	0.2
Sedatives	0.1	0.2	0.3	0.1	0.1	0.1	0.0	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA²	10.2	10.5	5.2	4.8	1.9	1.9	1.5	1.6

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.15A Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Numbers in Thousands, 2012 and 2013

Drug	Total (2012)	Total (2013)	Aged 12-20 (2012)	Aged 12-20 (2013)	Aged 21+ (2012)	Aged 21+ (2013)
ILLCIT DRUGS¹	18,588	19,213	3,683	3,417	14,904	15,796
Marijuana and Hashish	15,384	16,200	3,338	3,168	12,046	13,032
Cocaine (Including Crack)	1,438	1,404	154	150	1,285	1,254
Heroin	190	180	30	14	160	165
Hallucinogens	999	1,163	388	332	611	831
Inhalants	353	316	138 ^b	62	216	254
Pain Relievers ²	3,428	3,143	611 ^b	433	2,817	2,710
Tranquilizers ²	1,640 ^a	1,295	234	175	1,406	1,120
Stimulants ^{2,3}	980	1,078	203	174	777	904
Methamphetamine ³	336	414	48	42	288	372
Sedatives ²	207	149	28	21	180	127
ILLCIT DRUGS OTHER THAN MARIJUANA¹	6,650	6,411	1,193 ^b	966	5,456	5,446

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.15B Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Percentages, 2012 and 2013

Drug	Total (2012)	Total (2013)	Aged 12-20 (2012)	Aged 12-20 (2013)	Aged 21+ (2012)	Aged 21+ (2013)
ILLCIT DRUGS¹	13.7	14.0	39.6	39.5	11.8	12.3
Marijuana and Hashish	11.4	11.8	35.9	36.6	9.5	10.2
Cocaine (Including Crack)	1.1	1.0	1.7	1.7	1.0	1.0
Heroin	0.1	0.1	0.3	0.2	0.1	0.1
Hallucinogens	0.7	0.8	4.2	3.8	0.5	0.6
Inhalants	0.3	0.2	1.5 ^b	0.7	0.2	0.2
Pain Relievers ²	2.5	2.3	6.6 ^b	5.0	2.2	2.1
Tranquilizers ²	1.2 ^a	0.9	2.5	2.0	1.1	0.9
Stimulants ^{2,3}	0.7	0.8	2.2	2.0	0.6	0.7
Methamphetamine ³	0.2	0.3	0.5	0.5	0.2	0.3
Sedatives ²	0.2	0.1	0.3	0.2	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	4.9	4.7	12.8 ^a	11.2	4.3	4.2

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.16A Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Numbers in Thousands, 2012 and 2013

Drug	Total (2012)	Total (2013)	Aged 12-20 (2012)	Aged 12-20 (2013)	Aged 21+ (2012)	Aged 21+ (2013)
ILLICIT DRUGS¹	9,188	8,939	1,881	1,685	7,307	7,253
Marijuana and Hashish	8,367	8,082	1,803	1,643	6,564	6,440
Cocaine (Including Crack)	502	477	35	22	468	455
Heroin	56	85	6	1	51	84
Hallucinogens	111	149	33	14	78	135
Inhalants	13	14	11	3	*	11
Pain Relievers ²	573	547	59 ^a	27	515	519
Tranquilizers ²	142	104	17	14	125	90
Stimulants ²	100	60	9	21	91	40
Methamphetamine ³	184	131	13	11	171	121
Sedatives ²	14	26	1	4	13	22
ILLICIT DRUGS OTHER THAN MARIJUANA¹	1,398	1,346	161 ^a	97	1,237	1,249

*Low precision; no estimate reported.

NOTE: Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.16B Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Drinking Age Groups: Percentages, 2012 and 2013

Drug	Total (2012)	Total (2013)	Aged 12-20 (2012)	Aged 12-20 (2013)	Aged 21+ (2012)	Aged 21+ (2013)
ILLCIT DRUGS¹	6.8	6.6	20.6	19.9	5.8	5.7
Marijuana and Hashish	6.2	5.9	19.8	19.5	5.2	5.0
Cocaine (Including Crack)	0.4	0.3	0.4	0.3	0.4	0.4
Heroin	0.0	0.1	0.1	0.0	0.0	0.1
Hallucinogens	0.1	0.1	0.4	0.2	0.1	0.1
Inhalants	0.0	0.0	0.1	0.0	*	0.0
Pain Relievers ²	0.4	0.4	0.6 ^a	0.3	0.4	0.4
Tranquilizers ²	0.1	0.1	0.2	0.2	0.1	0.1
Stimulants ²	0.1	0.0	0.1	0.2	0.1	0.0
Methamphetamine ³	0.1	0.1	0.1	0.1	0.1	0.1
Sedatives ²	0.0	0.0	0.0	0.0	0.0	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1.0	1.0	1.7 ^a	1.1	1.0	1.0

*Low precision; no estimate reported.

NOTE: Estimates are based on nonimputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.17A Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2012 and 2013

Drug	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
ILLICIT DRUGS¹	18,588	19,213	1,248 ^a	1,116	6,111	6,294	11,228	11,803
Marijuana and Hashish	15,384	16,200	1,099	1,026	5,505	5,687	8,780	9,486
Cocaine (Including Crack)	1,438	1,404	23	28	367	378	1,049	998
Heroin	190	180	*	6	73	52	117	122
Hallucinogens	999	1,163	110	102	547	546	342	515
Inhalants	353	316	87 ^b	36	98	76	169	204
Pain Relievers ²	3,428	3,143	239 ^b	163	1,016	899	2,173	2,081
Tranquilizers ²	1,640 ^a	1,295	80	57	456	363	1,104	874
Stimulants ^{2,3}	980	1,078	77 ^a	42	362	410	540	626
Methamphetamine ³	336	414	21	11	84	108	231	296
Sedatives ²	207	149	11	8	45	27	151	113
ILLICIT DRUGS OTHER THAN MARIJUANA¹	6,650	6,411	417 ^b	309	2,007	1,937	4,226	4,166

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.17B Types of Illicit Drug Use in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Percentages, 2012 and 2013

Drug	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
ILLCIT DRUGS¹	13.7	14.0	38.8	38.8	29.3	30.4	10.1	10.4
Marijuana and Hashish	11.4	11.8	34.2	35.7	26.4	27.4	7.9	8.4
Cocaine (Including Crack)	1.1	1.0	0.7	1.0	1.8	1.8	0.9	0.9
Heroin	0.1	0.1	*	0.2	0.4	0.2	0.1	0.1
Hallucinogens	0.7	0.8	3.4	3.6	2.6	2.6	0.3	0.5
Inhalants	0.3	0.2	2.7 ^b	1.2	0.5	0.4	0.2	0.2
Pain Relievers ²	2.5	2.3	7.4 ^a	5.7	4.9	4.3	2.0	1.8
Tranquilizers ²	1.2 ^a	0.9	2.5	2.0	2.2	1.8	1.0	0.8
Stimulants ^{2,3}	0.7	0.8	2.4 ^a	1.5	1.7	2.0	0.5	0.6
Methamphetamine ³	0.2	0.3	0.7	0.4	0.4	0.5	0.2	0.3
Sedatives ²	0.2	0.1	0.3	0.3	0.2	0.1	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA¹	4.9	4.7	13.0 ^a	10.7	9.6	9.3	3.8	3.7

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.18A Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2012 and 2013

Drug	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
ILLCIT DRUGS¹	9,188	8,939	576	499	3,200	3,209	5,411	5,231
Marijuana and Hashish	8,367	8,082	546	486	3,048	3,026	4,773	4,570
Cocaine (Including Crack)	502	477	5	3	83	109	414	365
Heroin	56	85	*	*	11	25	46	60
Hallucinogens	111	149	9	4	56	40	46	105
Inhalants	13	14	5	*	8	5	*	8
Pain Relievers ²	573	547	23	14	164	110	387	423
Tranquilizers ²	142	104	6	7	42	29	94	67
Stimulants ²	100	60	6	4	24	26	71	29
Methamphetamine ³	184	131	3	2	31	24	150	105
Sedatives ²	14	26	1	*	2	8	11	18
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1,398	1,346	55	32	355	327	988	987

*Low precision; no estimate reported.

NOTE: Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.18B Types of Illicit Drugs Used with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month among Past Month Alcohol Users Aged 12 or Older, by Age Group: Percentages, 2012 and 2013

Drug	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
ILLCIT DRUGS¹	6.8	6.6	18.5	18.0	15.5	15.6	4.9	4.6
Marijuana and Hashish	6.2	5.9	17.5	17.6	14.8	14.8	4.3	4.0
Cocaine (Including Crack)	0.4	0.3	0.2	0.1	0.4	0.5	0.4	0.3
Heroin	0.0	0.1	*	*	0.1	0.1	0.0	0.1
Hallucinogens	0.1	0.1	0.3	0.1	0.3	0.2	0.0	0.1
Inhalants	0.0	0.0	0.2	*	0.0	0.0	*	0.0
Pain Relievers ²	0.4	0.4	0.7	0.5	0.8	0.5	0.3	0.4
Tranquilizers ²	0.1	0.1	0.2	0.3	0.2	0.1	0.1	0.1
Stimulants ²	0.1	0.0	0.2	0.1	0.1	0.1	0.1	0.0
Methamphetamine ³	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Sedatives ²	0.0	0.0	0.0	*	0.0	0.0	0.0	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1.0	1.0	1.7	1.1	1.7	1.6	0.9	0.9

*Low precision; no estimate reported.

NOTE: Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. These estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Refers to nonmedical use; does not include over-the-counter drugs.

³ Estimates in the designated row include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.