

**Table 6.24A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2012 and 2013**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	11,973	11,101
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	3,483	3,383	5,553	5,430
Cigars	7,261	6,846	6,156	5,592
Pipe Tobacco	1,283	1,360	1,247 <sup>a</sup>	910
<b>ALCOHOL</b>	37,643	36,345	97,859	100,523
Binge Alcohol Use <sup>2</sup>	25,082	23,912	34,604	36,152
Heavy Alcohol Use <sup>2</sup>	9,084	8,782	7,937	7,759

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 6.24B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Percentages, 2012 and 2013**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	5.9 <sup>a</sup>	5.4
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	6.1	6.1	2.7	2.6
Cigars	12.6	12.3	3.0	2.7
Pipe Tobacco	2.2	2.4	0.6 <sup>a</sup>	0.4
<b>ALCOHOL</b>	65.4	65.2	48.3	48.7
Binge Alcohol Use <sup>2</sup>	43.6	42.9	17.1	17.5
Heavy Alcohol Use <sup>2</sup>	15.8	15.7	3.9	3.8

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 6.25A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Numbers in Thousands, 2012 and 2013**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	518	554
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	259	211	257	276
Cigars	393 <sup>a</sup>	311	252	265
Pipe Tobacco	104	90	61	70
<b>ALCOHOL</b>	1,003 <sup>b</sup>	782	2,214	2,095
Binge Alcohol Use <sup>2</sup>	742 <sup>b</sup>	588	1,057	968
Heavy Alcohol Use <sup>2</sup>	199	167	127	125

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 6.25B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Percentages, 2012 and 2013**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	2.2	2.4
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	15.8	15.1	1.1	1.2
Cigars	24.1	22.3	1.1	1.1
Pipe Tobacco	6.4	6.4	0.3	0.3
<b>ALCOHOL</b>	61.4 <sup>a</sup>	55.9	9.5	8.9
Binge Alcohol Use <sup>2</sup>	45.4	42.1	4.5	4.1
Heavy Alcohol Use <sup>2</sup>	12.2	12.0	0.5	0.5

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 6.26A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Numbers in Thousands, 2012 and 2013**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	2,159	2,226
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1,137	1,209	774	810
Cigars	2,321	2,138	1,395	1,335
Pipe Tobacco	386	419	239 <sup>a</sup>	339
<b>ALCOHOL</b>	8,417	8,270	12,411	12,469
Binge Alcohol Use <sup>2</sup>	6,658	6,399	7,004	6,788
Heavy Alcohol Use <sup>2</sup>	2,641	2,397	1,746	1,542

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 6.26B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Percentages, 2012 and 2013**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	9.2	9.2
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	10.3	11.3	3.3	3.4
Cigars	21.1	20.1	5.9	5.5
Pipe Tobacco	3.5	3.9	1.0 <sup>a</sup>	1.4
<b>ALCOHOL</b>	76.5	77.6	52.6	51.7
Binge Alcohol Use <sup>2</sup>	60.5	60.0	29.7 <sup>a</sup>	28.1
Heavy Alcohol Use <sup>2</sup>	24.0	22.5	7.4 <sup>a</sup>	6.4

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 6.27A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2012 and 2013**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	9,296 <sup>a</sup>	8,322
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	2,087	1,963	4,522	4,344
Cigars	4,546	4,397	4,509	3,992
Pipe Tobacco	794	851	947 <sup>b</sup>	502
<b>ALCOHOL</b>	28,224	27,293	83,234	85,959
Binge Alcohol Use <sup>2</sup>	17,682	16,925	26,542	28,396
Heavy Alcohol Use <sup>2</sup>	6,245	6,218	6,064	6,092

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 6.27B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Percentages, 2012 and 2013**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	6.0 <sup>a</sup>	5.2
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	4.7	4.5	2.9	2.7
Cigars	10.1	10.1	2.9	2.5
Pipe Tobacco	1.8	1.9	0.6 <sup>b</sup>	0.3
<b>ALCOHOL</b>	62.9	62.4	53.5	54.1
Binge Alcohol Use <sup>2</sup>	39.4	38.7	17.1	17.9
Heavy Alcohol Use <sup>2</sup>	13.9	14.2	3.9	3.8

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.