

Table 6.28A Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2012 and 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	10,959	10,385	19,743	18,669	15,765	15,601	23,031	22,224
Cigarettes	9,084	8,782	15,998	15,130	12,561	12,432	19,882	19,433
Smokeless Tobacco	2,121	2,005	2,806	2,729	1,585	1,540	2,524	2,540
Cigars	2,946	2,552	4,542	4,669	3,201	2,973	2,727 ^a	2,244
Pipe Tobacco	330	389	642	728	794 ^a	528	765	626

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.28B Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2012 and 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	64.4	62.8	46.3 ^b	42.9	20.8	20.3	18.5	17.7
Cigarettes	53.4	53.1	37.5 ^b	34.8	16.6	16.2	16.0	15.5
Smokeless Tobacco	12.5	12.1	6.6	6.3	2.1	2.0	2.0	2.0
Cigars	17.3	15.4	10.6	10.7	4.2	3.9	2.2 ^a	1.8
Pipe Tobacco	1.9	2.4	1.5	1.7	1.0 ^a	0.7	0.6	0.5

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.29A Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2012 and 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	239	209	683 ^a	580	346	284	885	879
Cigarettes	199	167	543 ^b	421	260 ^a	194	632	615
Smokeless Tobacco	63	78	162	142	48	61	242	206
Cigars	89	95	245	200	110 ^a	74	201	207
Pipe Tobacco	17	19	52	57	44 ^a	18	53	66

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.29B Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2012 and 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	73.2	71.3	46.4	45.9	24.4	21.5	4.1	4.0
Cigarettes	61.0	57.1	36.9	33.3	18.4	14.7	2.9	2.8
Smokeless Tobacco	19.5	26.8	11.0	11.2	3.4	4.6	1.1	0.9
Cigars	27.2	32.6	16.7	15.8	7.8	5.6	0.9	0.9
Pipe Tobacco	5.2	6.4	3.5	4.5	3.1 ^a	1.4	0.2	0.3

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.30A Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2012 and 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	3,121 ^a	2,805	4,807	4,857	2,214	2,356	3,024	2,864
Cigarettes	2,641	2,397	4,017	4,002	1,759	1,871	2,591	2,387
Smokeless Tobacco	722	677	647	733	199	249	344	359
Cigars	1,088 ^a	929	1,356	1,370	622	602	650	572
Pipe Tobacco	156	151	215 ^a	306	129	160	124	141

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.30B Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2012 and 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	71.2	71.2	51.8	52.5	30.9	31.2	22.0	20.4
Cigarettes	60.2	60.9	43.3	43.3	24.5	24.8	18.8 ^a	17.0
Smokeless Tobacco	16.5	17.2	7.0	7.9	2.8	3.3	2.5	2.6
Cigars	24.8	23.6	14.6	14.8	8.7	8.0	4.7	4.1
Pipe Tobacco	3.6	3.8	2.3 ^a	3.3	1.8	2.1	0.9	1.0

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.31A Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2012 and 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	7,599	7,372	14,253	13,232	13,204	12,961	19,122	18,481
Cigarettes	6,245	6,218	11,437	10,708	10,542	10,368	16,659	16,431
Smokeless Tobacco	1,336	1,250	1,998	1,854	1,337	1,230	1,938	1,974
Cigars	1,769	1,528	2,941	3,099	2,469	2,297	1,877	1,465
Pipe Tobacco	157	220	374	365	621 ^a	349	588	419

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.31B Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2012 and 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	61.7	59.9	44.7 ^b	40.1	19.6	19.1	21.5	20.7
Cigarettes	50.7	50.5	35.8 ^a	32.4	15.7	15.3	18.7	18.4
Smokeless Tobacco	10.9	10.2	6.3	5.6	2.0	1.8	2.2	2.2
Cigars	14.4	12.4	9.2	9.4	3.7	3.4	2.1	1.6
Pipe Tobacco	1.3	1.8	1.2	1.1	0.9 ^a	0.5	0.7	0.5

*Low precision; no estimate reported.

^a Difference between estimate and 2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2013 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.