

**Table 6.47A Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	1,947 <sup>a</sup>	2,287	235	216	495	476	1,217 <sup>b</sup>	1,594
From More Than One Doctor	204	276	29	32	46	46	128	198
Wrote Fake Prescription	24	13	6	3	11	3	7	6
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	50	78	13	7	13	7	24	65
From Friend or Relative for Free	5,822	5,713	559 <sup>b</sup>	428	1,796 <sup>b</sup>	1,597	3,466	3,688
Bought from Friend or Relative	1,306	1,149	113	92	513 <sup>a</sup>	428	679	629
Took from Friend or Relative without Asking	473	429	123 <sup>a</sup>	92	148	157	202	180
Bought from Drug Dealer or Other Stranger	423	460	46	37	222	225	156	198
Bought on the Internet	27	10	0	2	3	7	23	1
Some Other Way	463	373	88	68	128	112	247	193

\*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.47B Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	18.1 <sup>a</sup>	21.2	19.3	22.1	14.7	15.6	19.8 <sup>a</sup>	23.6
From More Than One Doctor	1.9	2.6	2.4	3.3	1.4	1.5	2.1	2.9
Wrote Fake Prescription	0.2	0.1	0.5	0.3	0.3	0.1	0.1	0.1
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.5	0.7	1.1	0.7	0.4	0.2	0.4	1.0
From Friend or Relative for Free	54.2	53.0	46.1	43.8	53.2	52.2	56.4	54.6
Bought from Friend or Relative	12.2	10.6	9.4	9.4	15.2	14.0	11.0	9.3
Took from Friend or Relative without Asking	4.4	4.0	10.2	9.4	4.4	5.1	3.3	2.7
Bought from Drug Dealer or Other Stranger	3.9	4.3	3.8	3.8	6.6	7.3	2.5	2.9
Bought on the Internet	0.3	0.1	0.0	0.2	0.1	0.2	0.4	0.0
Some Other Way	4.3	3.5	7.2	7.0	3.8	3.7	4.0	2.9

\*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.48A Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Pain Relievers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	4,243	4,359	282 <sup>b</sup>	218	1,165	1,101	2,795	3,040
From More Than One Doctor	160	173	31	24	70	54	58	94
Wrote Fake Prescription	17	1	1	0	3	1	14	0
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	18	13	5	2	11	9	2	2
From Friend or Relative for Free	286	264	41	34	123	103	122	127
Bought from Friend or Relative	202	193	29	23	117 <sup>b</sup>	73	56	97
Took from Friend or Relative without Asking	92	60	43 <sup>b</sup>	16	38	34	11	10
Bought from Drug Dealer or Other Stranger	96	72	17	12	51	34	28	26
Bought on the Internet	9	15	0	2	1	5	8	7
Some Other Way	77	49	23	15	18	16	36	18

\*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.48B Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Pain Relievers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	81.6	83.8	59.5	63.1	73.0 <sup>a</sup>	77.0	89.3	88.8
From More Than One Doctor	3.1	3.3	6.6	7.1	4.4	3.8	1.9	2.8
Wrote Fake Prescription	0.3	0.0	0.2	0.1	0.2	0.0	0.5	0.0
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3	0.2	1.1	0.4	0.7	0.7	0.1	0.1
From Friend or Relative for Free	5.5	5.1	8.7	9.9	7.7	7.2	3.9	3.7
Bought from Friend or Relative	3.9	3.7	6.0	6.6	7.3 <sup>a</sup>	5.1	1.8	2.8
Took from Friend or Relative without Asking	1.8	1.2	9.1 <sup>b</sup>	4.6	2.4	2.4	0.3	0.3
Bought from Drug Dealer or Other Stranger	1.9	1.4	3.7	3.4	3.2	2.4	0.9	0.8
Bought on the Internet	0.2	0.3	0.0	0.5	0.0	0.4	0.2	0.2
Some Other Way	1.5	0.9	4.9	4.3	1.1	1.1	1.2	0.5

\*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.49A Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	1,146	1,166	107	107	284	237	755	823
From More Than One Doctor	194	232	24	25	46	47	123	160
Wrote Fake Prescription	27	20	4	5	4	2	19	12
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	49	56	6	6	15 <sup>a</sup>	4	29	46
From Friend or Relative for Free	2,565	2,465	234 <sup>b</sup>	173	812 <sup>a</sup>	688	1,519	1,604
Bought from Friend or Relative	1,127	1,045	79 <sup>b</sup>	45	421 <sup>a</sup>	335	628	666
Took from Friend or Relative without Asking	498 <sup>a</sup>	364	71 <sup>a</sup>	45	179 <sup>b</sup>	110	247	208
Bought from Drug Dealer or Other Stranger	514	504	47 <sup>a</sup>	28	264	226	204	250
Bought on the Internet	40	23	1	2	15	11	24	11
Some Other Way	254	182	37	34	77	71	140	77

\*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.49B Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	26.7	27.6	23.3	28.3	22.4	21.3	29.4	30.0
From More Than One Doctor	4.5	5.5	5.3	6.6	3.6	4.2	4.8	5.8
Wrote Fake Prescription	0.6	0.5	0.8	1.3	0.3	0.2	0.7	0.5
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.1	1.3	1.2	1.5	1.2 <sup>a</sup>	0.4	1.1	1.7
From Friend or Relative for Free	59.7	58.2	50.8	45.9	64.0	61.8	59.2	58.5
Bought from Friend or Relative	26.3	24.7	17.1 <sup>a</sup>	11.9	33.2	30.1	24.5	24.3
Took from Friend or Relative without Asking	11.6 <sup>a</sup>	8.6	15.5	12.1	14.2 <sup>a</sup>	9.9	9.6	7.6
Bought from Drug Dealer or Other Stranger	12.0	11.9	10.2	7.5	20.8	20.3	8.0	9.1
Bought on the Internet	0.9	0.5	0.2	0.5	1.2	1.0	0.9	0.4
Some Other Way	6.1	4.4	8.3	9.4	6.2	6.6	5.6	2.9

\*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.50A Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Pain Relievers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	1,814	1,863	111	97	534	488	1,169	1,278
From More Than One Doctor	166	159	25	18	70	67	72	75
Wrote Fake Prescription	37	17	0	2	4	3	33	12
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	26	23	2	3	13	11	11	*
From Friend or Relative for Free	328	267	29	18	129	115	170	134
Bought from Friend or Relative	294	271	27	17	142	108	125	145
Took from Friend or Relative without Asking	98	98	23 <sup>b</sup>	9	45	46	30	43
Bought from Drug Dealer or Other Stranger	203	150	20	10	100	70	83	70
Bought on the Internet	6	28	0	2	4	9	1	16
Some Other Way	79	49	16	8	25	20	39	21

\*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.50B Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Pain Relievers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	77.5 <sup>a</sup>	82.3	55.1 <sup>a</sup>	65.1	71.4 <sup>a</sup>	77.1	83.9	86.3
From More Than One Doctor	7.1	7.1	12.4	11.9	9.4	10.6	5.2	5.1
Wrote Fake Prescription	1.6	0.8	0.1	1.4	0.5	0.5	2.4	0.8
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.1	1.0	0.8	2.0	1.8	1.7	0.8	*
From Friend or Relative for Free	14.1	12.0	14.7	12.2	17.5	18.3	12.2	9.2
Bought from Friend or Relative	12.6	12.1	13.5	11.7	19.2	17.2	9.0	10.0
Took from Friend or Relative without Asking	4.2	4.4	11.5	6.1	6.1	7.3	2.2	3.0
Bought from Drug Dealer or Other Stranger	8.7	6.7	9.8	7.0	13.5	11.1	6.0	4.8
Bought on the Internet	0.2	1.2	0.1	1.5	0.6	1.5	0.1	1.1
Some Other Way	3.5	2.2	8.0	5.5	3.4	3.2	2.8	1.5

\*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.51A Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	661	690	31	24	116	110	514	557
From More Than One Doctor	51	43	5	3	15	22	31	18
Wrote Fake Prescription	4	2	3	1	1	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	14	17	1	0	9	2	4	15
From Friend or Relative for Free	2,901 <sup>a</sup>	3,245	192	161	830	779	1,879 <sup>b</sup>	2,305
Bought from Friend or Relative	669	640	61	57	293	297	314	286
Took from Friend or Relative without Asking	165	208	33	29	71	76	61	104
Bought from Drug Dealer or Other Stranger	296	262	39 <sup>a</sup>	21	159	158	98	84
Bought on the Internet	17	5	1	2	2	3	*	*
Some Other Way	120	175	18	16	42	67	60	91

\*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.51B Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	13.5	13.1	8.0	7.6	7.6	7.3	17.3	16.1
From More Than One Doctor	1.0	0.8	1.3	0.9	1.0	1.5	1.0	0.5
Wrote Fake Prescription	0.1	0.0	0.7	0.3	0.1	0.1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3	0.3	0.4	0.1	0.6	0.1	0.1	0.4
From Friend or Relative for Free	59.2	61.4	50.0	51.4	53.9	51.4	63.2	66.6
Bought from Friend or Relative	13.7	12.1	16.0	18.3	19.0	19.6	10.6	8.3
Took from Friend or Relative without Asking	3.4	3.9	8.6	9.1	4.6	5.0	2.0	3.0
Bought from Drug Dealer or Other Stranger	6.0	5.0	10.3	6.6	10.3	10.4	3.3	2.4
Bought on the Internet	0.4	0.1	0.2	0.6	0.2	0.2	*	*
Some Other Way	2.4	3.3	4.6	5.1	2.7	4.4	2.0	2.6

\*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.52A Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Tranquilizers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	2,141	2,417	82	73	524	485	1,534 <sup>a</sup>	1,859
From More Than One Doctor	69	75	8	6	35	22	26	47
Wrote Fake Prescription	2	0	*	0	2	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	6	6	*	1	1	2	5	3
From Friend or Relative for Free	221	186	22	21	85	73	115	92
Bought from Friend or Relative	93	123	14	15	41	59	38	49
Took from Friend or Relative without Asking	25	32	8	10	13	13	4	10
Bought from Drug Dealer or Other Stranger	62	65	12	5	32	33	18	27
Bought on the Internet	7	10	*	*	1	*	5	6
Some Other Way	21	31	8	*	4	13	10	17

\*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.52B Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Tranquilizers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	80.9	82.1	53.1	55.8	71.1	69.0	87.4	88.1
From More Than One Doctor	2.6	2.6	5.0	4.6	4.7	3.2	1.5	2.2
Wrote Fake Prescription	0.1	0.0	*	0.3	0.2	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2	0.2	*	0.6	0.1	0.4	0.3	0.1
From Friend or Relative for Free	8.4	6.3	14.0	15.9	11.5	10.4	6.5	4.4
Bought from Friend or Relative	3.5	4.2	9.2	11.2	5.6	8.3	2.2	2.3
Took from Friend or Relative without Asking	0.9	1.1	5.4	7.3	1.7	1.8	0.2	0.5
Bought from Drug Dealer or Other Stranger	2.3	2.2	7.8	3.8	4.4	4.6	1.0	1.3
Bought on the Internet	0.3	0.3	*	*	0.2	*	0.3	0.3
Some Other Way	0.8	1.0	4.9	*	0.5	1.8	0.6	0.8

\*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.53A Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	334	354	10	*	63	44	261	299
From More Than One Doctor	14	15	3	2	8	10	4	2
Wrote Fake Prescription	3	1	1	*	2	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	21	4	1	*	4	*	*	4
From Friend or Relative for Free	1,040	1,060	45	49	255	229	740	782
Bought from Friend or Relative	376	307	29	19	130	123	217	166
Took from Friend or Relative without Asking	91	92	9	10	43	32	39	51
Bought from Drug Dealer or Other Stranger	146	131	17	11	84 <sup>a</sup>	57	45	63
Bought on the Internet	12	6	0	*	4	*	8	*
Some Other Way	41	48	6	6	9	17	26	25

\*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.53B Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	18.6	19.6	9.1	*	12.5	9.8	22.1	23.8
From More Than One Doctor	0.8	0.8	3.0	2.6	1.5	2.3	0.3	0.2
Wrote Fake Prescription	0.2	0.0	1.2	*	0.4	0.0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.2	0.2	1.0	*	0.9	*	*	0.3
From Friend or Relative for Free	57.9	58.8	41.0	52.0	50.6	50.8	62.6	62.2
Bought from Friend or Relative	20.9	17.0	26.1	19.8	25.8	27.2	18.3	13.2
Took from Friend or Relative without Asking	5.1	5.1	7.7	10.3	8.5	7.0	3.3	4.0
Bought from Drug Dealer or Other Stranger	8.1	7.3	15.6	11.2	16.7	12.7	3.8	5.0
Bought on the Internet	0.6	0.4	0.4	*	0.7	*	0.6	*
Some Other Way	2.3	2.7	6.0	6.4	1.8	3.9	2.2	2.0

\*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.54A Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Tranquilizers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	756	789	*	*	163	140	572	631
From More Than One Doctor	55	42	*	*	20 <sup>a</sup>	6	34	33
Wrote Fake Prescription	1	*	*	*	1	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	7	*	*	*	3	*	*	*
From Friend or Relative for Free	132	75	*	*	36	28	87	39
Bought from Friend or Relative	61	74	*	*	25	40	30	23
Took from Friend or Relative without Asking	23	16	*	*	8	2	*	12
Bought from Drug Dealer or Other Stranger	47	44	*	*	19	17	26	23
Bought on the Internet	6	16	*	*	*	*	*	12
Some Other Way	12	20	*	*	3	*	8	*

\*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.54B Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Tranquilizers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	77.9	79.6	*	*	70.5	64.6	81.8	85.6
From More Than One Doctor	5.7	4.2	*	*	8.6 <sup>a</sup>	2.9	4.9	4.5
Wrote Fake Prescription	0.1	*	*	*	0.4	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.7	*	*	*	1.3	*	*	*
From Friend or Relative for Free	13.6 <sup>a</sup>	7.6	*	*	15.8	12.9	12.5	5.4
Bought from Friend or Relative	6.3	7.5	*	*	10.8	18.6	4.3	3.2
Took from Friend or Relative without Asking	2.4	1.7	*	*	3.4	0.9	*	1.7
Bought from Drug Dealer or Other Stranger	4.8	4.5	*	*	8.1	8.0	3.7	3.2
Bought on the Internet	0.7	1.6	*	*	*	*	*	1.6
Some Other Way	1.3	2.0	*	*	1.2	*	1.1	*

\*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.55A Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

Source for Most Recent Nonmedical Use among Past Year Stimulant Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	211	278	19	27	65	62	126	190
From More Than One Doctor	13	32	5	3	8	21	*	8
Wrote Fake Prescription	3	2	3	0	0	2	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	14	6	2	1	4	2	*	3
From Friend or Relative for Free	1,065 <sup>a</sup>	1,304	94	106	504	534	466 <sup>a</sup>	664
Bought from Friend or Relative	426 <sup>b</sup>	560	41	46	265	321	120	194
Took from Friend or Relative without Asking	86	95	18	17	30	25	38	53
Bought from Drug Dealer or Other Stranger	146	129	16	15	61	80	68	34
Bought on the Internet	21	27	1	*	6	12	14	15
Some Other Way	72	61	5	6	24	28	43	26

\*Low precision; no estimate reported.

NOTE: Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.55B Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source for Most Recent Nonmedical Use among Past Year Stimulant Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	10.3	11.2	9.4	12.1	6.8	5.7	14.3	16.0
From More Than One Doctor	0.6	1.3	2.3	1.5	0.8	1.9	*	0.6
Wrote Fake Prescription	0.2	0.1	1.6	0.0	0.0	0.2	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.7	0.2	1.0	0.3	0.4	0.2	*	0.3
From Friend or Relative for Free	51.8	52.3	46.1	48.2	52.1	49.1	52.8	55.9
Bought from Friend or Relative	20.7	22.5	20.0	20.8	27.4	29.5	13.5	16.3
Took from Friend or Relative without Asking	4.2	3.8	8.9	7.6	3.1	2.3	4.3	4.5
Bought from Drug Dealer or Other Stranger	7.1	5.2	8.0	6.7	6.3	7.4	7.7	2.9
Bought on the Internet	1.0	1.1	0.5	*	0.6	1.1	1.6	1.3
Some Other Way	3.5	2.4	2.4	2.8	2.5	2.6	4.9	2.2

\*Low precision; no estimate reported.

NOTE: Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.56A Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Stimulants	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	766 <sup>a</sup>	938	48	60	338	387	380	491
From More Than One Doctor	30	36	6	6	20	24	4	6
Wrote Fake Prescription	1	0	*	*	0	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	2	*	0	2	2	*	*
From Friend or Relative for Free	70	76	9	11	43	28	18	37
Bought from Friend or Relative	55	59	6	*	37	31	12	24
Took from Friend or Relative without Asking	12	5	*	2	6	3	*	*
Bought from Drug Dealer or Other Stranger	26	37	4	3	13	21	*	13
Bought on the Internet	4	3	1	*	*	1	4	*
Some Other Way	2	5	1	2	1	2	*	2

\*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.56B Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Stimulants	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	79.1	80.7	58.1	67.1	73.6	77.5	89.2	85.7
From More Than One Doctor	3.1	3.1	7.7	6.4	4.3	4.8	0.8	1.1
Wrote Fake Prescription	0.1	0.0	*	*	0.1	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2	0.2	*	0.1	0.4	0.4	*	*
From Friend or Relative for Free	7.2	6.5	11.3	11.8	9.3	5.6	4.2	6.5
Bought from Friend or Relative	5.7	5.1	7.5	*	8.1	6.2	2.8	4.1
Took from Friend or Relative without Asking	1.2	0.4	*	2.1	1.3	0.6	*	*
Bought from Drug Dealer or Other Stranger	2.7	3.2	5.1	3.6	2.8	4.2	*	2.3
Bought on the Internet	0.5	0.3	0.8	*	*	0.3	0.9	*
Some Other Way	0.2	0.4	1.3	1.7	0.2	0.3	*	0.4

\*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.57A Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

Source for Past Month Nonmedical Use among Past Month Stimulant Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	151	140	9	13	42	26	*	101
From More Than One Doctor	10	22	3	3	7	8	*	*
Wrote Fake Prescription	3	*	*	*	0	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	3	*	0	*	3	2	*	*
From Friend or Relative for Free	342 <sup>a</sup>	455	28	37	165	190	150	228
Bought from Friend or Relative	210	276	18	20	120	125	72	*
Took from Friend or Relative without Asking	54	46	8	7	25	16	20	24
Bought from Drug Dealer or Other Stranger	97	82	12	12	41	36	43	34
Bought on the Internet	10	23	1	*	4	6	5	*
Some Other Way	29	35	4	2	4 <sup>a</sup>	15	21	18

\*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.57B Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source for Past Month Nonmedical Use among Past Month Stimulant Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	20.9	16.6	12.7	19.5	13.5 <sup>a</sup>	7.4	*	23.4
From More Than One Doctor	1.4	2.6	3.6	4.1	2.4	2.2	*	*
Wrote Fake Prescription	0.4	*	*	*	0.0	0.3	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.4	*	0.6	*	0.9	0.5	*	*
From Friend or Relative for Free	47.2	53.7	39.7 <sup>a</sup>	54.4	53.1	54.3	43.4	53.0
Bought from Friend or Relative	29.0	32.7	26.0	30.1	38.9	36.0	20.8	*
Took from Friend or Relative without Asking	7.4	5.5	11.9	10.0	8.3	4.5	5.8	5.6
Bought from Drug Dealer or Other Stranger	13.4	9.7	17.8	18.1	13.4	10.4	12.4	7.8
Bought on the Internet	1.4	2.7	1.0	*	1.3	1.8	1.5	*
Some Other Way	4.2	4.2	6.1	3.4	1.3 <sup>a</sup>	4.4	6.3	4.1

\*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.58A Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Stimulants	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	257	322	*	*	119	154	*	*
From More Than One Doctor	17	18	*	*	12	5	*	*
Wrote Fake Prescription	0	*	*	*	*	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1	*	*	*	*	*	*	*
From Friend or Relative for Free	30	52	*	*	16	14	*	*
Bought from Friend or Relative	22	53	*	*	14	10	*	*
Took from Friend or Relative without Asking	10	*	*	*	7	1	*	*
Bought from Drug Dealer or Other Stranger	18	43	*	*	6	7	*	*
Bought on the Internet	6	*	*	*	*	*	*	*
Some Other Way	2	4	*	*	0	3	*	*

\*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.58B Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Stimulants	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	82.3	78.8	*	*	80.8	86.0	*	*
From More Than One Doctor	5.4	4.5	*	*	8.4	3.0	*	*
Wrote Fake Prescription	0.0	*	*	*	*	0.1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.5	*	*	*	*	*	*	*
From Friend or Relative for Free	9.9	12.9	*	*	10.9	7.9	*	*
Bought from Friend or Relative	7.4	13.0	*	*	9.6	5.4	*	*
Took from Friend or Relative without Asking	3.4	*	*	*	4.6	0.5	*	*
Bought from Drug Dealer or Other Stranger	6.0	10.5	*	*	4.4	4.0	*	*
Bought on the Internet	1.9	*	*	*	*	*	*	*
Some Other Way	0.7	1.0	*	*	0.1	1.7	*	*

\*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.59A Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

<b>Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users</b>	<b>Total (2010-2011)</b>	<b>Total (2012-2013)</b>	<b>Aged 12-17 (2010-2011)</b>	<b>Aged 12-17 (2012-2013)</b>	<b>Aged 18-25 (2010-2011)</b>	<b>Aged 18-25 (2012-2013)</b>	<b>Aged 26+ (2010-2011)</b>	<b>Aged 26+ (2012-2013)</b>
From Friend or Relative for Free	356	420	*	*	110	136	204	254
Bought from Friend or Relative	250	229	*	10	43	52	*	167
Took from Friend or Relative without Asking	26	29	2	3	*	5	13	*
Bought from Drug Dealer or Other Stranger	189	279	*	14	48 <sup>a</sup>	84	*	180
Bought on the Internet	9	2	*	*	3	0	*	1
Some Other Way	28	71	*	1	6	12	15	*

\*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.59B Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From Friend or Relative for Free	41.5	40.8	*	*	49.6	47.0	36.7	37.2
Bought from Friend or Relative	29.2	22.3	*	16.7	19.5	18.0	*	24.6
Took from Friend or Relative without Asking	3.0	2.9	3.0	4.9	*	1.7	2.3	*
Bought from Drug Dealer or Other Stranger	22.0	27.1	*	24.6	21.7	29.1	*	26.5
Bought on the Internet	1.0	0.1	*	*	1.5	0.2	*	0.1
Some Other Way	3.3	6.9	*	2.1	2.9	4.2	2.7	*

\*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.60A Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Methamphetamine	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From Friend or Relative for Free	108	116	*	*	35	42	*	*
Bought from Friend or Relative	84	118	*	*	25	37	*	*
Took from Friend or Relative without Asking	*	4	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	48	69	*	*	20	36	*	*
Bought on the Internet	2	*	*	*	*	*	*	*
Some Other Way	18	*	*	*	9	8	*	*

\*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.60B Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Methamphetamine	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From Friend or Relative for Free	40.1	34.3	*	*	37.9	33.4	*	*
Bought from Friend or Relative	31.4	34.9	*	*	26.5	29.7	*	*
Took from Friend or Relative without Asking	*	1.2	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	17.8	20.4	*	*	21.0	28.8	*	*
Bought on the Internet	0.7	*	*	*	*	*	*	*
Some Other Way	6.9	*	*	*	9.3	6.7	*	*

\*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.61A Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

<b>Source for Past Month Nonmedical Use among Past Month Methamphetamine Users</b>	<b>Total (2010-2011)</b>	<b>Total (2012-2013)</b>	<b>Aged 12-17 (2010-2011)</b>	<b>Aged 12-17 (2012-2013)</b>	<b>Aged 18-25 (2010-2011)</b>	<b>Aged 18-25 (2012-2013)</b>	<b>Aged 26+ (2010-2011)</b>	<b>Aged 26+ (2012-2013)</b>
From Friend or Relative for Free	166	196	*	*	*	*	*	*
Bought from Friend or Relative	*	150	*	*	*	*	*	*
Took from Friend or Relative without Asking	9	*	*	*	*	3	*	*
Bought from Drug Dealer or Other Stranger	101	112	*	*	*	*	*	*
Bought on the Internet	7	1	*	*	*	0	*	*
Some Other Way	9	*	*	*	*	4	*	*

\*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.61B Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

<b>Source for Past Month Nonmedical Use among Past Month Methamphetamine Users</b>	<b>Total (2010-2011)</b>	<b>Total (2012-2013)</b>	<b>Aged 12-17 (2010-2011)</b>	<b>Aged 12-17 (2012-2013)</b>	<b>Aged 18-25 (2010-2011)</b>	<b>Aged 18-25 (2012-2013)</b>	<b>Aged 26+ (2010-2011)</b>	<b>Aged 26+ (2012-2013)</b>
From Friend or Relative for Free	51.1	43.6	*	*	*	*	*	*
Bought from Friend or Relative	*	33.3	*	*	*	*	*	*
Took from Friend or Relative without Asking	2.9	*	*	*	*	2.8	*	*
Bought from Drug Dealer or Other Stranger	31.0	24.8	*	*	*	*	*	*
Bought on the Internet	2.0	0.1	*	*	*	0.1	*	*
Some Other Way	2.7	*	*	*	*	4.7	*	*

\*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.62A Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Methamphetamine	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	5	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.62B Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Methamphetamine	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	3.0	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.63A Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

Source for Most Recent Nonmedical Use among Past Year Sedative Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	132	102	13	*	21	17	*	69
From More Than One Doctor	*	12	*	*	2	*	*	5
Wrote Fake Prescription	1	*	*	*	*	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	*	4	*	*	3	0	*
From Friend or Relative for Free	385	290	36 <sup>b</sup>	14	77	69	*	*
Bought from Friend or Relative	53	64	9	6	15	13	29	46
Took from Friend or Relative without Asking	25	22	4	8	9	4	*	*
Bought from Drug Dealer or Other Stranger	24	13	3	2	5	6	16	5
Bought on the Internet	0	1	0	*	*	0	*	*
Some Other Way	17	23	*	*	3	*	*	20

\*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.63B Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source for Most Recent Nonmedical Use among Past Year Sedative Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	20.1	18.2	16.9	*	16.0	14.3	*	17.6
From More Than One Doctor	*	2.2	*	*	1.4	*	*	1.4
Wrote Fake Prescription	0.2	*	*	*	*	0.6	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.7	*	5.7	*	*	2.6	0.1	*
From Friend or Relative for Free	58.9	51.6	48.7 <sup>b</sup>	27.2	58.5	57.9	*	*
Bought from Friend or Relative	8.1	11.4	12.4	10.9	11.7	10.5	6.4	11.8
Took from Friend or Relative without Asking	3.8	3.9	5.8 <sup>a</sup>	16.1	6.7	3.4	*	*
Bought from Drug Dealer or Other Stranger	3.7	2.3	3.6	3.9	3.7	4.9	3.7	1.4
Bought on the Internet	0.0	0.2	0.1	*	*	0.1	*	*
Some Other Way	2.5	4.2	*	*	2.0	*	*	5.1

\*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.64A Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Sedatives	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	326	238	*	*	*	*	*	*
From More Than One Doctor	6	8	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	3	*	*	*	*	*	*
From Friend or Relative for Free	7	12	*	*	*	*	*	*
Bought from Friend or Relative	5	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	7	3	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	7	8	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	3	0	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.64B Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Sedatives	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	89.3	86.1	*	*	*	*	*	*
From More Than One Doctor	1.7	2.9	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.7	1.0	*	*	*	*	*	*
From Friend or Relative for Free	1.8	4.3	*	*	*	*	*	*
Bought from Friend or Relative	1.5	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	1.8	1.1	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	2.0	2.8	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	0.7	0.0	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.65A Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

Source for Past Month Nonmedical Use among Past Month Sedative Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	0	7	*	*	*	*	*	*
Wrote Fake Prescription	1	1	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	7	7	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	11	*	*	*	*	*	*	*
Bought on the Internet	0	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.65B Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source for Past Month Nonmedical Use among Past Month Sedative Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	0.0	3.0	*	*	*	*	*	*
Wrote Fake Prescription	0.4	0.3	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.8	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	2.7	3.0	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	4.1	*	*	*	*	*	*	*
Bought on the Internet	0.1	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.66A Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Sedatives	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.66B Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, Annual Averages Based on 2010-2011 and 2012-2013**

Source Where Friend or Relative Obtained Sedatives	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

\*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

<sup>a</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.