

Table 6.71A Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
ILLCIT DRUGS²	6,530 ^a	7,039	113	122	6,390 ^a	6,886
Marijuana and Hashish	4,983 ^b	5,498	94	110	4,869 ^b	5,368
Cocaine	372	345	2	5	368	338
Crack	66	67	2	*	63	65
Heroin	69	88	5	4	63	83
Hallucinogens	374	356	1	3	369	349
LSD	30	56	*	*	29	56
PCP	6	12	0	*	6	12
Ecstasy	247	223	1	2	243	220
Inhalants	136	95	1	4	133	91
Nonmedical Use of Psychotherapeutics ^{3,4}	2,179	2,285	24	28	2,142	2,240
Pain Relievers	1,493	1,541	20	15	1,461	1,512
OxyContin [®]	145 ^a	96	3	3	141 ^a	92
Tranquilizers	727	689	3	8	721	675
Stimulants ⁴	385 ^a	492	4	10	380	479
Methamphetamine ⁴	131	134	*	10	130	121
Sedatives	75	90	0	1	74	89
ILLCIT DRUGS OTHER THAN MARIJUANA²	2,695	2,723	30	35	2,650	2,672

*Low precision; no estimate reported.

^a Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.71B Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
ILLCIT DRUGS²	10.5 ^a	11.2	5.0	5.4	10.8 ^a	11.4
Marijuana and Hashish	8.0 ^b	8.7	4.1	4.9	8.2 ^b	8.9
Cocaine	0.6	0.5	0.1	0.2	0.6	0.6
Crack	0.1	0.1	0.1	*	0.1	0.1
Heroin	0.1	0.1	0.2	0.2	0.1	0.1
Hallucinogens	0.6	0.6	0.1	0.1	0.6	0.6
LSD	0.0	0.1	*	*	0.0	0.1
PCP	0.0	0.0	0.0	*	0.0	0.0
Ecstasy	0.4	0.4	0.1	0.1	0.4	0.4
Inhalants	0.2	0.2	0.0	0.2	0.2	0.2
Nonmedical Use of Psychotherapeutics ^{3,4}	3.5	3.6	1.0	1.3	3.6	3.7
Pain Relievers	2.4	2.4	0.9	0.6	2.5	2.5
OxyContin [®]	0.2 ^a	0.2	0.1	0.2	0.2 ^a	0.2
Tranquilizers	1.2	1.1	0.1	0.4	1.2	1.1
Stimulants ⁴	0.6	0.8	0.2	0.4	0.6	0.8
Methamphetamine ⁴	0.2	0.2	*	0.4	0.2	0.2
Sedatives	0.1	0.1	0.0	0.1	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	4.4	4.3	1.3	1.5	4.5	4.4

*Low precision; no estimate reported.

^a Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.72A Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	6,530 ^a	7,039	113	122	6,390 ^a	6,886
AGE						
15-17	867	807	12	8	848	794
18-25	2,906	2,972	71	69	2,817	2,881
26-44	2,757 ^b	3,260	30	45	2,726 ^b	3,211
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	5,530 ^a	5,988	88	98	5,422 ^a	5,864
White	4,215	4,451	59	63	4,143	4,371
Black or African American	967	1,105	25	32	938	1,064
American Indian or Alaska Native	51	39	*	*	50	39
Native Hawaiian or Other Pacific Islander	20	30	*	*	19	30
Asian	140	148	*	*	140	146
Two or More Races	137 ^b	214	*	*	133 ^b	213
Hispanic or Latino	1,000	1,051	25	24	968	1,023
TRIMESTER²						
First	N/A	N/A	62	66	N/A	N/A
Second	N/A	N/A	37	38	N/A	N/A
Third	N/A	N/A	14	17	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.72B Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	10.5 ^a	11.2	5.0	5.4	10.8 ^a	11.4
AGE						
15-17	14.0	13.0	20.9	14.6	13.9	13.0
18-25	17.1	17.2	8.2	8.6	17.6	17.6
26-44	7.1 ^b	8.3	2.2	3.2	7.3 ^b	8.5
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	10.9 ^b	11.9	4.9	5.4	11.2 ^b	12.1
White	11.5	12.3	4.5	5.0	11.8	12.6
Black or African American	11.0	12.8	7.6	8.8	11.2	12.9
American Indian or Alaska Native	13.2	10.4	*	*	13.4	10.8
Native Hawaiian or Other Pacific Islander	7.3	12.3	*	*	7.2	13.1
Asian	4.0	3.7	*	*	4.1	3.8
Two or More Races	13.7	18.5	*	*	13.9 ^a	19.2
Hispanic or Latino	8.8	8.4	5.3	5.8	8.9	8.5
TRIMESTER²						
First	N/A	N/A	9.2	9.0	N/A	N/A
Second	N/A	N/A	4.4	4.8	N/A	N/A
Third	N/A	N/A	1.8	2.4	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.73A Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	4,983 ^b	5,498	94	110	4,869 ^b	5,368
AGE						
15-17	675	677	11	8	659	666
18-25	2,432	2,482	58	58	2,359	2,409
26-44	1,876 ^b	2,339	25	44	1,851 ^b	2,293
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	4,296 ^b	4,736	77	90	4,205 ^b	4,630
White	3,281	3,497	51	59	3,221	3,430
Black or African American	768	884	23	29	741	848
American Indian or Alaska Native	32	31	*	*	32	31
Native Hawaiian or Other Pacific Islander	14	26	*	*	13	26
Asian	89	111	*	*	89	111
Two or More Races	112 ^b	185	*	*	109 ^b	184
Hispanic or Latino	687	763	17	21	664	737
TRIMESTER²						
First	N/A	N/A	55	59	N/A	N/A
Second	N/A	N/A	25	36	N/A	N/A
Third	N/A	N/A	12	14	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.73B Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	8.0 ^b	8.7	4.1	4.9	8.2 ^b	8.9
AGE						
15-17	10.9	10.9	18.5	14.6	10.8	10.9
18-25	14.3	14.4	6.7	7.2	14.7	14.7
26-44	4.8 ^b	5.9	1.8	3.2	5.0 ^b	6.1
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	8.5 ^b	9.4	4.3	4.9	8.7 ^b	9.6
White	9.0 ^a	9.7	3.9	4.7	9.2	9.9
Black or African American	8.8	10.2	7.0	8.1	8.8	10.3
American Indian or Alaska Native	8.4	8.3	*	*	8.7	8.6
Native Hawaiian or Other Pacific Islander	5.2	10.7	*	*	5.0	11.5
Asian	2.5	2.8	*	*	2.6	2.9
Two or More Races	11.1 ^a	16.0	*	*	11.3 ^a	16.7
Hispanic or Latino	6.0	6.1	3.5	5.0	6.1	6.1
TRIMESTER²						
First	N/A	N/A	8.2	8.1	N/A	N/A
Second	N/A	N/A	3.0	4.6	N/A	N/A
Third	N/A	N/A	1.6	1.9	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.74A Illicit Drug, Tobacco Product, and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
ILLCIT DRUGS²	6,530 ^a	7,039	113	122	6,390 ^a	6,886
TOBACCO PRODUCTS³	16,519	15,997	409	355	16,056	15,569
Cigarettes	15,555	14,937	401	345	15,109	14,524
Smokeless Tobacco	326	281	5	10	318	270
Cigars	2,212	2,021	34	24	2,166	1,975
Pipe Tobacco	316	349	2	5	308	340
ALCOHOL	33,089	33,797	214	212	32,752	33,448
Binge Alcohol Use ⁴	14,665	15,009	58	51	14,555	14,887
Heavy Alcohol Use ⁴	3,166	3,195	10	9	3,149	3,175
ILLCIT DRUGS, TOBACCO PRODUCTS, OR ALCOHOL^{2,3}	38,761	39,376	602	563	38,019	38,641
ILLCIT DRUGS OR ALCOHOL²	34,417	35,227	301	297	33,983	34,789
TOBACCO PRODUCTS OR ALCOHOL³	38,127	38,656	566	522	37,423	37,965

*Low precision; no estimate reported.

^a Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.74B Illicit Drug, Tobacco Product, and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
ILLCIT DRUGS²	10.5 ^a	11.2	5.0	5.4	10.8 ^a	11.4
TOBACCO PRODUCTS³	26.7 ^a	25.4	18.0	15.8	27.0 ^a	25.8
Cigarettes	25.1 ^b	23.7	17.6	15.4	25.4 ^b	24.0
Smokeless Tobacco	0.5	0.4	0.2	0.5	0.5	0.4
Cigars	3.6 ^a	3.2	1.5	1.1	3.6 ^a	3.3
Pipe Tobacco	0.5	0.6	0.1	0.2	0.5	0.6
ALCOHOL	53.4	53.7	9.4	9.4	55.1	55.4
Binge Alcohol Use ⁴	23.7	23.9	2.6	2.3	24.5	24.6
Heavy Alcohol Use ⁴	5.1	5.1	0.4	0.4	5.3	5.3
ILLCIT DRUGS, TOBACCO PRODUCTS, OR ALCOHOL^{2,3}	62.6	62.6	26.4	25.1	64.0	64.0
ILLCIT DRUGS OR ALCOHOL²	55.6	56.0	13.2	13.2	57.2	57.6
TOBACCO PRODUCTS OR ALCOHOL³	61.5	61.4	24.8	23.2	63.0	62.9

*Low precision; no estimate reported.

^a Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.75A Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	15,555	14,937	401	345	15,109	14,524
AGE						
15-17	770 ^b	610	*	*	748 ^b	592
18-25	5,010 ^b	4,488	194	170	4,790 ^b	4,291
26-44	9,775	9,839	194	162	9,571	9,641
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	13,712	13,177	387	317	13,285	12,796
White	11,143	10,557	306	244	10,806	10,273
Black or African American	1,677	1,740	54	51	1,616	1,674
American Indian or Alaska Native	150	193	*	*	146	179
Native Hawaiian or Other Pacific Islander	*	60	*	*	*	59
Asian	351	280	*	*	351	280
Two or More Races	336	346	*	*	318	331
Hispanic or Latino	1,844	1,759	14	29	1,824	1,728
TRIMESTER²						
First	N/A	N/A	174	145	N/A	N/A
Second	N/A	N/A	127	106	N/A	N/A
Third	N/A	N/A	99	93	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.75B Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	25.1 ^b	23.7	17.6	15.4	25.4 ^b	24.0
AGE						
15-17	12.4 ^b	9.8	*	*	12.3 ^b	9.7
18-25	29.5 ^b	26.0	22.4	21.0	29.9 ^b	26.2
26-44	25.2	25.0	14.3	11.8	25.7	25.4
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	27.1	26.1	21.4	17.3	27.4	26.4
White	30.5	29.3	23.4	19.3	30.8	29.6
Black or African American	19.1	20.1	16.2	14.2	19.2	20.3
American Indian or Alaska Native	39.1 ^a	50.9	*	*	39.2	49.9
Native Hawaiian or Other Pacific Islander	*	24.5	*	*	*	25.5
Asian	10.0 ^a	7.1	*	*	10.3 ^a	7.4
Two or More Races	33.5	29.8	*	*	33.2	29.9
Hispanic or Latino	16.2 ^a	14.1	2.9	7.0	16.8 ^a	14.4
TRIMESTER²						
First	N/A	N/A	25.8	19.9	N/A	N/A
Second	N/A	N/A	15.2	13.4	N/A	N/A
Third	N/A	N/A	13.0	12.8	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.76A Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	33,089	33,797	214	212	32,752	33,448
AGE						
15-17	1,329	1,272	5	*	1,316	1,257
18-25	9,752	9,892	71	50	9,636	9,791
26-44	22,008	22,633	138	154	21,800	22,400
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	28,256	28,584	191	187	27,963	28,277
White	22,021	21,994	145	138	21,794	21,779
Black or African American	3,998	4,268	28	41	3,951	4,210
American Indian or Alaska Native	170	188	*	*	169	185
Native Hawaiian or Other Pacific Islander	*	102	*	*	*	102
Asian	1,409	1,320	*	*	1,397	1,304
Two or More Races	558 ^a	713	*	*	551 ^a	698
Hispanic or Latino	4,833	5,212	23	25	4,789	5,171
TRIMESTER²						
First	N/A	N/A	124	138	N/A	N/A
Second	N/A	N/A	67	39	N/A	N/A
Third	N/A	N/A	23	32	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.76B Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	53.4	53.7	9.4	9.4	55.1	55.4
AGE						
15-17	21.4	20.5	8.8	*	21.6	20.5
18-25	57.5	57.2	8.2	6.2	60.1	59.8
26-44	56.7	57.4	10.2	11.2	58.5	59.1
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	55.9	56.7	10.6	10.2	57.6	58.4
White	60.2	61.0	11.0	10.9	62.0	62.8
Black or African American	45.6 ^b	49.4	8.4	11.3	47.1 ^b	51.1
American Indian or Alaska Native	44.2	49.5	*	*	45.6	51.5
Native Hawaiian or Other Pacific Islander	*	41.4	*	*	*	44.1
Asian	40.0 ^a	33.4	*	*	41.1 ^a	34.3
Two or More Races	55.6	61.4	*	*	57.5	63.1
Hispanic or Latino	42.4	41.8	4.8	6.0	44.0	43.1
TRIMESTER²						
First	N/A	N/A	18.4	19.0	N/A	N/A
Second	N/A	N/A	7.9	5.0	N/A	N/A
Third	N/A	N/A	3.0	4.4	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.77A Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	14,665	15,009	58	51	14,555	14,887
AGE						
15-17	722	667	4	*	714	662
18-25	5,583	5,579	27	25	5,529	5,524
26-44	8,360	8,763	28	24	8,312	8,701
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	12,276	12,453	56	40	12,182	12,353
White	9,876	9,811	42 ^a	18	9,803	9,750
Black or African American	1,581	1,751	12	14	1,563	1,727
American Indian or Alaska Native	110	113	*	*	110	109
Native Hawaiian or Other Pacific Islander	47	54	*	*	46	54
Asian	443	425	*	*	443	423
Two or More Races	219 ^a	299	*	*	218	290
Hispanic or Latino	2,389	2,556	2	10	2,373	2,533
TRIMESTER²						
First	N/A	N/A	45	35	N/A	N/A
Second	N/A	N/A	11	14	N/A	N/A
Third	N/A	N/A	2	1	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

^a Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.77B Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	23.7	23.9	2.6	2.3	24.5	24.6
AGE						
15-17	11.6	10.7	6.4	*	11.7	10.8
18-25	32.9	32.3	3.1	3.1	34.5	33.7
26-44	21.6	22.2	2.1	1.8	22.3	23.0
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	24.3	24.7	3.1	2.2	25.1	25.5
White	27.0	27.2	3.2 ^a	1.4	27.9	28.1
Black or African American	18.0 ^a	20.3	3.5	3.8	18.6 ^a	21.0
American Indian or Alaska Native	28.7	29.7	*	*	29.5	30.5
Native Hawaiian or Other Pacific Islander	16.9	21.9	*	*	17.1	23.3
Asian	12.6	10.7	*	*	13.0	11.1
Two or More Races	21.8	25.8	*	*	22.7	26.2
Hispanic or Latino	20.9	20.5	0.5	2.5	21.8	21.1
TRIMESTER²						
First	N/A	N/A	6.7	4.8	N/A	N/A
Second	N/A	N/A	1.3	1.8	N/A	N/A
Third	N/A	N/A	0.3	0.1	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

^a Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.78A Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	3,166	3,195	10	9	3,149	3,175
AGE						
15-17	135	119	1	*	134	118
18-25	1,397	1,325	9	2	1,384	1,315
26-44	1,634	1,752	*	7	1,631	1,741
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	2,785	2,789	8	9	2,772	2,770
White	2,368	2,290	5	4	2,358	2,277
Black or African American	250	317	2	5	248	312
American Indian or Alaska Native	45	27	*	*	45	27
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	55	84	*	*	55	83
Two or More Races	51	55	*	*	50	55
Hispanic or Latino	381	406	2	*	377	405
TRIMESTER²						
First	N/A	N/A	6	4	N/A	N/A
Second	N/A	N/A	4	5	N/A	N/A
Third	N/A	N/A	*	0	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

^a Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.78B Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	5.1	5.1	0.4	0.4	5.3	5.3
AGE						
15-17	2.2	1.9	1.6	*	2.2	1.9
18-25	8.2	7.7	1.0	0.3	8.6	8.0
26-44	4.2	4.4	*	0.5	4.4	4.6
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	5.5	5.5	0.4	0.5	5.7	5.7
White	6.5	6.4	0.4	0.3	6.7	6.6
Black or African American	2.9	3.7	0.7	1.3	3.0	3.8
American Indian or Alaska Native	11.9	7.0	*	*	12.2	7.5
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.6	2.1	*	*	1.6	2.2
Two or More Races	5.1	4.7	*	*	5.2	4.9
Hispanic or Latino	3.3	3.3	0.4	*	3.5	3.4
TRIMESTER²						
First	N/A	N/A	0.8	0.6	N/A	N/A
Second	N/A	N/A	0.5	0.6	N/A	N/A
Third	N/A	N/A	*	0.0	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

^a Difference between estimate and 2012-2013 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2012-2013 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.