

Table 6.19C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2012 and 2013

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLCIT DRUGS¹	311	323	150	187	322	309	150	159
Marijuana and Hashish	282	296	126	171	281	268	110	112
Cocaine	123	106	36	36	58	70	31	25
Crack	82	52	32	12	25	25	20	24
Heroin	37	33	23	19	4	*	25	18
Hallucinogens	83	78	16	14	29	92	13	32
LSD	30	20	4	7	7	30	5	3
PCP	5	11	2	1	11	*	*	10
Ecstasy	68	56	8	7	23	52	8	11
Inhalants	26	41	7	21	55	35	18	27
Nonmedical Use of Psychotherapeutics ^{2,3}	151	165	82	92	165	133	105	109
Pain Relievers	119	139	67	78	123	104	92	89
OxyContin [®]	35	46	25	38	8	22	12	51
Tranquilizers	104	81	44	48	96	60	68	38
Stimulants ³	67	91	37	37	64	50	11	33
Methamphetamine ³	58	68	32	34	31	25	7	27
Sedatives	13	26	17	8	43	45	12	44
ILLCIT DRUGS OTHER THAN MARIJUANA¹	195	201	91	103	182	173	109	115

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.19D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2012 and 2013

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLCIT DRUGS¹	0.70	0.74	0.72	0.89	0.32	0.30	0.14	0.15
Marijuana and Hashish	0.66	0.69	0.62	0.82	0.28	0.26	0.10	0.10
Cocaine	0.32	0.29	0.18	0.19	0.06	0.07	0.03	0.02
Crack	0.22	0.14	0.16	0.06	0.03	0.02	0.02	0.02
Heroin	0.10	0.09	0.11	0.10	0.00	*	0.02	0.02
Hallucinogens	0.22	0.21	0.08	0.07	0.03	0.09	0.01	0.03
LSD	0.08	0.06	0.02	0.04	0.01	0.03	0.00	0.00
PCP	0.01	0.03	0.01	0.01	0.01	*	*	0.01
Ecstasy	0.18	0.15	0.04	0.04	0.02	0.05	0.01	0.01
Inhalants	0.07	0.11	0.04	0.11	0.06	0.03	0.02	0.03
Nonmedical Use of Psychotherapeutics ^{2,3}	0.38	0.43	0.41	0.47	0.17	0.13	0.10	0.10
Pain Relievers	0.31	0.37	0.34	0.40	0.13	0.10	0.09	0.08
OxyContin [®]	0.09	0.13	0.12	0.19	0.01	0.02	0.01	0.05
Tranquilizers	0.27	0.22	0.22	0.25	0.10	0.06	0.06	0.04
Stimulants ³	0.18	0.25	0.18	0.19	0.07	0.05	0.01	0.03
Methamphetamine ³	0.15	0.19	0.16	0.18	0.03	0.02	0.01	0.03
Sedatives	0.03	0.07	0.09	0.04	0.04	0.04	0.01	0.04
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.49	0.51	0.45	0.52	0.18	0.17	0.10	0.11

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.19P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLCIT DRUGS¹	0.1372		0.5892		0.5940		0.9529	
Marijuana and Hashish	0.1257		0.5599		0.2563		0.7680	
Cocaine	0.6062		0.3143		0.3735		0.7385	
Crack	0.7477		0.0949		0.7006		0.5760	
Heroin	0.9922		0.5652		*		0.5598	
Hallucinogens	0.9306		0.7660		0.0451		0.3822	
LSD	0.8650		0.2220		0.1309		0.8201	
PCP	0.7078		0.9685		*		*	
Ecstasy	0.6712		0.9241		0.1558		0.9933	
Inhalants	0.6473		0.3476		0.3788		0.5574	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.7532		0.4261		0.0709		0.5727	
Pain Relievers	0.8046		0.5703		0.2104		0.3857	
OxyContin [®]	0.6357		0.5423		0.1496		0.3134	
Tranquilizers	0.2729		0.7720		0.1042		0.1755	
Stimulants ³	0.0660		0.4410		0.2181		0.0884	
Methamphetamine ³	0.2230		0.4956		0.5732		0.0931	
Sedatives	0.5128		0.7281		0.1999		0.3026	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.6722		0.5128		0.2622		0.6777	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.20C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2012 and 2013

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLICIT DRUGS¹	36	29	22	21	33	35	45	42
Marijuana and Hashish	34	28	21	20	32	34	32	32
Cocaine	6	6	3	3	1	3	1	5
Crack	1	1	*	*	0	1	*	*
Heroin	*	3	*	3	*	1	*	1
Hallucinogens	14	11	8	7	3	7	8	8
LSD	5	8	3	4	1	3	1	3
PCP	2	3	2	1	0	1	*	*
Ecstasy	9	6	5	3	3	4	4	4
Inhalants	11	5	4	4	8	5	13	13
Nonmedical Use of Psychotherapeutics ^{2,3}	18	15	9	8	15	12	29	25
Pain Relievers	15	13	8	7	14	10	25	23
OxyContin [®]	8	4	1	0	6	4	3	3
Tranquilizers	10	8	5	1	5	3	15	8
Stimulants ³	10	6	5	5	6	6	6	6
Methamphetamine ³	4	2	2	4	3	3	3	3
Sedatives	4	3	1	1	1	1	5	6
ILLICIT DRUGS OTHER THAN MARIJUANA¹	22	18	12	11	16	13	33	29

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.20D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2012 and 2013

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLCIT DRUGS¹	2.13	2.20	2.47	2.63	1.31	1.40	0.20	0.19
Marijuana and Hashish	2.11	2.32	2.43	2.56	1.28	1.37	0.15	0.15
Cocaine	0.62	0.76	0.43	0.49	0.05	0.15	0.01	0.03
Crack	0.07	0.06	*	*	0.01	0.05	*	*
Heroin	*	0.44	*	0.47	*	0.05	*	0.01
Hallucinogens	1.29	1.39	1.17	1.07	0.16	0.34	0.04	0.04
LSD	0.49	1.01	0.41	0.62	0.03	0.16	0.01	0.01
PCP	0.23	0.38	0.27	0.17	0.01	0.04	*	*
Ecstasy	0.92	0.78	0.72	0.53	0.13	0.20	0.02	0.02
Inhalants	1.03	0.66	0.65	0.58	0.36	0.23	0.06	0.06
Nonmedical Use of Psychotherapeutics ^{2,3}	1.62	1.71	1.37	1.29	0.65	0.55	0.13	0.12
Pain Relievers	1.37	1.54	1.15	1.08	0.59	0.50	0.11	0.11
OxyContin [®]	0.74	0.51	0.17	0.05	0.26	0.17	0.02	0.02
Tranquilizers	0.96	1.03	0.77	0.17	0.23	0.15	0.07	0.04
Stimulants ³	1.00	0.74	0.81	0.81	0.26	0.26	0.03	0.03
Methamphetamine ³	0.44	0.30	0.38	0.59	0.12	0.14	0.01	0.02
Sedatives	0.38	0.34	0.22	0.24	0.05	0.06	0.02	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1.89	1.96	1.79	1.69	0.72	0.63	0.15	0.13

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.20P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLCIT DRUGS¹	0.2693		0.2742		0.7858		0.6080	
Marijuana and Hashish	0.2293		0.6239		0.2184		0.2946	
Cocaine	0.6581		0.8477		0.0774		0.1251	
Crack	0.6965		*		0.3851		*	
Heroin	*		*		*		*	
Hallucinogens	0.9527		0.8690		0.0907		0.9684	
LSD	0.0306		0.5189		0.1806		0.3248	
PCP	0.5693		0.7412		0.1981		*	
Ecstasy	0.0775		0.6894		0.2840		0.9120	
Inhalants	0.0296		0.6482		0.0570		0.4275	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.7086		0.3088		0.0400		0.2035	
Pain Relievers	0.6448		0.2326		0.0408		0.1315	
OxyContin [®]	0.3279		0.1118		0.5225		0.9706	
Tranquilizers	0.8983		0.0016		0.0938		0.5005	
Stimulants ³	0.0475		0.7658		0.7186		0.9650	
Methamphetamine ³	0.1689		0.6987		0.8695		0.9144	
Sedatives	0.6499		0.9133		0.4009		0.8787	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.9435		0.6224		0.0305		0.2258	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.21C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2012 and 2013

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLCIT DRUGS¹	118	117	50	40	93	91	42	45
Marijuana and Hashish	115	111	44	37	89	84	36	36
Cocaine	29	29	11	6	12	16	3	2
Crack	7	11	4	5	4	*	2	*
Heroin	14	11	13	12	4	1	11	1
Hallucinogens	39	37	9	11	19	23	9	15
LSD	15	15	2	4	7	8	5	2
PCP	3	1	0	*	3	*	*	2
Ecstasy	35	27	5	6	15	15	6	4
Inhalants	18	12	4	10	6	8	11	6
Nonmedical Use of Psychotherapeutics ^{2,3}	58	57	29	18	36	36	18	22
Pain Relievers	49	49	29	15	27	28	17	20
OxyContin [®]	17	17	16	4	5	7	*	*
Tranquilizers	33	29	23	11	16	17	5	7
Stimulants ³	24	28	7	8	17	21	5	7
Methamphetamine ³	14	17	5	4	3	6	2	2
Sedatives	5	6	1	*	11	6	3	4
ILLCIT DRUGS OTHER THAN MARIJUANA¹	71	67	31	23	40	42	24	27

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.21D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2012 and 2013

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLCIT DRUGS¹	0.90	0.98	1.59	1.45	0.61	0.59	0.36	0.37
Marijuana and Hashish	0.92	0.96	1.49	1.37	0.59	0.55	0.32	0.31
Cocaine	0.33	0.34	0.41	0.24	0.09	0.13	0.02	0.02
Crack	0.08	0.13	0.15	0.21	0.03	*	0.02	*
Heroin	0.16	0.13	0.50	0.48	0.03	0.01	0.10	0.01
Hallucinogens	0.45	0.44	0.34	0.46	0.15	0.19	0.08	0.13
LSD	0.18	0.18	0.09	0.17	0.06	0.07	0.04	0.02
PCP	0.04	0.01	0.01	*	0.03	*	*	0.02
Ecstasy	0.40	0.32	0.18	0.25	0.12	0.12	0.05	0.04
Inhalants	0.22	0.14	0.14	0.41	0.05	0.07	0.10	0.05
Nonmedical Use of Psychotherapeutics ^{2,3}	0.62	0.61	1.07	0.73	0.27	0.27	0.16	0.18
Pain Relievers	0.54	0.55	1.07	0.60	0.22	0.22	0.15	0.17
OxyContin [®]	0.20	0.20	0.60	0.15	0.04	0.06	*	*
Tranquilizers	0.38	0.33	0.86	0.47	0.12	0.13	0.04	0.06
Stimulants ³	0.28	0.33	0.27	0.33	0.13	0.16	0.05	0.06
Methamphetamine ³	0.16	0.21	0.19	0.16	0.02	0.05	0.01	0.01
Sedatives	0.06	0.07	0.05	*	0.09	0.05	0.03	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.73	0.70	1.12	0.91	0.30	0.32	0.21	0.22

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.21P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLCIT DRUGS¹	0.3604		0.1398		0.1899		0.6008	
Marijuana and Hashish	0.3709		0.5743		0.1787		0.6588	
Cocaine	0.7827		0.4833		0.7705		0.6249	
Crack	0.6156		0.5618		*		*	
Heroin	0.3974		0.7812		0.1575		0.1164	
Hallucinogens	0.8870		0.2462		0.9298		0.8224	
LSD	0.6276		0.5662		0.9810		0.4180	
PCP	0.1914		*		*		*	
Ecstasy	0.6696		0.3883		0.3222		0.5374	
Inhalants	0.1860		0.3086		0.4380		0.6449	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.1602		0.0494		0.7183		0.2540	
Pain Relievers	0.1621		0.0161		0.6372		0.5147	
OxyContin [®]	0.6558		0.0843		0.7764		*	
Tranquilizers	0.0285		0.6879		0.8674		0.6712	
Stimulants ³	0.5006		0.6119		0.3065		0.2242	
Methamphetamine ³	0.4938		0.9900		0.1203		0.8387	
Sedatives	0.8081		*		0.1917		0.6401	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.7641		0.2914		0.6754		0.5895	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.22C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2012 and 2013

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLCIT DRUGS¹	289	301	137	181	306	281	135	146
Marijuana and Hashish	259	274	114	164	266	244	99	99
Cocaine	119	102	35	36	57	68	31	25
Crack	82	51	32	11	25	25	20	24
Heroin	34	32	18	15	*	*	23	18
Hallucinogens	73	67	11	6	20	90	7	27
LSD	26	11	*	4	*	29	*	*
PCP	4	11	*	1	11	*	*	10
Ecstasy	59	49	5	*	17	51	5	10
Inhalants	13	39	5	18	54	34	6	23
Nonmedical Use of Psychotherapeutics ^{2,3}	139	155	76	90	159	128	97	104
Pain Relievers	112	129	59	77	120	101	86	84
OxyContin [®]	30	41	19	37	4	20	11	51
Tranquilizers	97	74	38	47	94	57	66	37
Stimulants ³	61	85	36	36	63	44	8	28
Methamphetamine ³	55	64	32	34	30	24	6	27
Sedatives	12	25	17	8	41	44	10	43
ILLCIT DRUGS OTHER THAN MARIJUANA¹	179	188	83	99	174	168	99	108

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.22D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2012 and 2013

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLCIT DRUGS¹	0.88	0.92	0.79	1.03	0.36	0.32	0.18	0.20
Marijuana and Hashish	0.82	0.86	0.67	0.95	0.31	0.28	0.14	0.13
Cocaine	0.42	0.37	0.21	0.22	0.07	0.08	0.04	0.03
Crack	0.29	0.19	0.19	0.07	0.03	0.03	0.03	0.03
Heroin	0.12	0.12	0.11	0.09	*	*	0.03	0.02
Hallucinogens	0.26	0.24	0.07	0.03	0.02	0.10	0.01	0.04
LSD	0.09	0.04	*	0.02	*	0.03	*	*
PCP	0.01	0.04	*	0.01	0.01	*	*	0.01
Ecstasy	0.21	0.18	0.03	*	0.02	0.06	0.01	0.01
Inhalants	0.04	0.14	0.03	0.11	0.07	0.04	0.01	0.03
Nonmedical Use of Psychotherapeutics ^{2,3}	0.47	0.54	0.45	0.54	0.19	0.15	0.13	0.14
Pain Relievers	0.38	0.46	0.35	0.46	0.14	0.12	0.12	0.11
OxyContin [®]	0.10	0.15	0.11	0.23	0.00	0.02	0.02	0.07
Tranquilizers	0.34	0.27	0.23	0.28	0.11	0.07	0.09	0.05
Stimulants ³	0.22	0.31	0.21	0.22	0.08	0.05	0.01	0.04
Methamphetamine ³	0.20	0.23	0.19	0.20	0.04	0.03	0.01	0.04
Sedatives	0.04	0.09	0.10	0.05	0.05	0.05	0.01	0.06
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.60	0.65	0.49	0.59	0.21	0.20	0.14	0.15

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.22P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes But Not Alcohol		Alcohol But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2012	2013	2012	2013	2012	2013	2012	2013
ILLCIT DRUGS¹	0.1865		0.2443		0.6974		0.8935	
Marijuana and Hashish	0.1743		0.3518		0.3726		0.8301	
Cocaine	0.5203		0.3798		0.4328		0.6125	
Crack	0.6872		0.0715		0.5666		0.5299	
Heroin	0.8421		0.5432		*		0.9229	
Hallucinogens	0.9433		0.4132		0.0416		0.3642	
LSD	0.6386		*		*		*	
PCP	0.5313		*		*		*	
Ecstasy	0.9620		*		0.0652		0.7109	
Inhalants	0.0509		0.4961		0.4741		0.0926	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.3482		0.1259		0.1178		0.6106	
Pain Relievers	0.7111		0.1266		0.3594		0.4967	
OxyContin [®]	0.3313		0.1824		0.0786		0.3119	
Tranquilizers	0.7076		0.4394		0.1106		0.1999	
Stimulants ³	0.0495		0.5134		0.1136		0.1019	
Methamphetamine ³	0.2326		0.5135		0.4232		0.0949	
Sedatives	0.3832		0.7859		0.2881		0.3348	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.5142		0.2739		0.4022		0.8017	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.23C Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: Standard Errors of Numbers in Thousands, 2012 and 2013

Cigarette Use/Level of Alcohol Use	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
TOTAL	492	509	63	62	138	143	452	472
Heavy Alcohol Use ¹	233	232	21	17	84	82	215	217
Binge Alcohol Use But Not Heavy Use ¹	270	267	32	35	94	103	246	237
Alcohol Use But Not Binge Use ¹	257	270	28	24	69	64	247	261
No Alcohol Use	217	245	53	47	65	61	195	231
PAST MONTH CIGARETTE USE	352	372	42	36	133	126	325	348
Heavy Alcohol Use ¹	179	176	18	13	68	67	166	168
Binge Alcohol Use But Not Heavy Use ¹	196	194	24	23	71	77	182	175
Alcohol Use But Not Binge Use ¹	143	173	16	12	48	43	131	165
No Alcohol Use	150	187	22	21	50	40	137	181
NO PAST MONTH CIGARETTE USE	349	355	56	55	103	103	329	320
Heavy Alcohol Use ¹	136	140	11	*	45	44	127	130
Binge Alcohol Use But Not Heavy Use ¹	172	170	22	27	55	62	161	149
Alcohol Use But Not Binge Use ¹	213	208	22	20	47	44	211	200
No Alcohol Use	150	159	45	42	42	45	135	146

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.23D Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: Standard Errors of Percentages, 2012 and 2013

Cigarette Use/Level of Alcohol Use	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
TOTAL	0.19	0.19	0.25	0.25	0.40	0.41	0.23	0.23
Heavy Alcohol Use ¹	1.12	1.17	3.64	4.26	1.28	1.35	1.47	1.49
Binge Alcohol Use But Not Heavy Use ¹	0.57	0.57	1.72	2.08	0.77	0.86	0.71	0.67
Alcohol Use But Not Binge Use ¹	0.33	0.34	1.60	1.61	0.84	0.79	0.36	0.37
No Alcohol Use	0.17	0.19	0.23	0.20	0.44	0.41	0.22	0.25
PAST MONTH CIGARETTE USE	0.53	0.58	1.61	1.70	0.81	0.84	0.63	0.70
Heavy Alcohol Use ¹	1.52	1.65	4.45	4.43	1.56	1.65	2.08	2.19
Binge Alcohol Use But Not Heavy Use ¹	1.05	1.10	2.91	3.36	1.26	1.38	1.38	1.40
Alcohol Use But Not Binge Use ¹	1.08	1.24	4.33	4.73	2.04	1.91	1.17	1.43
No Alcohol Use	0.72	0.89	2.47	2.63	1.59	1.45	0.79	1.03
NO PAST MONTH CIGARETTE USE	0.17	0.17	0.22	0.22	0.37	0.37	0.21	0.20
Heavy Alcohol Use ¹	1.54	1.62	5.75	*	2.03	2.15	1.91	1.94
Binge Alcohol Use But Not Heavy Use ¹	0.61	0.59	2.01	2.41	0.90	0.96	0.75	0.65
Alcohol Use But Not Binge Use ¹	0.33	0.32	1.66	1.58	0.81	0.75	0.37	0.34
No Alcohol Use	0.14	0.15	0.20	0.19	0.36	0.37	0.18	0.20

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.23P Illicit Drug Use in the Past Month, by Past Month Cigarette Use, Levels of Past Month Alcohol Use, and Age Group: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Cigarette Use/Level of Alcohol Use	Total (2012)	Total (2013)	Aged 12-17 (2012)	Aged 12-17 (2013)	Aged 18-25 (2012)	Aged 18-25 (2013)	Aged 26+ (2012)	Aged 26+ (2013)
TOTAL	0.4809		0.0476		0.7348		0.3451	
Heavy Alcohol Use ¹	0.0968		0.3020		0.3444		0.0635	
Binge Alcohol Use But Not Heavy Use ¹	0.9645		0.0840		0.0009		0.3040	
Alcohol Use But Not Binge Use ¹	0.7327		0.2848		0.2745		0.4999	
No Alcohol Use	0.8904		0.3343		0.2345		0.5086	
PAST MONTH CIGARETTE USE	0.1399		0.7501		0.6943		0.1008	
Heavy Alcohol Use ¹	0.2786		0.4120		0.1895		0.2903	
Binge Alcohol Use But Not Heavy Use ¹	0.4159		0.2194		0.0632		0.9182	
Alcohol Use But Not Binge Use ¹	0.3864		0.4509		0.1247		0.2203	
No Alcohol Use	0.5892		0.2742		0.1398		0.2443	
NO PAST MONTH CIGARETTE USE	0.6021		0.4808		0.2858		0.7070	
Heavy Alcohol Use ¹	0.1769		*		0.8414		0.0915	
Binge Alcohol Use But Not Heavy Use ¹	0.9670		0.0720		0.0018		0.3969	
Alcohol Use But Not Binge Use ¹	0.9625		0.2742		0.7501		0.9123	
No Alcohol Use	0.9529		0.6080		0.6008		0.8935	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.