

**Table 6.24C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2012 and 2013**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	366	359
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	174	160	247	256
Cigars	268	272	284	244
Pipe Tobacco	120	120	140	93
<b>ALCOHOL</b>	700	675	1,532	1,519
Binge Alcohol Use <sup>2</sup>	527	517	728	762
Heavy Alcohol Use <sup>2</sup>	311	308	334	313

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 6.24D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2012 and 2013**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	0.18	0.17
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.29	0.28	0.12	0.12
Cigars	0.42	0.46	0.14	0.12
Pipe Tobacco	0.20	0.21	0.07	0.04
<b>ALCOHOL</b>	0.73	0.74	0.45	0.48
Binge Alcohol Use <sup>2</sup>	0.70	0.71	0.32	0.32
Heavy Alcohol Use <sup>2</sup>	0.48	0.52	0.16	0.15

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 6.24P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2013 Versus 2012**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A		0.0287	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.9780		0.5088	
Cigars	0.5717		0.0702	
Pipe Tobacco	0.4853		0.0305	
<b>ALCOHOL</b>	0.7906		0.5947	
Binge Alcohol Use <sup>2</sup>	0.4626		0.3573	
Heavy Alcohol Use <sup>2</sup>	0.9465		0.4415	

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 6.25C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2012 and 2013**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	29	34
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	23	19	20	24
Cigars	26	23	20	23
Pipe Tobacco	14	12	11	13
<b>ALCOHOL</b>	47	39	73	70
Binge Alcohol Use <sup>2</sup>	40	33	48	48
Heavy Alcohol Use <sup>2</sup>	21	16	15	18

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 6.25D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Percentages, 2012 and 2013**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	0.12	0.14
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1.25	1.24	0.08	0.10
Cigars	1.37	1.50	0.08	0.10
Pipe Tobacco	0.83	0.81	0.05	0.05
<b>ALCOHOL</b>	1.62	1.81	0.28	0.28
Binge Alcohol Use <sup>2</sup>	1.69	1.72	0.19	0.19
Heavy Alcohol Use <sup>2</sup>	1.14	1.08	0.06	0.07

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 6.25P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2013 Versus 2012**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A		0.4547	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.6720		0.5666	
Cigars	0.3738		0.7116	
Pipe Tobacco	0.9423		0.6140	
<b>ALCOHOL</b>	0.0279		0.1454	
Binge Alcohol Use <sup>2</sup>	0.1768		0.1309	
Heavy Alcohol Use <sup>2</sup>	0.9035		0.8971	

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 6.26C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2012 and 2013**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	85	86
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	50	58	49	45
Cigars	86	81	62	65
Pipe Tobacco	31	35	27	37
<b>ALCOHOL</b>	190	173	276	280
Binge Alcohol Use <sup>2</sup>	169	155	196	195
Heavy Alcohol Use <sup>2</sup>	97	90	98	77

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 6.26D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Percentages, 2012 and 2013**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	0.32	0.33
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.43	0.50	0.20	0.19
Cigars	0.64	0.63	0.24	0.26
Pipe Tobacco	0.27	0.32	0.11	0.15
<b>ALCOHOL</b>	0.74	0.64	0.62	0.65
Binge Alcohol Use <sup>2</sup>	0.87	0.81	0.57	0.59
Heavy Alcohol Use <sup>2</sup>	0.70	0.67	0.37	0.29

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 6.26P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: P Values from Tests of Differences of Percentages, 2013 Versus 2012**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A		0.8799	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.1400		0.7920	
Cigars	0.2528		0.2742	
Pipe Tobacco	0.3046		0.0424	
<b>ALCOHOL</b>	0.2441		0.2761	
Binge Alcohol Use <sup>2</sup>	0.7006		0.0457	
Heavy Alcohol Use <sup>2</sup>	0.1152		0.0299	

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 6.27C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2012 and 2013**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	350	345
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	165	148	237	251
Cigars	250	258	277	233
Pipe Tobacco	114	114	137	86
<b>ALCOHOL</b>	665	640	1,476	1,461
Binge Alcohol Use <sup>2</sup>	493	488	693	718
Heavy Alcohol Use <sup>2</sup>	296	298	318	295

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 6.27D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2012 and 2013**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	0.22	0.21
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.36	0.33	0.15	0.15
Cigars	0.52	0.55	0.18	0.14
Pipe Tobacco	0.25	0.25	0.09	0.05
<b>ALCOHOL</b>	0.90	0.93	0.56	0.58
Binge Alcohol Use <sup>2</sup>	0.85	0.87	0.40	0.40
Heavy Alcohol Use <sup>2</sup>	0.59	0.64	0.20	0.18

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

**Table 6.27P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2013 Versus 2012**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2012	2013	2012	2013
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A		0.0161	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.7472		0.4237	
Cigars	0.9228		0.0947	
Pipe Tobacco	0.6213		0.0042	
<b>ALCOHOL</b>	0.7216		0.4481	
Binge Alcohol Use <sup>2</sup>	0.5724		0.1480	
Heavy Alcohol Use <sup>2</sup>	0.7302		0.8081	

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.