

Table 6.28C Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2012 and 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	347	337	436	445	465	453	601	590
Cigarettes	311	308	385	406	422	398	564	560
Smokeless Tobacco	126	122	152	154	126	133	182	175
Cigars	160	154	205	219	202	182	188	152
Pipe Tobacco	42	56	76	83	109	74	109	85

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.28D Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2012 and 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	1.32	1.32	0.79	0.80	0.56	0.55	0.43	0.41
Cigarettes	1.33	1.27	0.74	0.77	0.53	0.49	0.41	0.40
Smokeless Tobacco	0.70	0.73	0.35	0.35	0.17	0.17	0.14	0.14
Cigars	0.87	0.87	0.46	0.48	0.26	0.24	0.15	0.12
Pipe Tobacco	0.25	0.33	0.18	0.19	0.14	0.10	0.09	0.07

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.28P Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2013 Versus 2012

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	0.3779		0.0023		0.5315		0.1861	
Cigarettes	0.8782		0.0090		0.5884		0.4034	
Smokeless Tobacco	0.7416		0.5359		0.7210		0.9858	
Cigars	0.1171		0.9026		0.3222		0.0328	
Pipe Tobacco	0.3228		0.5048		0.0362		0.2969	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.29C Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2012 and 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	23	18	35	35	25	22	39	41
Cigarettes	21	16	33	28	23	18	33	34
Smokeless Tobacco	11	12	17	15	10	10	19	21
Cigars	11	12	20	22	15	10	17	19
Pipe Tobacco	5	5	9	11	10	5	9	11

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.29D Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2012 and 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	3.38	4.13	1.81	1.98	1.53	1.49	0.18	0.18
Cigarettes	3.69	4.14	1.81	1.86	1.44	1.29	0.15	0.15
Smokeless Tobacco	3.08	3.51	1.11	1.16	0.70	0.73	0.09	0.09
Cigars	3.07	3.61	1.24	1.53	1.02	0.78	0.08	0.09
Pipe Tobacco	1.56	1.68	0.60	0.88	0.69	0.38	0.04	0.05

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.29P Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2013 Versus 2012

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	0.7131		0.8636		0.1735		0.7409	
Cigarettes	0.4880		0.1705		0.0517		0.5950	
Smokeless Tobacco	0.1185		0.8796		0.2360		0.1726	
Cigars	0.2386		0.6621		0.0871		0.8925	
Pipe Tobacco	0.6015		0.3562		0.0286		0.3819	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.30C Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2012 and 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	110	100	129	124	78	83	103	92
Cigarettes	97	90	116	110	71	70	97	83
Smokeless Tobacco	43	40	40	44	23	26	30	30
Cigars	57	57	60	60	42	41	44	38
Pipe Tobacco	19	22	23	31	18	23	19	20

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.30D Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2012 and 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	1.35	1.33	0.85	0.94	0.91	0.89	0.66	0.57
Cigarettes	1.48	1.39	0.84	0.93	0.86	0.81	0.63	0.52
Smokeless Tobacco	0.88	0.94	0.42	0.45	0.32	0.34	0.22	0.21
Cigars	1.06	1.19	0.58	0.59	0.54	0.53	0.30	0.27
Pipe Tobacco	0.41	0.53	0.24	0.32	0.25	0.29	0.14	0.14

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.30P Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2013 Versus 2012

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	0.9782		0.5655		0.8049		0.0654	
Cigarettes	0.7508		0.9778		0.8434		0.0228	
Smokeless Tobacco	0.5724		0.1250		0.2546		0.8510	
Cigars	0.4335		0.8119		0.3499		0.0936	
Pipe Tobacco	0.7069		0.0113		0.4172		0.5911	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.31C Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2012 and 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	327	322	415	419	456	438	578	571
Cigarettes	296	298	370	384	415	388	540	543
Smokeless Tobacco	115	114	144	146	125	130	178	173
Cigars	150	139	196	207	199	175	181	149
Pipe Tobacco	36	51	71	78	107	69	106	81

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.31D Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2012 and 2013

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	1.77	1.71	1.02	1.01	0.61	0.61	0.58	0.56
Cigarettes	1.77	1.65	0.96	0.97	0.58	0.54	0.55	0.54
Smokeless Tobacco	0.89	0.90	0.45	0.43	0.18	0.19	0.20	0.19
Cigars	1.13	1.06	0.59	0.59	0.28	0.26	0.20	0.16
Pipe Tobacco	0.29	0.41	0.22	0.24	0.16	0.10	0.12	0.09

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.

Table 6.31P Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2013 Versus 2012

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2012	2013	2012	2013	2012	2013	2012	2013
TOBACCO PRODUCTS²	0.4410		0.0013		0.5097		0.3158	
Cigarettes	0.9266		0.0107		0.5932		0.6661	
Smokeless Tobacco	0.5810		0.3072		0.4982		0.9140	
Cigars	0.2053		0.8352		0.4560		0.0679	
Pipe Tobacco	0.3150		0.8321		0.0297		0.2001	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012 and 2013.