

Table 6.47C Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	100	119	15	16	27	28	93	116
From More Than One Doctor	36	49	5	6	8	9	35	47
Wrote Fake Prescription	8	4	3	2	4	2	7	3
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	15	26	4	2	5	2	14	26
From Friend or Relative for Free	158	173	24	20	54	51	146	162
Bought from Friend or Relative	70	67	10	9	27	25	63	61
Took from Friend or Relative without Asking	37	41	11	9	14	17	31	37
Bought from Drug Dealer or Other Stranger	30	35	7	6	17	18	22	30
Bought on the Internet	14	5	0	1	1	4	14	1
Some Other Way	47	41	10	7	13	13	44	37

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.47D Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	0.83	0.94	1.08	1.42	0.75	0.81	1.35	1.44
From More Than One Doctor	0.33	0.44	0.40	0.61	0.23	0.28	0.57	0.69
Wrote Fake Prescription	0.08	0.04	0.24	0.24	0.11	0.05	0.12	0.05
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.14	0.24	0.33	0.24	0.16	0.08	0.22	0.38
From Friend or Relative for Free	0.96	1.05	1.43	1.57	0.98	1.14	1.61	1.62
Bought from Friend or Relative	0.60	0.58	0.76	0.91	0.70	0.74	0.93	0.85
Took from Friend or Relative without Asking	0.33	0.37	0.84	0.91	0.40	0.53	0.50	0.54
Bought from Drug Dealer or Other Stranger	0.27	0.32	0.58	0.58	0.48	0.58	0.35	0.43
Bought on the Internet	0.13	0.04	0.03	0.14	0.04	0.14	0.22	0.02
Some Other Way	0.42	0.37	0.77	0.74	0.38	0.41	0.69	0.55

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.47P Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011**

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	0.0111		0.1137		0.4126		0.0474	
From More Than One Doctor	0.2418		0.2332		0.7316		0.3515	
Wrote Fake Prescription	0.2399		0.6369		0.0856		0.8253	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3568		0.3565		0.3436		0.1975	
From Friend or Relative for Free	0.3800		0.2674		0.5144		0.4458	
Bought from Friend or Relative	0.0720		0.9737		0.2411		0.1680	
Took from Friend or Relative without Asking	0.3857		0.5532		0.2711		0.4049	
Bought from Drug Dealer or Other Stranger	0.4437		0.9822		0.2933		0.4781	
Bought on the Internet	0.2554		0.1803		0.3746		0.1111	
Some Other Way	0.1354		0.8077		0.8279		0.1924	

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.48C Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Pain Relievers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	137	156	16	15	44	42	130	149
From More Than One Doctor	22	27	6	4	11	8	17	25
Wrote Fake Prescription	11	1	1	0	2	1	10	0
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	5	5	4	1	4	4	2	2
From Friend or Relative for Free	34	27	6	6	13	12	31	23
Bought from Friend or Relative	20	24	5	5	13	9	14	22
Took from Friend or Relative without Asking	13	12	7	3	8	9	8	7
Bought from Drug Dealer or Other Stranger	15	15	5	3	8	7	12	13
Bought on the Internet	5	9	0	1	1	4	5	7
Some Other Way	15	11	5	4	5	4	14	9

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.48D Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Pain Relievers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	0.95	0.99	2.25	2.48	1.38	1.38	1.29	1.35
From More Than One Doctor	0.41	0.52	1.22	1.25	0.67	0.57	0.55	0.73
Wrote Fake Prescription	0.20	0.02	0.12	0.10	0.12	0.05	0.33	0.01
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.10	0.09	0.74	0.25	0.22	0.31	0.05	0.06
From Friend or Relative for Free	0.65	0.51	1.23	1.67	0.80	0.81	0.96	0.67
Bought from Friend or Relative	0.38	0.45	1.07	1.29	0.79	0.63	0.43	0.64
Took from Friend or Relative without Asking	0.25	0.23	1.33	0.90	0.47	0.63	0.27	0.21
Bought from Drug Dealer or Other Stranger	0.30	0.28	1.00	0.98	0.51	0.46	0.37	0.37
Bought on the Internet	0.09	0.17	0.05	0.43	0.04	0.29	0.15	0.22
Some Other Way	0.29	0.21	0.95	1.10	0.30	0.29	0.43	0.27

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.48P Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Source Where Friend or Relative Obtained Pain Relievers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	0.0853		0.2947		0.0483		0.7935	
From More Than One Doctor	0.6984		0.7886		0.4822		0.3308	
Wrote Fake Prescription	0.1292		0.6759		0.3760		0.1833	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.4558		0.3750		0.9510		0.9385	
From Friend or Relative for Free	0.6160		0.5993		0.6976		0.8732	
Bought from Friend or Relative	0.7683		0.7274		0.0221		0.1783	
Took from Friend or Relative without Asking	0.0765		0.0046		0.9805		0.9014	
Bought from Drug Dealer or Other Stranger	0.2572		0.8232		0.2478		0.8135	
Bought on the Internet	0.5478		0.2511		0.2631		0.9046	
Some Other Way	0.1339		0.6812		0.9646		0.2205	

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.49C Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	75	76	10	11	20	20	72	72
From More Than One Doctor	33	36	5	6	8	9	31	34
Wrote Fake Prescription	12	9	2	3	2	1	12	9
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	15	19	2	2	5	2	14	19
From Friend or Relative for Free	103	111	15	12	36	34	95	106
Bought from Friend or Relative	66	70	8	6	24	23	59	66
Took from Friend or Relative without Asking	46	41	8	6	16	13	42	39
Bought from Drug Dealer or Other Stranger	39	39	7	4	20	19	32	34
Bought on the Internet	15	10	1	1	5	5	14	9
Some Other Way	33	25	5	5	10	12	31	22

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.49D Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	1.47	1.54	1.87	2.31	1.43	1.53	2.35	2.24
From More Than One Doctor	0.75	0.83	0.98	1.42	0.63	0.76	1.16	1.23
Wrote Fake Prescription	0.28	0.21	0.51	0.67	0.14	0.09	0.45	0.32
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.35	0.45	0.39	0.59	0.36	0.16	0.55	0.68
From Friend or Relative for Free	1.50	1.73	2.25	2.40	1.59	1.83	2.35	2.58
Bought from Friend or Relative	1.29	1.46	1.53	1.56	1.57	1.68	1.98	2.15
Took from Friend or Relative without Asking	1.00	0.91	1.56	1.60	1.13	1.13	1.53	1.34
Bought from Drug Dealer or Other Stranger	0.86	0.90	1.33	1.13	1.29	1.48	1.21	1.21
Bought on the Internet	0.34	0.24	0.12	0.28	0.43	0.47	0.53	0.31
Some Other Way	0.76	0.60	1.08	1.39	0.79	1.03	1.19	0.80

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.49P Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	0.6789		0.0988		0.5851		0.8599	
From More Than One Doctor	0.3879		0.4523		0.5420		0.5424	
Wrote Fake Prescription	0.6706		0.5426		0.6115		0.6023	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.7580		0.6594		0.0395		0.5135	
From Friend or Relative for Free	0.5151		0.1423		0.3406		0.8364	
Bought from Friend or Relative	0.4026		0.0174		0.1860		0.9337	
Took from Friend or Relative without Asking	0.0264		0.1254		0.0102		0.3081	
Bought from Drug Dealer or Other Stranger	0.9502		0.1313		0.7876		0.4940	
Bought on the Internet	0.3438		0.4413		0.6937		0.3757	
Some Other Way	0.0896		0.5502		0.7817		0.0575	

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.50C Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Pain Relievers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	90	99	9	10	28	29	85	94
From More Than One Doctor	22	23	6	3	11	12	18	20
Wrote Fake Prescription	15	9	0	1	2	2	15	9
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	7	10	1	2	4	5	6	*
From Friend or Relative for Free	32	28	4	4	13	16	29	23
Bought from Friend or Relative	27	29	4	3	14	15	23	25
Took from Friend or Relative without Asking	16	20	5	3	8	12	13	15
Bought from Drug Dealer or Other Stranger	25	20	5	3	14	10	20	17
Bought on the Internet	3	12	0	2	2	5	1	11
Some Other Way	16	11	4	2	6	6	14	9

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.50D Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Pain Relievers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	1.60	1.49	3.41	3.65	2.10	1.98	2.33	2.01
From More Than One Doctor	0.92	0.98	2.56	2.25	1.35	1.75	1.29	1.31
Wrote Fake Prescription	0.64	0.41	0.05	0.85	0.28	0.32	1.07	0.61
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.31	0.44	0.31	1.02	0.59	0.84	0.41	*
From Friend or Relative for Free	1.29	1.20	2.11	2.43	1.59	2.22	1.99	1.55
Bought from Friend or Relative	1.14	1.21	2.05	2.23	1.68	2.11	1.62	1.62
Took from Friend or Relative without Asking	0.67	0.86	2.15	1.75	1.08	1.78	0.93	1.04
Bought from Drug Dealer or Other Stranger	1.05	0.88	2.37	1.73	1.68	1.44	1.43	1.16
Bought on the Internet	0.11	0.55	0.09	1.10	0.32	0.84	0.07	0.76
Some Other Way	0.68	0.49	1.97	1.43	0.80	0.91	1.02	0.61

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.50P Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Source Where Friend or Relative Obtained Pain Relievers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	0.0176		0.0461		0.0486		0.4224	
From More Than One Doctor	0.9885		0.8917		0.5914		0.9809	
Wrote Fake Prescription	0.2980		0.1009		0.8887		0.2042	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.8671		0.2512		0.9245		*	
From Friend or Relative for Free	0.2225		0.4379		0.7636		0.2299	
Bought from Friend or Relative	0.7486		0.5516		0.4476		0.6831	
Took from Friend or Relative without Asking	0.8713		0.0584		0.5448		0.5797	
Bought from Drug Dealer or Other Stranger	0.1384		0.3290		0.2819		0.5098	
Bought on the Internet	0.0772		0.2121		0.2984		0.1873	
Some Other Way	0.1483		0.3071		0.8654		0.2620	

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.51C Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	63	68	5	5	12	12	62	66
From More Than One Doctor	15	15	2	1	5	7	14	13
Wrote Fake Prescription	1	1	1	1	1	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	6	12	1	0	5	1	4	11
From Friend or Relative for Free	115	137	14	13	36	34	107	133
Bought from Friend or Relative	48	45	8	7	20	21	43	40
Took from Friend or Relative without Asking	19	30	5	5	9	12	16	26
Bought from Drug Dealer or Other Stranger	28	26	6	4	14	16	22	20
Bought on the Internet	14	3	1	1	2	3	*	*
Some Other Way	19	31	4	4	8	13	18	28

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.51D Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	1.19	1.17	1.31	1.56	0.78	0.75	1.88	1.73
From More Than One Doctor	0.31	0.28	0.63	0.42	0.34	0.46	0.47	0.38
Wrote Fake Prescription	0.03	0.02	0.27	0.26	0.06	0.05	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.13	0.22	0.18	0.08	0.34	0.08	0.12	0.33
From Friend or Relative for Free	1.53	1.41	2.63	2.77	1.49	1.56	2.30	2.06
Bought from Friend or Relative	0.92	0.86	1.90	2.17	1.14	1.20	1.35	1.14
Took from Friend or Relative without Asking	0.39	0.54	1.30	1.57	0.58	0.80	0.53	0.74
Bought from Drug Dealer or Other Stranger	0.56	0.50	1.61	1.15	0.85	0.99	0.73	0.59
Bought on the Internet	0.29	0.05	0.14	0.40	0.12	0.17	*	*
Some Other Way	0.39	0.57	1.01	1.11	0.49	0.80	0.58	0.79

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.51P Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011**

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	0.7895		0.8432		0.7801		0.6385	
From More Than One Doctor	0.6045		0.6440		0.3991		0.3966	
Wrote Fake Prescription	0.2099		0.3333		0.7300		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.9163		0.1368		0.1756		0.3917	
From Friend or Relative for Free	0.2878		0.7221		0.2522		0.2542	
Bought from Friend or Relative	0.2187		0.4093		0.7455		0.1941	
Took from Friend or Relative without Asking	0.3868		0.7863		0.6768		0.2959	
Bought from Drug Dealer or Other Stranger	0.1167		0.0634		0.9350		0.3159	
Bought on the Internet	0.3757		0.4013		0.8177		*	
Some Other Way	0.2034		0.7489		0.0572		0.5318	

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.52C Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Tranquilizers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	103	120	9	9	28	25	98	117
From More Than One Doctor	14	20	3	2	7	6	12	19
Wrote Fake Prescription	1	0	*	0	1	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	3	*	1	1	2	4	2
From Friend or Relative for Free	27	22	5	5	12	10	24	19
Bought from Friend or Relative	15	17	4	5	7	9	12	14
Took from Friend or Relative without Asking	6	7	2	3	5	4	3	5
Bought from Drug Dealer or Other Stranger	13	11	4	2	7	6	10	9
Bought on the Internet	4	7	*	*	1	*	4	6
Some Other Way	8	10	3	*	2	6	7	8

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.52D Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Tranquilizers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	1.31	1.23	4.10	4.38	1.99	2.10	1.74	1.48
From More Than One Doctor	0.51	0.66	1.79	1.33	0.92	0.82	0.66	0.88
Wrote Fake Prescription	0.06	0.01	*	0.26	0.20	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.15	0.09	*	0.44	0.08	0.22	0.23	0.11
From Friend or Relative for Free	0.98	0.73	2.73	3.29	1.55	1.32	1.31	0.87
Bought from Friend or Relative	0.53	0.57	2.54	3.78	0.88	1.23	0.69	0.62
Took from Friend or Relative without Asking	0.23	0.24	1.50	1.93	0.62	0.58	0.16	0.25
Bought from Drug Dealer or Other Stranger	0.47	0.39	2.31	1.22	0.92	0.86	0.55	0.44
Bought on the Internet	0.15	0.24	*	*	0.11	*	0.21	0.28
Some Other Way	0.31	0.34	2.11	*	0.25	0.78	0.40	0.40

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.52P Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Source Where Friend or Relative Obtained Tranquilizers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	0.4988		0.6358		0.4738		0.7836	
From More Than One Doctor	0.9719		0.8696		0.2101		0.4984	
Wrote Fake Prescription	0.3060		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.9714		*		0.2583		0.5502	
From Friend or Relative for Free	0.0933		0.6722		0.5736		0.1660	
Bought from Friend or Relative	0.4057		0.6739		0.0681		0.8491	
Took from Friend or Relative without Asking	0.6775		0.4248		0.9537		0.4363	
Bought from Drug Dealer or Other Stranger	0.8236		0.1294		0.8321		0.6861	
Bought on the Internet	0.8004		*		*		0.9875	
Some Other Way	0.6122		*		0.1172		0.6660	

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.53C Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	36	52	3	*	9	7	36	51
From More Than One Doctor	5	4	2	1	4	3	2	2
Wrote Fake Prescription	1	1	1	*	1	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	13	4	1	*	3	*	*	4
From Friend or Relative for Free	75	79	6	7	20	17	72	76
Bought from Friend or Relative	41	32	5	4	12	14	38	28
Took from Friend or Relative without Asking	18	18	3	3	8	8	16	15
Bought from Drug Dealer or Other Stranger	16	20	5	3	10	9	12	18
Bought on the Internet	7	6	0	*	2	*	7	*
Some Other Way	11	13	2	3	3	6	11	11

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.53D Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	1.85	2.47	2.65	*	1.63	1.50	2.73	3.45
From More Than One Doctor	0.29	0.24	1.81	1.36	0.84	0.76	0.18	0.18
Wrote Fake Prescription	0.08	0.04	0.68	*	0.24	0.03	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.72	0.21	0.55	*	0.54	*	*	0.30
From Friend or Relative for Free	2.43	2.62	4.16	5.26	2.52	2.81	3.46	3.61
Bought from Friend or Relative	1.97	1.60	3.85	4.34	2.07	2.56	2.86	2.10
Took from Friend or Relative without Asking	0.99	0.95	2.72	2.79	1.47	1.67	1.34	1.20
Bought from Drug Dealer or Other Stranger	0.92	1.11	3.85	2.72	1.76	1.88	1.01	1.42
Bought on the Internet	0.39	0.31	0.33	*	0.49	*	0.56	*
Some Other Way	0.59	0.72	2.02	2.59	0.57	1.40	0.89	0.87

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.53P Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	0.7299		*		0.2429		0.6974	
From More Than One Doctor	0.9340		0.8656		0.4898		0.6352	
Wrote Fake Prescription	0.1199		*		0.1123		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.1949		*		*		*	
From Friend or Relative for Free	0.7884		0.0969		0.9582		0.9434	
Bought from Friend or Relative	0.1355		0.2875		0.6911		0.1549	
Took from Friend or Relative without Asking	0.9620		0.5032		0.5086		0.6938	
Bought from Drug Dealer or Other Stranger	0.5423		0.3542		0.1253		0.4555	
Bought on the Internet	0.5549		*		*		*	
Some Other Way	0.6638		0.9083		0.1415		0.8738	

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.54C Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Tranquilizers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	65	69	*	*	16	13	64	67
From More Than One Doctor	15	16	*	*	5	3	14	16
Wrote Fake Prescription	1	*	*	*	1	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	*	*	*	2	*	*	*
From Friend or Relative for Free	25	14	*	*	7	7	24	12
Bought from Friend or Relative	13	13	*	*	4	8	12	9
Took from Friend or Relative without Asking	10	6	*	*	3	1	*	6
Bought from Drug Dealer or Other Stranger	14	11	*	*	5	5	13	10
Bought on the Internet	4	9	*	*	*	*	*	8
Some Other Way	6	9	*	*	2	*	6	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.54D Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Tranquilizers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	2.55	2.48	*	*	3.60	3.80	3.36	2.93
From More Than One Doctor	1.51	1.56	*	*	2.25	1.23	1.96	2.08
Wrote Fake Prescription	0.07	*	*	*	0.27	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.43	*	*	*	0.79	*	*	*
From Friend or Relative for Free	2.42	1.42	*	*	2.82	2.93	3.23	1.68
Bought from Friend or Relative	1.30	1.34	*	*	1.92	3.48	1.63	1.24
Took from Friend or Relative without Asking	1.00	0.61	*	*	1.38	0.65	*	0.81
Bought from Drug Dealer or Other Stranger	1.43	1.10	*	*	1.99	2.08	1.82	1.33
Bought on the Internet	0.40	0.91	*	*	*	*	*	1.11
Some Other Way	0.67	0.94	*	*	0.71	*	0.88	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.54P Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Source Where Friend or Relative Obtained Tranquilizers	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	0.6405		*		0.2601		0.3882	
From More Than One Doctor	0.5083		*		0.0287		0.8857	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	0.0364		*		0.4610		0.0522	
Bought from Friend or Relative	0.5158		*		0.0520		0.5804	
Took from Friend or Relative without Asking	0.5142		*		0.1002		*	
Bought from Drug Dealer or Other Stranger	0.8596		*		0.9689		0.8277	
Bought on the Internet	0.3591		*		*		*	
Some Other Way	0.5097		*		*		*	

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.55C Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source for Most Recent Nonmedical Use among Past Year Stimulant Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	28	35	3	5	9	9	26	34
From More Than One Doctor	4	9	2	2	3	8	*	5
Wrote Fake Prescription	2	1	2	0	0	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	8	3	1	1	2	1	*	3
From Friend or Relative for Free	62	77	9	9	28	30	54	72
Bought from Friend or Relative	31	41	6	7	19	25	24	31
Took from Friend or Relative without Asking	15	18	4	4	6	6	13	18
Bought from Drug Dealer or Other Stranger	30	17	4	4	9	11	29	12
Bought on the Internet	7	9	1	*	3	4	7	8
Some Other Way	17	13	2	2	5	6	16	11

*Low precision; no estimate reported.

NOTE: Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.55D Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source for Most Recent Nonmedical Use among Past Year Stimulant Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	1.25	1.30	1.64	1.99	0.90	0.76	2.74	2.63
From More Than One Doctor	0.19	0.37	1.13	0.68	0.32	0.71	*	0.41
Wrote Fake Prescription	0.09	0.05	0.89	0.01	0.01	0.11	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.40	0.14	0.54	0.30	0.19	0.13	*	0.26
From Friend or Relative for Free	1.96	1.94	3.44	3.13	1.73	1.92	4.18	3.68
Bought from Friend or Relative	1.43	1.47	2.80	2.72	1.62	1.84	2.51	2.41
Took from Friend or Relative without Asking	0.71	0.73	1.69	1.64	0.59	0.52	1.52	1.45
Bought from Drug Dealer or Other Stranger	1.39	0.66	1.89	1.56	0.87	0.96	3.08	1.02
Bought on the Internet	0.36	0.36	0.35	*	0.30	0.34	0.75	0.69
Some Other Way	0.81	0.53	0.97	0.81	0.55	0.58	1.77	0.95

*Low precision; no estimate reported.

NOTE: Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.55P Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011**

Source for Most Recent Nonmedical Use among Past Year Stimulant Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	0.6158		0.3137		0.3506		0.6446	
From More Than One Doctor	0.1046		0.5629		0.1411		*	
Wrote Fake Prescription	0.4057		0.0804		0.1510		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3190		0.2557		0.4690		*	
From Friend or Relative for Free	0.8690		0.6464		0.2246		0.5728	
Bought from Friend or Relative	0.3923		0.8369		0.4010		0.4191	
Took from Friend or Relative without Asking	0.7186		0.5931		0.3342		0.9311	
Bought from Drug Dealer or Other Stranger	0.2142		0.6117		0.3722		0.1358	
Bought on the Internet	0.8756		*		0.2497		0.7541	
Some Other Way	0.2615		0.7444		0.9126		0.1853	

*Low precision; no estimate reported.

NOTE: Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.56C Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Stimulants	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	56	63	7	7	23	25	50	59
From More Than One Doctor	6	9	2	2	6	6	2	4
Wrote Fake Prescription	1	0	*	*	0	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1	1	*	0	1	1	*	*
From Friend or Relative for Free	12	14	3	3	7	6	10	12
Bought from Friend or Relative	9	14	2	*	7	7	5	12
Took from Friend or Relative without Asking	4	2	*	1	2	2	*	*
Bought from Drug Dealer or Other Stranger	10	10	2	1	5	6	*	8
Bought on the Internet	3	2	0	*	*	1	3	*
Some Other Way	1	2	1	1	1	1	*	2

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.56D Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Stimulants	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	2.00	2.02	5.04	4.58	2.43	2.23	3.13	3.22
From More Than One Doctor	0.68	0.73	2.80	2.61	1.18	1.21	0.52	0.76
Wrote Fake Prescription	0.07	0.03	*	*	0.10	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.14	0.13	*	0.05	0.29	0.30	*	*
From Friend or Relative for Free	1.23	1.19	3.08	2.85	1.46	1.21	2.19	2.06
Bought from Friend or Relative	0.95	1.20	2.27	*	1.51	1.37	1.26	2.08
Took from Friend or Relative without Asking	0.38	0.17	*	1.08	0.53	0.34	*	*
Bought from Drug Dealer or Other Stranger	0.97	0.82	2.05	1.45	1.00	1.10	*	1.33
Bought on the Internet	0.29	0.14	0.55	*	*	0.23	0.65	*
Some Other Way	0.10	0.20	0.98	1.08	0.13	0.17	*	0.33

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.56P Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Source Where Friend or Relative Obtained Stimulants	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	0.5725		0.1865		0.2371		0.4311	
From More Than One Doctor	0.9690		0.7267		0.7460		0.8073	
Wrote Fake Prescription	0.3201		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.6921		*		0.8773		*	
From Friend or Relative for Free	0.6988		0.8918		0.0586		0.4422	
Bought from Friend or Relative	0.7043		*		0.3516		0.5717	
Took from Friend or Relative without Asking	0.0580		*		0.2868		*	
Bought from Drug Dealer or Other Stranger	0.6844		0.5343		0.3350		*	
Bought on the Internet	0.5170		*		*		*	
Some Other Way	0.2268		0.7826		0.1886		*	

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, (c) respondents with unknown or invalid responses to the corresponding other-specify questions, and (d) respondents reporting methamphetamine use as their only use of a stimulant in the past year.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.57C Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source for Past Month Nonmedical Use among Past Month Stimulant Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	25	26	3	3	8	5	*	25
From More Than One Doctor	3	10	1	1	3	4	*	*
Wrote Fake Prescription	2	*	*	*	0	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	*	0	*	2	1	*	*
From Friend or Relative for Free	31	41	4	6	17	17	26	38
Bought from Friend or Relative	23	35	4	4	13	13	19	*
Took from Friend or Relative without Asking	11	11	2	2	6	4	9	10
Bought from Drug Dealer or Other Stranger	16	15	4	3	8	6	13	14
Bought on the Internet	4	10	0	*	3	3	3	*
Some Other Way	9	10	2	1	2	5	9	8

*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.57D Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source for Past Month Nonmedical Use among Past Month Stimulant Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	2.98	2.68	3.71	3.82	2.26	1.41	*	5.04
From More Than One Doctor	0.44	1.16	1.99	1.77	0.91	1.26	*	*
Wrote Fake Prescription	0.24	*	*	*	0.03	0.26	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.23	*	0.46	*	0.53	0.29	*	*
From Friend or Relative for Free	3.16	3.23	5.30	5.47	3.14	3.11	5.81	5.67
Bought from Friend or Relative	2.68	3.37	4.82	5.31	3.04	3.02	4.65	*
Took from Friend or Relative without Asking	1.43	1.32	2.98	3.08	1.89	1.17	2.57	2.39
Bought from Drug Dealer or Other Stranger	1.96	1.74	4.60	4.44	2.16	1.65	3.55	3.07
Bought on the Internet	0.57	1.19	0.63	*	0.82	0.78	0.95	*
Some Other Way	1.25	1.14	2.66	1.41	0.57	1.42	2.53	1.88

*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.57P Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Source for Past Month Nonmedical Use among Past Month Stimulant Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	0.2783		0.2090		0.0281		*	
From More Than One Doctor	0.3435		0.8749		0.8699		*	
Wrote Fake Prescription	*		*		0.3048		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		0.4621		*	
From Friend or Relative for Free	0.1509		0.0498		0.7914		0.2297	
Bought from Friend or Relative	0.3880		0.5757		0.5035		*	
Took from Friend or Relative without Asking	0.3257		0.6479		0.0947		0.9626	
Bought from Drug Dealer or Other Stranger	0.1527		0.9602		0.2481		0.3190	
Bought on the Internet	0.3148		*		0.7074		*	
Some Other Way	0.9966		0.3660		0.0473		0.4932	

*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.58C Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Stimulants	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	28	34	*	*	14	15	*	*
From More Than One Doctor	5	9	*	*	5	3	*	*
Wrote Fake Prescription	0	*	*	*	*	0	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1	*	*	*	*	*	*	*
From Friend or Relative for Free	8	14	*	*	5	4	*	*
Bought from Friend or Relative	6	16	*	*	4	2	*	*
Took from Friend or Relative without Asking	4	*	*	*	3	1	*	*
Bought from Drug Dealer or Other Stranger	7	14	*	*	2	2	*	*
Bought on the Internet	3	*	*	*	*	*	*	*
Some Other Way	1	2	*	*	0	2	*	*

*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.58D Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Stimulants	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	3.26	3.85	*	*	4.06	2.74	*	*
From More Than One Doctor	1.69	2.24	*	*	3.16	1.73	*	*
Wrote Fake Prescription	0.03	*	*	*	*	0.14	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.27	*	*	*	*	*	*	*
From Friend or Relative for Free	2.43	3.11	*	*	3.11	2.02	*	*
Bought from Friend or Relative	2.01	3.64	*	*	2.60	1.39	*	*
Took from Friend or Relative without Asking	1.16	*	*	*	2.15	0.43	*	*
Bought from Drug Dealer or Other Stranger	2.11	3.18	*	*	1.58	1.30	*	*
Bought on the Internet	1.09	*	*	*	*	*	*	*
Some Other Way	0.46	0.52	*	*	0.09	1.06	*	*

*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.58P Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Source Where Friend or Relative Obtained Stimulants	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	0.4973		*		0.2878		*	
From More Than One Doctor	0.7354		*		0.1331		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	0.4550		*		0.4183		*	
Bought from Friend or Relative	0.1755		*		0.1554		*	
Took from Friend or Relative without Asking	*		*		0.0634		*	
Bought from Drug Dealer or Other Stranger	0.2443		*		0.8539		*	
Bought on the Internet	*		*		*		*	
Some Other Way	0.7067		*		0.1372		*	

*Low precision; no estimate reported.

NOTE: Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.59C Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From Friend or Relative for Free	36	43	*	*	12	16	33	40
Bought from Friend or Relative	50	35	*	3	8	12	*	33
Took from Friend or Relative without Asking	9	18	1	1	*	2	7	*
Bought from Drug Dealer or Other Stranger	35	37	*	3	8	13	*	33
Bought on the Internet	5	1	*	*	2	0	*	1
Some Other Way	8	27	*	1	3	4	6	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.59D Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From Friend or Relative for Free	3.69	3.39	*	*	4.13	4.34	5.15	4.74
Bought from Friend or Relative	4.41	2.94	*	4.41	3.40	3.24	*	4.29
Took from Friend or Relative without Asking	1.05	1.72	1.30	1.97	*	0.76	1.21	*
Bought from Drug Dealer or Other Stranger	3.61	3.03	*	5.12	3.29	3.33	*	4.30
Bought on the Internet	0.55	0.08	*	*	0.78	0.13	*	0.09
Some Other Way	0.92	2.51	*	1.32	1.43	1.26	1.16	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.59P Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From Friend or Relative for Free	0.8772		*		0.6635		0.9325	
Bought from Friend or Relative	0.2018		*		0.7549		*	
Took from Friend or Relative without Asking	0.9401		0.4138		*		*	
Bought from Drug Dealer or Other Stranger	0.2870		*		0.1296		*	
Bought on the Internet	0.1313		*		0.0956		*	
Some Other Way	0.1756		*		0.5050		*	

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.60C Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Methamphetamine	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From Friend or Relative for Free	20	21	*	*	8	8	*	*
Bought from Friend or Relative	18	22	*	*	5	8	*	*
Took from Friend or Relative without Asking	*	3	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	10	14	*	*	4	7	*	*
Bought on the Internet	1	*	*	*	*	*	*	*
Some Other Way	5	*	*	*	3	4	*	*

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.60D Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Methamphetamine	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From Friend or Relative for Free	5.38	4.88	*	*	5.72	5.28	*	*
Bought from Friend or Relative	5.26	5.11	*	*	5.01	5.41	*	*
Took from Friend or Relative without Asking	*	0.81	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	3.46	3.83	*	*	4.43	5.13	*	*
Bought on the Internet	0.49	*	*	*	*	*	*	*
Some Other Way	1.88	*	*	*	2.81	2.88	*	*

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.60P Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year, by Age Group: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Source Where Friend or Relative Obtained Methamphetamine	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From Friend or Relative for Free	0.4226		*		0.5584		*	
Bought from Friend or Relative	0.6379		*		0.6697		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	0.6105		*		0.2381		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		0.5265		*	

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.61C Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From Friend or Relative for Free	26	33	*	*	*	*	*	*
Bought from Friend or Relative	*	26	*	*	*	*	*	*
Took from Friend or Relative without Asking	4	*	*	*	*	2	*	*
Bought from Drug Dealer or Other Stranger	19	21	*	*	*	*	*	*
Bought on the Internet	3	0	*	*	*	0	*	*
Some Other Way	4	*	*	*	*	2	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.61D Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From Friend or Relative for Free	5.60	5.77	*	*	*	*	*	*
Bought from Friend or Relative	*	5.16	*	*	*	*	*	*
Took from Friend or Relative without Asking	1.18	*	*	*	*	1.65	*	*
Bought from Drug Dealer or Other Stranger	5.30	4.38	*	*	*	*	*	*
Bought on the Internet	1.02	0.10	*	*	*	0.15	*	*
Some Other Way	1.24	*	*	*	*	2.03	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.61P Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From Friend or Relative for Free	0.3512		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	0.3928		*		*		*	
Bought on the Internet	0.0655		*		*		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.62C Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Methamphetamine	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	3	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.62D Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Methamphetamine	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	1.66	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.62P Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Source Where Friend or Relative Obtained Methamphetamine	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.63C Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source for Most Recent Nonmedical Use among Past Year Sedative Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	26	20	3	*	5	4	*	19
From More Than One Doctor	*	5	*	*	1	*	*	4
Wrote Fake Prescription	1	*	*	*	*	1	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	*	2	*	*	2	0	*
From Friend or Relative for Free	49	37	7	3	10	10	*	*
Bought from Friend or Relative	13	16	3	2	4	5	12	15
Took from Friend or Relative without Asking	9	7	2	3	4	2	*	*
Bought from Drug Dealer or Other Stranger	9	4	1	1	2	2	8	4
Bought on the Internet	0	1	0	*	*	0	*	*
Some Other Way	8	11	*	*	1	*	*	10

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.63D Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source for Most Recent Nonmedical Use among Past Year Sedative Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	3.67	3.32	4.04	*	3.61	3.50	*	4.58
From More Than One Doctor	*	0.88	*	*	0.85	*	*	1.02
Wrote Fake Prescription	0.15	*	*	*	*	0.43	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.31	*	2.72	*	*	1.40	0.06	*
From Friend or Relative for Free	4.45	4.59	5.58	5.21	4.82	5.07	*	*
Bought from Friend or Relative	1.93	2.74	3.72	3.76	3.17	3.62	2.54	3.73
Took from Friend or Relative without Asking	1.38	1.24	2.17	4.44	2.84	1.90	*	*
Bought from Drug Dealer or Other Stranger	1.29	0.81	1.52	1.78	1.48	1.63	1.81	1.01
Bought on the Internet	0.01	0.16	0.05	*	*	0.14	*	*
Some Other Way	1.16	1.85	*	*	1.07	*	*	2.60

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

**Table 6.63P Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011**

Source for Most Recent Nonmedical Use among Past Year Sedative Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	0.6950		*		0.7230		*	
From More Than One Doctor	*		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	0.2659		0.0057		0.9346		*	
Bought from Friend or Relative	0.3228		0.7801		0.8107		0.2301	
Took from Friend or Relative without Asking	0.9688		0.0374		0.3359		*	
Bought from Drug Dealer or Other Stranger	0.3790		0.9134		0.6123		0.2711	
Bought on the Internet	0.2596		*		*		*	
Some Other Way	0.4455		*		*		*	

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment of this drug for their past month nonmedical use and (b) all other past month and past year users who reported their most recent source of obtainment of this drug for their most recent nonmedical use.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.64C Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Sedatives	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	48	35	*	*	*	*	*	*
From More Than One Doctor	3	3	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	2	*	*	*	*	*	*
From Friend or Relative for Free	3	5	*	*	*	*	*	*
Bought from Friend or Relative	2	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	3	2	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	4	4	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	2	0	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.64D Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Sedatives	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	2.41	3.17	*	*	*	*	*	*
From More Than One Doctor	0.84	1.23	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.52	0.70	*	*	*	*	*	*
From Friend or Relative for Free	0.76	1.68	*	*	*	*	*	*
Bought from Friend or Relative	0.65	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	0.88	0.65	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	1.17	1.29	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	0.49	0.05	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.64P Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Source Where Friend or Relative Obtained Sedatives	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	0.4348		*		*		*	
From More Than One Doctor	0.4234		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.7528		*		*		*	
From Friend or Relative for Free	0.1905		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	0.5041		*		*		*	
Bought from Drug Dealer or Other Stranger	0.6469		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	0.1804		*		*		*	

*Low precision; no estimate reported.

NOTE: Excluded from the analysis are (a) past month users who reported multiple sources for where their friend or relative obtained this drug, (b) respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use, and (c) respondents with unknown or invalid responses to the corresponding other-specify questions.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free, then reported a single valid source for where their friend or relative obtained this drug; and (b) all other past month and past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free, then reported a valid source for where their friend or relative obtained this drug.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.65C Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source for Past Month Nonmedical Use among Past Month Sedative Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	0	4	*	*	*	*	*	*
Wrote Fake Prescription	1	0	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	3	3	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	5	*	*	*	*	*	*	*
Bought on the Internet	0	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.65D Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source for Past Month Nonmedical Use among Past Month Sedative Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	0.04	1.77	*	*	*	*	*	*
Wrote Fake Prescription	0.34	0.21	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.47	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	1.15	1.41	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	1.96	*	*	*	*	*	*	*
Bought on the Internet	0.04	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.65P Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Source for Past Month Nonmedical Use among Past Month Sedative Users	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	*		*		*		*	
From More Than One Doctor	0.1019		*		*		*	
Wrote Fake Prescription	0.8127		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	0.8955		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.66C Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Sedatives	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.66D Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Source Where Friend or Relative Obtained Sedatives	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.66P Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Source Where Friend or Relative Obtained Sedatives	Total (2010-2011)	Total (2012-2013)	Aged 12-17 (2010-2011)	Aged 12-17 (2012-2013)	Aged 18-25 (2010-2011)	Aged 18-25 (2012-2013)	Aged 26+ (2010-2011)	Aged 26+ (2012-2013)
From One Doctor	*		*		*		*	
From More Than One Doctor	*		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.