

Table 6.71C Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
ILLCIT DRUGS²	147	160	16	18	146	157
Marijuana and Hashish	120	132	15	18	118	130
Cocaine	34	41	1	3	34	41
Crack	18	16	1	*	18	16
Heroin	15	15	3	2	14	15
Hallucinogens	29	31	1	2	29	31
LSD	4	15	*	*	4	15
PCP	2	7	0	*	2	7
Ecstasy	22	26	1	1	22	26
Inhalants	18	16	1	3	18	15
Nonmedical Use of Psychotherapeutics ^{3,4}	85	91	6	8	84	91
Pain Relievers	65	73	6	4	64	73
OxyContin [®]	18	14	1	2	18	14
Tranquilizers	49	48	1	4	49	48
Stimulants ⁴	31	41	2	7	31	40
Methamphetamine ⁴	20	20	*	7	20	18
Sedatives	13	25	0	1	13	25
ILLCIT DRUGS OTHER THAN MARIJUANA²	93	100	7	9	93	99

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.71D Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
ILLCIT DRUGS²	0.22	0.23	0.67	0.79	0.22	0.24
Marijuana and Hashish	0.18	0.19	0.63	0.78	0.19	0.20
Cocaine	0.05	0.07	0.05	0.13	0.06	0.07
Crack	0.03	0.03	0.05	*	0.03	0.03
Heroin	0.02	0.02	0.15	0.10	0.02	0.02
Hallucinogens	0.05	0.05	0.04	0.07	0.05	0.05
LSD	0.01	0.02	*	*	0.01	0.02
PCP	0.00	0.01	0.01	*	0.00	0.01
Ecstasy	0.04	0.04	0.04	0.06	0.04	0.04
Inhalants	0.03	0.02	0.04	0.12	0.03	0.03
Nonmedical Use of Psychotherapeutics ^{3,4}	0.13	0.14	0.27	0.37	0.14	0.14
Pain Relievers	0.10	0.11	0.25	0.20	0.11	0.12
OxyContin [®]	0.03	0.02	0.05	0.09	0.03	0.02
Tranquilizers	0.08	0.07	0.06	0.17	0.08	0.08
Stimulants ⁴	0.05	0.06	0.08	0.29	0.05	0.07
Methamphetamine ⁴	0.03	0.03	*	0.29	0.03	0.03
Sedatives	0.02	0.04	0.01	0.05	0.02	0.04
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.14	0.15	0.29	0.39	0.15	0.16

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.71P Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
ILLCIT DRUGS²	0.0381		0.6430		0.0455	
Marijuana and Hashish	0.0072		0.4236		0.0097	
Cocaine	0.5342		0.3659		0.4877	
Crack	0.9940		*		0.9734	
Heroin	0.3910		0.8451		0.3760	
Hallucinogens	0.5767		0.3105		0.5338	
LSD	0.0972		*		0.0914	
PCP	0.4169		*		0.4053	
Ecstasy	0.4172		0.6231		0.4297	
Inhalants	0.0748		0.3499		0.0627	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.5504		0.6476		0.6022	
Pain Relievers	0.8036		0.4608		0.7801	
OxyContin [®]	0.0326		0.6965		0.0277	
Tranquilizers	0.4584		0.1380		0.3799	
Stimulants ⁴	0.0508		0.3572		0.0666	
Methamphetamine ⁴	0.9731		*		0.6814	
Sedatives	0.6141		0.4450		0.6208	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.9189		0.6472		0.8681	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.72C Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	147	160	16	18	146	157
AGE						
15-17	30	29	3	3	29	29
18-25	66	60	11	10	81	69
26-44	116	134	11	14	114	133
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	135	147	13	17	134	145
White	116	135	12	13	115	134
Black or African American	59	62	6	9	59	61
American Indian or Alaska Native	9	6	*	*	9	6
Native Hawaiian or Other Pacific Islander	8	7	*	*	8	7
Asian	23	22	*	*	23	22
Two or More Races	14	24	*	*	14	24
Hispanic or Latino	58	59	8	7	57	59
TRIMESTER²						
First	N/A	N/A	12	15	N/A	N/A
Second	N/A	N/A	9	8	N/A	N/A
Third	N/A	N/A	4	4	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.72D Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	0.22	0.23	0.67	0.79	0.22	0.24
AGE						
15-17	0.43	0.43	4.72	4.25	0.44	0.43
18-25	0.39	0.35	1.21	1.23	0.41	0.36
26-44	0.28	0.32	0.77	1.01	0.29	0.32
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.24	0.26	0.72	0.88	0.25	0.27
White	0.29	0.33	0.88	1.02	0.30	0.34
Black or African American	0.62	0.62	1.73	2.53	0.64	0.65
American Indian or Alaska Native	2.04	1.61	*	*	2.10	1.69
Native Hawaiian or Other Pacific Islander	2.89	2.75	*	*	2.96	2.92
Asian	0.66	0.55	*	*	0.68	0.57
Two or More Races	1.46	1.97	*	*	1.52	2.06
Hispanic or Latino	0.47	0.45	1.76	1.69	0.49	0.47
TRIMESTER²						
First	N/A	N/A	1.69	2.01	N/A	N/A
Second	N/A	N/A	1.12	1.02	N/A	N/A
Third	N/A	N/A	0.48	0.62	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.72P Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	0.0381		0.6430		0.0455	
AGE						
15-17	0.1011		0.3080		0.1245	
18-25	0.9111		0.8397		0.9842	
26-44	0.0052		0.4157		0.0069	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0083		0.6666		0.0094	
White	0.0539		0.6889		0.0625	
Black or African American	0.0505		0.6895		0.0628	
American Indian or Alaska Native	0.2756		*		0.3376	
Native Hawaiian or Other Pacific Islander	0.2119		*		0.1561	
Asian	0.7858		*		0.7489	
Two or More Races	0.0553		*		0.0412	
Hispanic or Latino	0.6112		0.8426		0.5807	
TRIMESTER²						
First	N/A		0.9456		N/A	
Second	N/A		0.7767		N/A	
Third	N/A		0.4006		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.73C Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	120	132	15	18	118	130
AGE						
15-17	26	26	3	3	26	26
18-25	62	57	10	10	74	62
26-44	93	113	10	14	91	112
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	112	123	13	16	111	121
White	97	111	11	13	96	111
Black or African American	52	53	6	9	52	51
American Indian or Alaska Native	5	5	*	*	5	6
Native Hawaiian or Other Pacific Islander	6	7	*	*	6	7
Asian	14	19	*	*	14	19
Two or More Races	13	22	*	*	13	22
Hispanic or Latino	42	45	7	7	41	45
TRIMESTER²						
First	N/A	N/A	12	15	N/A	N/A
Second	N/A	N/A	8	8	N/A	N/A
Third	N/A	N/A	3	4	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.73D Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	0.18	0.19	0.63	0.78	0.19	0.20
AGE						
15-17	0.39	0.39	4.53	4.24	0.39	0.39
18-25	0.36	0.33	1.11	1.16	0.38	0.34
26-44	0.23	0.27	0.75	1.01	0.24	0.28
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.21	0.22	0.69	0.87	0.21	0.23
White	0.25	0.27	0.85	1.01	0.26	0.28
Black or African American	0.54	0.56	1.71	2.50	0.57	0.57
American Indian or Alaska Native	1.23	1.47	*	*	1.27	1.54
Native Hawaiian or Other Pacific Islander	2.27	2.62	*	*	2.32	2.78
Asian	0.39	0.49	*	*	0.40	0.51
Two or More Races	1.33	1.80	*	*	1.39	1.88
Hispanic or Latino	0.35	0.35	1.50	1.64	0.37	0.36
TRIMESTER²						
First	N/A	N/A	1.67	1.99	N/A	N/A
Second	N/A	N/A	0.95	1.02	N/A	N/A
Third	N/A	N/A	0.47	0.55	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.73P Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	0.0072		0.4236		0.0097	
AGE						
15-17	0.9784		0.5174		0.9071	
18-25	0.9690		0.7914		0.9700	
26-44	0.0015		0.2748		0.0024	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0029		0.5793		0.0034	
White	0.0449		0.5633		0.0538	
Black or African American	0.0573		0.7008		0.0704	
American Indian or Alaska Native	0.9289		*		0.9453	
Native Hawaiian or Other Pacific Islander	0.1084		*		0.0752	
Asian	0.6358		*		0.6267	
Two or More Races	0.0361		*		0.0272	
Hispanic or Latino	0.8422		0.5000		0.9296	
TRIMESTER²						
First	N/A		0.9779		N/A	
Second	N/A		0.2541		N/A	
Third	N/A		0.6846		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.74C Illicit Drug, Tobacco Product, and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
ILLCIT DRUGS²	147	160	16	18	146	157
TOBACCO PRODUCTS³	274	266	34	31	268	262
Cigarettes	261	259	33	31	255	255
Smokeless Tobacco	33	29	2	6	33	29
Cigars	90	79	8	6	89	79
Pipe Tobacco	25	25	1	2	25	25
ALCOHOL	436	455	24	26	433	454
Binge Alcohol Use ⁴	258	263	10	10	257	263
Heavy Alcohol Use ⁴	116	118	4	4	116	118
ILLCIT DRUGS, TOBACCO PRODUCTS, OR ALCOHOL^{2,3}	466	486	41	41	458	483
ILLCIT DRUGS OR ALCOHOL²	445	460	28	30	440	459
TOBACCO PRODUCTS OR ALCOHOL³	462	482	40	39	455	479

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.74D Illicit Drug, Tobacco Product, and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
ILLCIT DRUGS²	0.22	0.23	0.67	0.79	0.22	0.24
TOBACCO PRODUCTS³	0.36	0.35	1.36	1.31	0.37	0.35
Cigarettes	0.35	0.34	1.35	1.29	0.36	0.35
Smokeless Tobacco	0.05	0.05	0.08	0.28	0.06	0.05
Cigars	0.14	0.12	0.34	0.25	0.14	0.13
Pipe Tobacco	0.04	0.04	0.04	0.09	0.04	0.04
ALCOHOL	0.40	0.42	1.02	1.13	0.41	0.43
Binge Alcohol Use ⁴	0.33	0.34	0.43	0.45	0.34	0.35
Heavy Alcohol Use ⁴	0.17	0.18	0.15	0.18	0.18	0.19
ILLCIT DRUGS, TOBACCO PRODUCTS, OR ALCOHOL^{2,3}	0.39	0.40	1.62	1.64	0.39	0.41
ILLCIT DRUGS OR ALCOHOL²	0.40	0.41	1.17	1.29	0.41	0.42
TOBACCO PRODUCTS OR ALCOHOL³	0.39	0.40	1.57	1.60	0.39	0.41

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.74P Illicit Drug, Tobacco Product, and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
ILLCIT DRUGS²	0.0381		0.6430		0.0455	
TOBACCO PRODUCTS³	0.0142		0.2460		0.0155	
Cigarettes	0.0057		0.2333		0.0061	
Smokeless Tobacco	0.2648		0.4189		0.2241	
Cigars	0.0486		0.3248		0.0468	
Pipe Tobacco	0.4142		0.2273		0.4455	
ALCOHOL	0.5784		0.9677		0.6607	
Binge Alcohol Use ⁴	0.6954		0.6352		0.7616	
Heavy Alcohol Use ⁴	0.8968		0.9148		0.8651	
ILLCIT DRUGS, TOBACCO PRODUCTS, OR ALCOHOL^{2,3}	0.9651		0.5532		0.9784	
ILLCIT DRUGS OR ALCOHOL²	0.4244		1.0000		0.4827	
TOBACCO PRODUCTS OR ALCOHOL³	0.8623		0.4749		0.8152	

*Low precision; no estimate reported.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006

³ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

⁴ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.75C Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	261	259	33	31	255	255
AGE						
15-17	26	24	*	*	26	23
18-25	74	72	15	15	90	83
26-44	230	238	29	26	226	236
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	242	239	33	29	236	235
White	217	209	30	24	211	207
Black or African American	85	93	10	15	84	91
American Indian or Alaska Native	17	26	*	*	17	25
Native Hawaiian or Other Pacific Islander	*	13	*	*	*	13
Asian	40	38	*	*	40	38
Two or More Races	30	33	*	*	30	33
Hispanic or Latino	95	91	4	9	95	90
TRIMESTER²						
First	N/A	N/A	24	20	N/A	N/A
Second	N/A	N/A	17	16	N/A	N/A
Third	N/A	N/A	16	17	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.75D Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	0.35	0.34	1.35	1.29	0.36	0.35
AGE						
15-17	0.39	0.37	*	*	0.39	0.36
18-25	0.44	0.42	1.53	1.68	0.46	0.43
26-44	0.50	0.51	2.01	1.78	0.51	0.52
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.39	0.39	1.66	1.50	0.40	0.40
White	0.46	0.46	2.04	1.80	0.47	0.47
Black or African American	0.82	0.90	2.96	3.64	0.84	0.93
American Indian or Alaska Native	3.19	4.22	*	*	3.29	4.35
Native Hawaiian or Other Pacific Islander	*	4.53	*	*	*	4.75
Asian	1.10	0.91	*	*	1.14	0.96
Two or More Races	2.22	2.53	*	*	2.27	2.60
Hispanic or Latino	0.75	0.68	0.84	2.09	0.78	0.70
TRIMESTER²						
First	N/A	N/A	3.03	2.56	N/A	N/A
Second	N/A	N/A	1.93	1.90	N/A	N/A
Third	N/A	N/A	2.03	2.21	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.75P Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	0.0057		0.2333		0.0061	
AGE						
15-17	0.0000		*		0.0000	
18-25	0.0000		0.5176		0.0000	
26-44	0.7513		0.3461		0.7633	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0746		0.0660		0.1039	
White	0.0709		0.1391		0.0918	
Black or African American	0.3908		0.6796		0.3807	
American Indian or Alaska Native	0.0287		*		0.0557	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.0479		*		0.0493	
Two or More Races	0.2855		*		0.3449	
Hispanic or Latino	0.0388		0.0715		0.0217	
TRIMESTER²						
First	N/A		0.1316		N/A	
Second	N/A		0.5098		N/A	
Third	N/A		0.9636		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.76C Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	436	455	24	26	433	454
AGE						
15-17	39	37	2	*	39	36
18-25	86	86	10	8	174	157
26-44	368	400	22	24	366	400
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	391	416	22	25	389	415
White	341	371	19	22	339	372
Black or African American	157	154	6	9	156	153
American Indian or Alaska Native	20	22	*	*	20	21
Native Hawaiian or Other Pacific Islander	*	18	*	*	*	18
Asian	100	97	*	*	99	93
Two or More Races	39	57	*	*	39	57
Hispanic or Latino	167	167	10	6	167	168
TRIMESTER²						
First	N/A	N/A	17	21	N/A	N/A
Second	N/A	N/A	15	9	N/A	N/A
Third	N/A	N/A	7	11	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.76D Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	0.40	0.42	1.02	1.13	0.41	0.43
AGE						
15-17	0.54	0.53	2.89	*	0.55	0.53
18-25	0.51	0.50	1.10	0.93	0.51	0.51
26-44	0.57	0.60	1.56	1.71	0.59	0.63
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.41	0.45	1.14	1.33	0.43	0.46
White	0.47	0.50	1.39	1.68	0.48	0.51
Black or African American	1.05	1.02	1.85	2.56	1.08	1.04
American Indian or Alaska Native	3.54	3.80	*	*	3.64	3.79
Native Hawaiian or Other Pacific Islander	*	5.37	*	*	*	5.55
Asian	2.00	1.93	*	*	2.07	1.94
Two or More Races	2.54	2.63	*	*	2.61	2.63
Hispanic or Latino	1.04	0.99	2.14	1.58	1.06	1.03
TRIMESTER²						
First	N/A	N/A	2.37	2.64	N/A	N/A
Second	N/A	N/A	1.75	1.17	N/A	N/A
Third	N/A	N/A	0.92	1.52	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.76P Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	0.5784		0.9677		0.6607	
AGE						
15-17	0.2077		*		0.1788	
18-25	0.6778		0.1836		0.5971	
26-44	0.3896		0.6615		0.4403	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.1950		0.8496		0.1790	
White	0.2252		0.9458		0.2343	
Black or African American	0.0065		0.3679		0.0045	
American Indian or Alaska Native	0.3221		*		0.2670	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.0203		*		0.0180	
Two or More Races	0.1003		*		0.1250	
Hispanic or Latino	0.6937		0.6754		0.5236	
TRIMESTER²						
First	N/A		0.8700		N/A	
Second	N/A		0.1587		N/A	
Third	N/A		0.4150		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.77C Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	258	263	10	10	257	263
AGE						
15-17	28	25	2	*	28	25
18-25	78	82	5	6	116	114
26-44	218	226	8	8	217	227
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	224	243	10	9	223	242
White	205	221	9	6	204	220
Black or African American	82	87	3	5	82	86
American Indian or Alaska Native	17	19	*	*	17	18
Native Hawaiian or Other Pacific Islander	13	12	*	*	13	12
Asian	51	46	*	*	51	46
Two or More Races	19	32	*	*	19	32
Hispanic or Latino	113	106	1	4	113	107
TRIMESTER²						
First	N/A	N/A	9	8	N/A	N/A
Second	N/A	N/A	4	5	N/A	N/A
Third	N/A	N/A	2	0	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.77D Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	0.33	0.34	0.43	0.45	0.34	0.35
AGE						
15-17	0.41	0.38	2.65	*	0.42	0.38
18-25	0.46	0.48	0.58	0.71	0.48	0.50
26-44	0.47	0.48	0.61	0.58	0.49	0.50
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.35	0.38	0.54	0.50	0.36	0.39
White	0.43	0.46	0.69	0.44	0.44	0.47
Black or African American	0.79	0.79	0.95	1.46	0.83	0.84
American Indian or Alaska Native	3.28	3.82	*	*	3.37	3.90
Native Hawaiian or Other Pacific Islander	4.46	4.34	*	*	4.57	4.56
Asian	1.32	1.07	*	*	1.36	1.12
Two or More Races	1.89	2.47	*	*	1.99	2.56
Hispanic or Latino	0.81	0.75	0.28	1.04	0.85	0.79
TRIMESTER²						
First	N/A	N/A	1.29	1.14	N/A	N/A
Second	N/A	N/A	0.53	0.68	N/A	N/A
Third	N/A	N/A	0.20	0.07	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.77P Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	0.6954		0.6352		0.7616	
AGE						
15-17	0.1217		*		0.1319	
18-25	0.3038		0.9579		0.2257	
26-44	0.3103		0.7171		0.3385	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.4397		0.2282		0.4308	
White	0.7280		0.0282		0.7212	
Black or African American	0.0402		0.8711		0.0397	
American Indian or Alaska Native	0.8436		*		0.8497	
Native Hawaiian or Other Pacific Islander	0.4290		*		0.3389	
Asian	0.2772		*		0.2738	
Two or More Races	0.2212		*		0.2895	
Hispanic or Latino	0.6947		0.0625		0.5509	
TRIMESTER²						
First	N/A		0.2841		N/A	
Second	N/A		0.5821		N/A	
Third	N/A		0.3754		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.78C Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	116	118	4	4	116	118
AGE						
15-17	12	11	1	*	12	11
18-25	48	47	3	1	56	50
26-44	96	105	*	4	96	105
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	107	110	3	4	107	110
White	97	101	3	3	97	101
Black or African American	36	39	1	3	36	39
American Indian or Alaska Native	12	6	*	*	12	6
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	15	22	*	*	15	22
Two or More Races	7	9	*	*	7	9
Hispanic or Latino	39	48	1	*	39	48
TRIMESTER²						
First	N/A	N/A	2	3	N/A	N/A
Second	N/A	N/A	3	3	N/A	N/A
Third	N/A	N/A	*	0	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.78D Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2010-2011 and 2012-2013

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	0.17	0.18	0.15	0.18	0.18	0.19
AGE						
15-17	0.18	0.18	1.09	*	0.19	0.18
18-25	0.28	0.27	0.39	0.17	0.30	0.28
26-44	0.24	0.26	*	0.28	0.25	0.27
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.20	0.21	0.18	0.22	0.20	0.22
White	0.24	0.26	0.22	0.21	0.25	0.27
Black or African American	0.40	0.44	0.39	0.85	0.42	0.46
American Indian or Alaska Native	2.82	1.65	*	*	2.91	1.73
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	0.40	0.53	*	*	0.41	0.56
Two or More Races	0.75	0.81	*	*	0.79	0.85
Hispanic or Latino	0.34	0.38	0.28	*	0.35	0.39
TRIMESTER²						
First	N/A	N/A	0.29	0.37	N/A	N/A
Second	N/A	N/A	0.35	0.38	N/A	N/A
Third	N/A	N/A	*	0.03	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.

Table 6.78P Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: P Values from Tests of Differences of Percentages, 2012-2013 Versus 2010-2011

Demographic/Pregnancy Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2010-2011	2012-2013	2010-2011	2012-2013	2010-2011	2012-2013
TOTAL	0.8968		0.9148		0.8651	
AGE						
15-17	0.2936		*		0.3103	
18-25	0.1084		0.0945		0.1070	
26-44	0.5054		*		0.5456	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.9469		0.8120		0.9706	
White	0.7309		0.8446		0.6918	
Black or African American	0.1638		0.4784		0.1714	
American Indian or Alaska Native	0.1435		*		0.1634	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.3983		*		0.4065	
Two or More Races	0.7694		*		0.7880	
Hispanic or Latino	0.8675		*		0.8664	
TRIMESTER²						
First	N/A		0.5576		N/A	
Second	N/A		0.8310		N/A	
Third	N/A		*		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column represent all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010-2013.