

**Table 1.11A Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2009 and 2010**

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOTAL</b>	118,705	119,508	37,954	38,806	21,813	22,622
12	384	382	243	250	132	121
13	584	630	359	427	137 <sup>a</sup>	195
14	878	782	604	571	291	281
15	1,338	1,256	1,009	953	479	471
16	1,560	1,424	1,194	1,110	633	619
17	1,845	1,777	1,400	1,419	798	780
18	2,376	2,325	1,833	1,794	1,057	1,048
19	2,478	2,347	1,781	1,706	987	1,083
20	2,431	2,561	1,592	1,667	985	1,029
21	2,519	2,493	1,651	1,599	974	986
22	2,509	2,410	1,522	1,445	905	908
23	2,434	2,400	1,331	1,308	765	809
24	2,544	2,447	1,333	1,246	821	776
25	2,218	2,435	1,051	1,171	631	690
26-29	10,204	10,385	4,290	4,228	2,424	2,460
30-34	11,279	11,413	3,535	3,957	2,043 <sup>a</sup>	2,571
35-39	10,534	10,360	2,703	2,735	1,584	1,564
40-44	11,929	11,717	2,637	2,627	1,364	1,413
45-49	13,678	13,346	2,624	2,784	1,459	1,595
50-54	13,383	13,325	2,382	2,574	1,539	1,588
55-59	9,534	10,548	1,553	1,777	998	787
60-64	6,425	6,504	801	777	483	437
65 or Older	5,641	6,239	527	680	323	414

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.11B Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2009 and 2010**

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOTAL</b>	47.1	47.1	15.1	15.3	8.7	8.9
12	10.5	10.0	6.6	6.6	3.6	3.2
13	15.2	15.5	9.3	10.5	3.6	4.8
14	21.2	20.1	14.6	14.7	7.0	7.2
15	30.4	30.2	22.9	22.9	10.9	11.3
16	37.1	34.3	28.4	26.7	15.1	14.9
17	42.4	41.9	32.2	33.4	18.3	18.4
18	48.0	49.0	37.0	37.8	21.3	22.1
19	54.2	53.6	39.0	38.9	21.6 <sup>a</sup>	24.7
20	59.0	56.1	38.6	36.5	23.9	22.5
21	60.8	58.5	39.8	37.5	23.5	23.1
22	62.2 <sup>a</sup>	58.6	37.7	35.1	22.4	22.1
23	60.7	59.7	33.2	32.5	19.1	20.1
24	61.8	60.4	32.4	30.7	19.9	19.1
25	61.2	62.1	29.0	29.8	17.4	17.6
26-29	60.7	62.3	25.5	25.4	14.4	14.8
30-34	58.1	57.5	18.2	19.9	10.5 <sup>a</sup>	12.9
35-39	53.0	53.6	13.6	14.2	8.0	8.1
40-44	57.2	57.2	12.6	12.8	6.5	6.9
45-49	60.9	60.0	11.7	12.5	6.5	7.2
50-54	60.1	60.5	10.7	11.7	6.9	7.2
55-59	51.5	54.8	8.4	9.2	5.4	4.1
60-64	41.5	39.6	5.2	4.7	3.1	2.7
65 or Older	14.9	16.0	1.4	1.7	0.9	1.1

\*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.12A Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2009 and 2010**

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOTAL</b>	104,446	106,232	28,521	29,206	16,718	17,373
12	50	39	33	35	17	10
13	158	191	118	162	45	60
14	443	411	365	354	183	165
15	870	838	718	705	354	356
16	1,127	1,122	928	906	507	516
17	1,532	1,538	1,176	1,238	693	690
18	2,080	2,020	1,582	1,559	897	915
19	2,206	2,076	1,565	1,475	897	946
20	2,169	2,265	1,377	1,449	856	913
21	2,273	2,272	1,399	1,358	829	825
22	2,271	2,184	1,283	1,238	774	795
23	2,242	2,210	1,103	1,123	637	687
24	2,343	2,202	1,109	1,034	670	663
25	1,961 <sup>a</sup>	2,186	842	924	523	548
26-29	9,138	9,433	3,436	3,329	1,914	1,897
30-34	10,062	10,254	2,609	3,083	1,565 <sup>a</sup>	1,955
35-39	9,293	9,057	1,871	1,891	1,065	1,127
40-44	10,807	10,623	1,710	1,692	878	1,012
45-49	12,490	12,497	1,786	1,871	1,070	1,118
50-54	12,532	12,379	1,691	1,598	1,093	1,072
55-59	8,572	9,671	1,048	1,253	720	602
60-64	5,565	5,992	511	541	322	316
65 or Older	4,264	4,771	262	385	209	187

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.12B Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2009 and 2010**

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOTAL</b>	41.5	41.9	11.3	11.5	6.6	6.9
12	1.4	1.0	0.9	0.9	0.5	0.2
13	4.1	4.7	3.1	4.0	1.2	1.5
14	10.7	10.6	8.8	9.1	4.4	4.2
15	19.8	20.2	16.3	17.0	8.0	8.6
16	26.8	27.0	22.1	21.8	12.1	12.4
17	35.2	36.2	27.0	29.2	15.9	16.2
18	42.0	42.6	31.9	32.9	18.1	19.3
19	48.3	47.4	34.2	33.7	19.6	21.6
20	52.6	49.6	33.4	31.7	20.8	20.0
21	54.8	53.3	33.7	31.8	20.0	19.3
22	56.3	53.1	31.8	30.1	19.2	19.3
23	55.9	55.0	27.5	27.9	15.9	17.1
24	56.9	54.3	26.9	25.5	16.3	16.4
25	54.1	55.7	23.2	23.6	14.4	14.0
26-29	54.4	56.6	20.4	20.0	11.4	11.4
30-34	51.8	51.6	13.4	15.5	8.1	9.8
35-39	46.8	46.9	9.4	9.8	5.4	5.8
40-44	51.8	51.8	8.2	8.3	4.2	4.9
45-49	55.6	56.2	8.0	8.4	4.8	5.0
50-54	56.3	56.2	7.6	7.3	4.9	4.9
55-59	46.3	50.2	5.7	6.5	3.9	3.1
60-64	36.0	36.5	3.3	3.3	2.1	1.9
65 or Older	11.2	12.3	0.7	1.0	0.6	0.5

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.13A Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2009 and 2010**

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOTAL</b>	36,599	37,210	4,797	4,499	1,637	1,466
12	10 <sup>a</sup>	1	7	0	6	*
13	4	10	3	2	0	*
14	20	22	12	14	3	1
15	62	73	43	57	6	15
16	107	101	61	71	18	22
17	200	156	114	94	35	22
18	333	285	178	157	38	47
19	483 <sup>a</sup>	370	239	204	61	58
20	546	505	217	199	51	56
21	707	607	302	234	84	92
22	735	678	286	247	82	63
23	695	703	179	167	58	63
24	783	681	193	184	50	53
25	682	699	178	183	38	64
26-29	3,116	3,175	660	657	165	157
30-34	3,452	3,791	556	645	203	247
35-39	3,295	3,126	317	343	112	116
40-44	4,553	4,508	306	329	139	96
45-49	5,955	6,285	361	321	199	154
50-54	5,457	5,473	386	231	161	112
55-59	3,228	3,625	105	103	63	4
60-64	1,363	1,474	80	56	62	24
65 or Older	812	864	13	*	*	*

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.13B Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2009 and 2010**

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOTAL</b>	14.5	14.7	1.9	1.8	0.7	0.6
12	0.3 <sup>a</sup>	0.0	0.2	0.0	0.2	*
13	0.1	0.2	0.1	0.1	0.0	*
14	0.5	0.6	0.3	0.4	0.1	0.0
15	1.4	1.8	1.0	1.4	0.1	0.4
16	2.5	2.4	1.4	1.7	0.4	0.5
17	4.6	3.7	2.6	2.2	0.8	0.5
18	6.7	6.0	3.6	3.3	0.8	1.0
19	10.6 <sup>a</sup>	8.4	5.2	4.7	1.3	1.3
20	13.3	11.1	5.3	4.4	1.2	1.2
21	17.1	14.2	7.3	5.5	2.0	2.2
22	18.2	16.5	7.1	6.0	2.0	1.5
23	17.3	17.5	4.4	4.2	1.4	1.6
24	19.0	16.8	4.7	4.5	1.2	1.3
25	18.8	17.8	4.9	4.7	1.0	1.6
26-29	18.5	19.1	3.9	3.9	1.0	0.9
30-34	17.8	19.1	2.9	3.2	1.0	1.2
35-39	16.6	16.2	1.6	1.8	0.6	0.6
40-44	21.8	22.0	1.5	1.6	0.7	0.5
45-49	26.5	28.3	1.6	1.4	0.9	0.7
50-54	24.5	24.9	1.7	1.0	0.7	0.5
55-59	17.4	18.8	0.6	0.5	0.3	0.0
60-64	8.8	9.0	0.5	0.3	0.4	0.1
65 or Older	2.1	2.2	0.0	*	*	*

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.14A Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2009 and 2010**

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOTAL</b>	8,359	9,158	1,016	871	492	378
12	2	1	2	0	2	*
13	2	5	2	0	0	*
14	4	6	4	1	2	*
15	9	9	7	4	3	2
16	22 <sup>a</sup>	7	6	4	0	0
17	18	32	8	9	1	1
18	43	51	19	26	2	10
19	57	57	14	31	7	14
20	92	86	20	8	5	*
21	145	117	32	17	3	7
22	121	139	19	31	0 <sup>b</sup>	16
23	166	151	18	22	5	16
24	180	129	14	18	12	5
25	157	146	32	23	5	5
26-29	790	758	105	96	43	13
30-34	926	1,106	84	37	49	4
35-39	1,054	959	115	107	44	37
40-44	1,128	1,299	113	132	75	68
45-49	1,380 <sup>a</sup>	1,839	145	166	92	91
50-54	1,250	1,248	185	90	78	73
55-59	567	648	35	4	29	4
60-64	191	290	34	44	34	13
65 or Older	54	76	*	*	*	*

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.14B Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2009 and 2010**

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOTAL</b>	3.3	3.6	0.4	0.3	0.2	0.1
12	0.1	0.0	0.1	0.0	0.0	*
13	0.1	0.1	0.0	0.0	0.0	*
14	0.1	0.1	0.1	0.0	0.0	*
15	0.2	0.2	0.1	0.1	0.1	0.0
16	0.5 <sup>a</sup>	0.2	0.2	0.1	0.0	0.0
17	0.4	0.8	0.2	0.2	0.0	0.0
18	0.9	1.1	0.4	0.6	0.0	0.2
19	1.2	1.3	0.3	0.7	0.2	0.3
20	2.2	1.9	0.5	0.2	0.1	*
21	3.5	2.7	0.8	0.4	0.1	0.2
22	3.0	3.4	0.5	0.7	0.0 <sup>b</sup>	0.4
23	4.1	3.8	0.5	0.6	0.1	0.4
24	4.4	3.2	0.3	0.5	0.3	0.1
25	4.3	3.7	0.9	0.6	0.1	0.1
26-29	4.7	4.5	0.6	0.6	0.3	0.1
30-34	4.8	5.6	0.4	0.2	0.3	0.0
35-39	5.3	5.0	0.6	0.6	0.2	0.2
40-44	5.4	6.3	0.5	0.6	0.4	0.3
45-49	6.1 <sup>a</sup>	8.3	0.6	0.7	0.4	0.4
50-54	5.6	5.7	0.8	0.4	0.3	0.3
55-59	3.1	3.4	0.2	0.0	0.2	0.0
60-64	1.2	1.8	0.2	0.3	0.2	0.1
65 or Older	0.1	0.2	*	*	*	*

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.



**Table 1.15A Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2009 and 2010**

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOTAL</b>	37,256	37,476	4,509	4,517	1,258	1,208
12	27	17	6	8	3	0
13	58	49	35	33	10	16
14	101	91	79	74	20	35
15	203	181	143	151	45	53
16	251	275	193	200	58	39
17	372	385	278	280	81	72
18	596	541	404	362	124	117
19	666	632	378	367	110	115
20	763	763	362	347	86	86
21	817	841	367	354	82	109
22	899	835	355	353	109	74
23	897	901	208	261	30 <sup>a</sup>	61
24	902	872	171	227	36	71
25	825	877	163	152	41	40
26-29	4,094	3,920	503	507	143	146
30-34	4,577	4,740	465	440	196	97
35-39	3,344	3,525	210	117	55 <sup>a</sup>	9
40-44	3,449	3,641	75	76	*	29
45-49	4,456	4,386	41	71	3	21
50-54	4,919	4,387	32	113	26	17
55-59	3,086	3,522	13	*	*	*
60-64	1,447	1,596	*	14	*	*
65 or Older	507	499	25	10	*	*

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.15B Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2009 and 2010**

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOTAL</b>	14.8	14.8	1.8	1.8	0.5	0.5
12	0.7	0.4	0.2	0.2	0.1	0.0
13	1.5	1.2	0.9	0.8	0.3	0.4
14	2.4	2.3	1.9	1.9	0.5	0.9
15	4.6	4.4	3.3	3.6	1.0	1.3
16	6.0	6.6	4.6	4.8	1.4	0.9
17	8.6	9.1	6.4	6.6	1.9	1.7
18	12.0	11.4	8.2	7.6	2.5	2.5
19	14.6	14.4	8.3	8.4	2.4	2.6
20	18.5	16.7	8.8	7.6	2.1	1.9
21	19.7	19.7	8.9	8.3	2.0	2.6
22	22.3	20.3	8.8	8.6	2.7	1.8
23	22.4	22.4	5.2	6.5	0.7 <sup>a</sup>	1.5
24	21.9	21.5	4.2	5.6	0.9	1.8
25	22.8	22.3	4.5	3.9	1.1	1.0
26-29	24.4	23.5	3.0	3.0	0.8	0.9
30-34	23.6	23.9	2.4	2.2	1.0	0.5
35-39	16.8	18.3	1.1	0.6	0.3 <sup>a</sup>	0.0
40-44	16.5	17.8	0.4	0.4	*	0.1
45-49	19.8	19.7	0.2	0.3	0.0	0.1
50-54	22.1	19.9	0.1	0.5	0.1	0.1
55-59	16.7	18.3	0.1	*	*	*
60-64	9.4	9.7	*	0.1	*	*
65 or Older	1.3	1.3	0.1	0.0	*	*

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.16A Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2009 and 2010**

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOTAL</b>	22,448	21,660	2,090	2,030	560	685
12	245	240	135	119	60	41
13	335	352	166	192	46	72
14	431 <sup>a</sup>	334	191	176	34	53
15	461	413	175	178	33	48
16	416	346	175 <sup>a</sup>	122	42	26
17	379	320	113	95	30	23
18	408	385	137	132	24	25
19	390	372	105	97	28	16
20	378	420	63	77	24	11
21	519	452	111	74	20	19
22	487	463	70	75	16	23
23	463	440	70	66	11	21
24	496 <sup>a</sup>	384	41	46	10	14
25	439	447	28	23	16	6
26-29	2,085	2,005	134	84	47	26
30-34	2,881	2,828	61	123	15	54
35-39	2,435	2,370	64	61	38	29
40-44	2,259	2,245	30	68	9	31
45-49	2,609	2,640	81	38	28	33
50-54	2,177	2,072	115	146	31	94
55-59	1,114	1,338	24	3	*	3
60-64	694	527	*	18	*	18
65 or Older	347	268	*	19	*	*

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.16B Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2009 and 2010**

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOTAL</b>	8.9	8.5	0.8	0.8	0.2	0.3
12	6.7	6.3	3.7	3.1	1.6	1.1
13	8.7	8.6	4.3	4.7	1.2	1.8
14	10.4 <sup>a</sup>	8.6	4.6	4.5	0.8	1.4
15	10.5	9.9	4.0	4.3	0.7	1.1
16	9.9	8.3	4.2 <sup>a</sup>	2.9	1.0	0.6
17	8.7	7.5	2.6	2.2	0.7	0.6
18	8.2	8.1	2.8	2.8	0.5	0.5
19	8.5	8.5	2.3	2.2	0.6	0.4
20	9.2	9.2	1.5	1.7	0.6	0.2
21	12.5	10.6	2.7	1.7	0.5	0.4
22	12.1	11.3	1.7	1.8	0.4	0.6
23	11.5	11.0	1.7	1.6	0.3	0.5
24	12.0 <sup>a</sup>	9.5	1.0	1.1	0.2	0.3
25	12.1	11.4	0.8	0.6	0.4	0.2
26-29	12.4	12.0	0.8	0.5	0.3	0.2
30-34	14.8	14.2	0.3	0.6	0.1	0.3
35-39	12.3	12.3	0.3	0.3	0.2	0.1
40-44	10.8	11.0	0.1	0.3	0.0	0.2
45-49	11.6	11.9	0.4	0.2	0.1	0.1
50-54	9.8	9.4	0.5	0.7	0.1	0.4
55-59	6.0	6.9	0.1	0.0	*	0.0
60-64	4.5	3.2	*	0.1	*	0.1
65 or Older	0.9	0.7	*	0.0	*	*

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.17A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2009 and 2010**

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOTAL</b>	51,771	51,641	16,006	16,031	6,953	6,967
12	154	179	104	118	61	75
13	218	239	140	171	59	81
14	348	291	233	199	109	98
15	572	526	419	369	174	142
16	699	601	462	437	172	162
17	805	747	526	517	196	166
18	1,138	1,036	735	673	315	278
19	1,121	1,092	683	659	253	266
20	1,192	1,234	677	659	278	268
21	1,294	1,286	723	691	288	292
22	1,352	1,270	644	636	306	262
23	1,327	1,224	570	504	230	210
24	1,310	1,290	540	523	246	214
25	1,084 <sup>b</sup>	1,347	456	516	203	211
26-29	5,183	5,142	1,816	1,788	780	739
30-34	5,395	5,415	1,590	1,673	610 <sup>a</sup>	854
35-39	4,377	4,467	1,254	1,167	629	497
40-44	4,833	4,638	1,131	1,132	535	485
45-49	5,888	5,486	1,107	1,093	433	499
50-54	5,364	5,656	930	1,236	446	613
55-59	3,779	4,317	676	658	350	200
60-64	2,259	1,974	347	247	165	122
65 or Older	2,078	2,182	242	364	114	232

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.17B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2009 and 2010**

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOTAL</b>	20.6	20.4	6.4	6.3	2.8	2.7
12	4.2	4.7	2.8	3.1	1.7	2.0
13	5.7	5.9	3.6	4.2	1.5	2.0
14	8.4	7.5	5.6	5.1	2.6	2.5
15	13.0	12.7	9.5	8.9	4.0	3.4
16	16.6	14.4	11.0	10.5	4.1	3.9
17	18.5	17.6	12.1	12.2	4.5	3.9
18	23.0	21.8	14.8	14.2	6.4	5.9
19	24.5	24.9	14.9	15.0	5.5	6.1
20	28.9	27.0	16.4	14.4	6.8	5.9
21	31.2	30.2	17.4	16.2	6.9	6.9
22	33.5	30.9	16.0	15.5	7.6	6.4
23	33.1	30.4	14.2	12.5	5.7	5.2
24	31.8	31.8	13.1	12.9	6.0	5.3
25	29.9 <sup>b</sup>	34.3	12.6	13.2	5.6	5.4
26-29	30.8	30.9	10.8	10.7	4.6	4.4
30-34	27.8	27.3	8.2	8.4	3.1 <sup>a</sup>	4.3
35-39	22.0	23.1	6.3	6.0	3.2	2.6
40-44	23.2	22.6	5.4	5.5	2.6	2.4
45-49	26.2	24.7	4.9	4.9	1.9	2.2
50-54	24.1	25.7	4.2	5.6	2.0	2.8
55-59	20.4	22.4	3.7	3.4	1.9	1.0
60-64	14.6	12.0	2.2	1.5	1.1	0.7
65 or Older	5.5	5.6	0.6	0.9	0.3	0.6

\*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.18A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2009 and 2010**

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOTAL</b>	35,046	34,776	12,405	12,213	5,257	5,100
12	126	154	85	107	50	71
13	185	204	117	149	55	67
14	282	243	206	160	83	83
15	481	463	364	314	148	121
16	590	537	380	374	150	124
17	712	636	464	417	167	144
18	983	865	598	528	258	222
19	956	943	563	530	208	189
20	994	1,038	522	512	216	202
21	1,070	1,041	554	517	213	211
22	1,090	1,024	496	502	209	206
23	1,096	1,016	457	369	163	154
24	1,089	1,056	446	419	180	161
25	879 <sup>b</sup>	1,115	365	391	154	173
26-29	4,088	4,083	1,467	1,468	557	543
30-34	4,061	4,131	1,215	1,365	476	632
35-39	2,989	3,341	921	916	509	418
40-44	3,033	3,084	802	795	403	345
45-49	3,359	3,112	843	821	297	392
50-54	3,254	2,745	689	758	315	367
55-59	2,043	1,930	504	451	239	104
60-64	1,030	1,150	190	178	126	67
65 or Older	657	864	159	171	83	103

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.18B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2009 and 2010**

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOTAL</b>	13.9	13.7	4.9	4.8	2.1	2.0
12	3.4	4.0	2.3	2.8	1.4	1.9
13	4.8	5.0	3.1	3.7	1.4	1.6
14	6.8	6.3	5.0	4.1	2.0	2.1
15	10.9	11.1	8.3	7.6	3.4	2.9
16	14.0	12.9	9.0	9.0	3.6	3.0
17	16.4	15.0	10.7	9.8	3.8	3.4
18	19.8	18.2	12.1	11.1	5.2	4.7
19	20.9	21.5	12.3	12.1	4.6	4.3
20	24.1	22.7	12.7	11.2	5.2	4.4
21	25.8	24.4	13.4	12.1	5.1	4.9
22	27.0	24.9	12.3	12.2	5.2	5.0
23	27.3	25.3	11.4 <sup>a</sup>	9.2	4.1	3.8
24	26.4	26.0	10.8	10.3	4.4	4.0
25	24.3 <sup>b</sup>	28.4	10.1	10.0	4.3	4.4
26-29	24.3	24.5	8.7	8.8	3.3	3.3
30-34	20.9	20.8	6.3	6.9	2.5	3.2
35-39	15.0	17.3	4.6	4.7	2.6	2.2
40-44	14.5	15.0	3.8	3.9	1.9	1.7
45-49	15.0	14.0	3.8	3.7	1.3	1.8
50-54	14.6	12.5	3.1	3.4	1.4	1.7
55-59	11.0	10.0	2.7	2.3	1.3	0.5
60-64	6.7	7.0	1.2	1.1	0.8	0.4
65 or Older	1.7	2.2	0.4	0.4	0.2	0.3

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.