

Table 1.49A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Numbers in Thousands, 2009 and 2010

| Demographic Characteristic | Lifetime (2009) | Lifetime (2010) | Past Year (2009) | Past Year (2010) | Past Month (2009) | Past Month (2010) |
|---|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 51,771 | 51,641 | 16,006 | 16,031 | 6,953 | 6,967 |
| AGE | | | | | | |
| 12-17 | 2,796 ^a | 2,583 | 1,885 | 1,812 | 772 | 725 |
| 18-25 | 9,819 | 9,780 | 5,027 | 4,862 | 2,119 | 2,003 |
| 26 or Older | 39,156 | 39,277 | 9,093 | 9,357 | 4,062 | 4,239 |
| GENDER | | | | | | |
| Male | 26,979 | 27,807 | 8,445 | 8,609 | 3,840 | 3,686 |
| Female | 24,792 | 23,834 | 7,561 | 7,422 | 3,114 | 3,281 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 46,010 | 45,931 | 13,823 | 13,850 | 6,107 | 6,030 |
| White | 40,401 | 39,573 | 11,880 | 11,662 | 5,119 | 5,145 |
| Black or African American | 3,519 | 3,995 | 1,216 | 1,335 | 599 | 602 |
| American Indian or Alaska Native | 399 | 327 | 130 | 129 | 76 | 55 |
| Native Hawaiian or Other Pacific Islander | * | 70 | * | 29 | * | 18 |
| Asian | 816 | 1,084 | 302 | 440 | 155 | 111 |
| Two or More Races | 759 | 882 | 226 | 256 | 100 | 99 |
| Hispanic or Latino | 5,760 | 5,710 | 2,183 | 2,180 | 847 | 937 |

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.49B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Percentages, 2009 and 2010

| Demographic Characteristic | Lifetime (2009) | Lifetime (2010) | Past Year (2009) | Past Year (2010) | Past Month (2009) | Past Month (2010) |
|---|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 20.6 | 20.4 | 6.4 | 6.3 | 2.8 | 2.7 |
| AGE | | | | | | |
| 12-17 | 11.4 ^a | 10.6 | 7.7 | 7.4 | 3.1 | 3.0 |
| 18-25 | 29.2 | 28.7 | 15.0 | 14.3 | 6.3 | 5.9 |
| 26 or Older | 20.2 | 20.1 | 4.7 | 4.8 | 2.1 | 2.2 |
| GENDER | | | | | | |
| Male | 22.1 | 22.5 | 6.9 | 7.0 | 3.1 | 3.0 |
| Female | 19.1 | 18.3 | 5.8 | 5.7 | 2.4 | 2.5 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 21.3 | 21.2 | 6.4 | 6.4 | 2.8 | 2.8 |
| White | 23.8 | 23.3 | 7.0 | 6.9 | 3.0 | 3.0 |
| Black or African American | 11.7 | 13.2 | 4.0 | 4.4 | 2.0 | 2.0 |
| American Indian or Alaska Native | 32.6 | 27.2 | 10.6 | 10.7 | 6.2 | 4.6 |
| Native Hawaiian or Other Pacific Islander | * | 9.6 | * | 4.0 | * | 2.4 |
| Asian | 7.4 | 9.5 | 2.7 | 3.8 | 1.4 | 1.0 |
| Two or More Races | 25.7 | 27.8 | 7.6 | 8.1 | 3.4 | 3.1 |
| Hispanic or Latino | 16.0 | 15.5 | 6.1 | 5.9 | 2.4 | 2.5 |

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.50A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Numbers in Thousands, 2009 and 2010

| Demographic Characteristic | Lifetime (2009) | Lifetime (2010) | Past Year (2009) | Past Year (2010) | Past Month (2009) | Past Month (2010) |
|---|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 2,796 ^a | 2,583 | 1,885 | 1,812 | 772 | 725 |
| GENDER | | | | | | |
| Male | 1,345 ^a | 1,200 | 876 | 811 | 352 | 288 |
| Female | 1,450 | 1,383 | 1,009 | 1,001 | 420 | 437 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 2,218 ^b | 2,013 | 1,515 | 1,414 | 615 | 560 |
| White | 1,639 | 1,502 | 1,158 | 1,095 | 446 | 411 |
| Black or African American | 405 | 343 | 260 | 208 | 129 | 94 |
| American Indian or Alaska Native | 22 | 22 | 14 | 15 | 4 | 4 |
| Native Hawaiian or Other Pacific Islander | * | 7 | * | * | * | 1 |
| Asian | 79 | 58 | 45 | 35 | 19 | 19 |
| Two or More Races | 59 | 81 | 32 ^a | 59 | 14 | 31 |
| Hispanic or Latino | 578 | 571 | 370 | 397 | 157 | 165 |
| GENDER/RACE/HISPANIC ORIGIN | | | | | | |
| Male, White, Not Hispanic | 779 ^a | 670 | 543 | 479 | 209 | 164 |
| Female, White, Not Hispanic | 860 | 831 | 615 | 616 | 237 | 246 |
| Male, Black, Not Hispanic | 205 | 149 | 126 ^a | 82 | 56 | 37 |
| Female, Black, Not Hispanic | 201 | 194 | 133 | 126 | 73 | 57 |
| Male, Hispanic | 282 | 305 | 174 | 204 | 74 | 70 |
| Female, Hispanic | 296 | 266 | 196 | 193 | 83 | 95 |

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.50B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Percentages, 2009 and 2010

| Demographic Characteristic | Lifetime (2009) | Lifetime (2010) | Past Year (2009) | Past Year (2010) | Past Month (2009) | Past Month (2010) |
|---|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 11.4 ^a | 10.6 | 7.7 | 7.4 | 3.1 | 3.0 |
| GENDER | | | | | | |
| Male | 10.7 ^a | 9.6 | 7.0 | 6.5 | 2.8 | 2.3 |
| Female | 12.0 | 11.6 | 8.4 | 8.4 | 3.5 | 3.7 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 11.2 ^a | 10.4 | 7.6 | 7.3 | 3.1 | 2.9 |
| White | 11.4 | 10.7 | 8.1 | 7.8 | 3.1 | 2.9 |
| Black or African American | 10.9 | 9.6 | 7.0 | 5.8 | 3.5 | 2.6 |
| American Indian or Alaska Native | 16.8 | 16.9 | 11.1 | 11.5 | 3.0 | 3.4 |
| Native Hawaiian or Other Pacific Islander | * | 9.7 | * | * | * | 1.6 |
| Asian | 8.0 | 5.5 | 4.5 | 3.3 | 1.9 | 1.8 |
| Two or More Races | 11.2 | 15.4 | 6.1 ^a | 11.2 | 2.6 | 5.8 |
| Hispanic or Latino | 12.0 | 11.6 | 7.7 | 8.1 | 3.3 | 3.4 |
| GENDER/RACE/HISPANIC ORIGIN | | | | | | |
| Male, White, Not Hispanic | 10.6 ^a | 9.3 | 7.4 | 6.6 | 2.8 | 2.3 |
| Female, White, Not Hispanic | 12.3 | 12.1 | 8.8 | 9.0 | 3.4 | 3.6 |
| Male, Black, Not Hispanic | 11.0 | 8.3 | 6.8 ^a | 4.6 | 3.0 | 2.1 |
| Female, Black, Not Hispanic | 10.9 | 10.9 | 7.2 | 7.1 | 3.9 | 3.2 |
| Male, Hispanic | 11.4 | 12.0 | 7.1 | 8.0 | 3.0 | 2.8 |
| Female, Hispanic | 12.6 | 11.2 | 8.4 | 8.2 | 3.5 | 4.0 |

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.51A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Numbers in Thousands, 2009 and 2010

| Demographic Characteristic | Lifetime (2009) | Lifetime (2010) | Past Year (2009) | Past Year (2010) | Past Month (2009) | Past Month (2010) |
|---|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 9,819 | 9,780 | 5,027 | 4,862 | 2,119 | 2,003 |
| GENDER | | | | | | |
| Male | 5,205 | 5,247 | 2,787 | 2,676 | 1,209 | 1,100 |
| Female | 4,614 | 4,533 | 2,240 | 2,186 | 910 | 903 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 8,402 | 8,323 | 4,331 | 4,220 | 1,807 | 1,754 |
| White | 7,237 | 7,130 | 3,752 | 3,652 | 1,567 | 1,515 |
| Black or African American | 736 | 731 | 361 | 351 | 148 | 163 |
| American Indian or Alaska Native | 85 | 59 | 47 | 28 | 30 | 19 |
| Native Hawaiian or Other Pacific Islander | * | * | * | 11 | * | * |
| Asian | 170 | 196 | 93 | 92 | 33 | 26 |
| Two or More Races | 146 | 181 | 62 | 86 | 23 | 25 |
| Hispanic or Latino | 1,417 | 1,457 | 696 | 642 | 312 | 249 |
| EDUCATION | | | | | | |
| < High School | 1,924 ^a | 1,699 | 994 | 866 | 480 | 416 |
| High School Graduate | 3,424 | 3,228 | 1,776 | 1,605 | 738 | 711 |
| Some College | 3,176 | 3,421 | 1,660 | 1,720 | 675 | 682 |
| College Graduate | 1,295 | 1,432 | 597 | 671 | 226 | 194 |
| CURRENT EMPLOYMENT | | | | | | |
| Full-Time | 3,774 | 3,783 | 1,793 | 1,768 | 764 | 771 |
| Part-Time | 2,598 | 2,744 | 1,409 | 1,421 | 611 ^a | 494 |
| Unemployed | 1,486 | 1,343 | 812 ^a | 685 | 345 | 350 |
| Other ¹ | 1,960 | 1,911 | 1,013 | 988 | 400 | 389 |

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.51B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Percentages, 2009 and 2010

| Demographic Characteristic | Lifetime (2009) | Lifetime (2010) | Past Year (2009) | Past Year (2010) | Past Month (2009) | Past Month (2010) |
|---|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 29.2 | 28.7 | 15.0 | 14.3 | 6.3 | 5.9 |
| GENDER | | | | | | |
| Male | 30.8 | 30.4 | 16.5 | 15.5 | 7.2 | 6.4 |
| Female | 27.6 | 27.0 | 13.4 | 13.0 | 5.5 | 5.4 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 30.6 | 30.3 | 15.8 | 15.3 | 6.6 | 6.4 |
| White | 35.5 | 35.1 | 18.4 | 18.0 | 7.7 | 7.4 |
| Black or African American | 15.3 | 15.0 | 7.5 | 7.2 | 3.1 | 3.3 |
| American Indian or Alaska Native | 36.7 | 32.2 | 20.3 | 15.4 | 12.9 | 10.5 |
| Native Hawaiian or Other Pacific Islander | * | * | * | 7.8 | * | * |
| Asian | 11.2 | 13.6 | 6.1 | 6.4 | 2.1 | 1.8 |
| Two or More Races | 34.1 | 33.2 | 14.5 | 15.8 | 5.4 | 4.6 |
| Hispanic or Latino | 23.1 | 22.2 | 11.4 | 9.8 | 5.1 | 3.8 |
| EDUCATION | | | | | | |
| < High School | 31.6 ^a | 28.8 | 16.3 | 14.7 | 7.9 | 7.0 |
| High School Graduate | 29.2 | 27.4 | 15.1 ^a | 13.6 | 6.3 | 6.0 |
| Some College | 28.7 | 29.4 | 15.0 | 14.8 | 6.1 | 5.9 |
| College Graduate | 27.5 | 30.1 | 12.7 | 14.1 | 4.8 | 4.1 |
| CURRENT EMPLOYMENT | | | | | | |
| Full-Time | 30.5 | 30.5 | 14.5 | 14.3 | 6.2 | 6.2 |
| Part-Time | 27.7 | 28.7 | 15.0 | 14.9 | 6.5 ^a | 5.2 |
| Unemployed | 32.6 | 29.6 | 17.8 ^a | 15.1 | 7.6 | 7.7 |
| Other ¹ | 26.9 | 25.2 | 13.9 | 13.0 | 5.5 | 5.1 |

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.52A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Numbers in Thousands, 2009 and 2010

| Demographic Characteristic | Lifetime (2009) | Lifetime (2010) | Past Year (2009) | Past Year (2010) | Past Month (2009) | Past Month (2010) |
|---|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 39,156 | 39,277 | 9,093 | 9,357 | 4,062 | 4,239 |
| GENDER | | | | | | |
| Male | 20,429 | 21,360 | 4,782 | 5,123 | 2,278 | 2,299 |
| Female | 18,727 | 17,917 | 4,311 | 4,234 | 1,784 | 1,941 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 35,390 | 35,595 | 7,977 | 8,216 | 3,685 | 3,716 |
| White | 31,526 | 30,941 | 6,969 | 6,915 | 3,106 | 3,220 |
| Black or African American | 2,378 | 2,920 | 596 | 775 | 323 | 345 |
| American Indian or Alaska Native | * | 246 | * | 85 | * | 32 |
| Native Hawaiian or Other Pacific Islander | * | * | * | * | * | * |
| Asian | 567 | 830 | 164 | 314 | 103 | 66 |
| Two or More Races | 554 | 620 | 131 | 111 | 63 | 43 |
| Hispanic or Latino | 3,766 | 3,682 | 1,116 | 1,141 | 377 | 523 |
| EDUCATION | | | | | | |
| < High School | 4,947 | 4,825 | 1,538 | 1,420 | 827 | 732 |
| High School Graduate | 11,153 | 10,656 | 2,619 | 2,495 | 1,190 | 1,142 |
| Some College | 10,948 | 11,619 | 2,350 | 2,716 | 1,103 | 1,185 |
| College Graduate | 12,108 | 12,177 | 2,587 | 2,725 | 942 | 1,181 |
| CURRENT EMPLOYMENT | | | | | | |
| Full-Time | 23,654 | 23,167 | 5,114 | 5,413 | 2,047 | 2,411 |
| Part-Time | 4,896 | 5,239 | 1,204 | 1,107 | 487 | 481 |
| Unemployed | 3,025 | 2,831 | 832 | 880 | 421 | 416 |
| Other ¹ | 7,580 | 8,040 | 1,944 | 1,957 | 1,107 | 931 |

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.52B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Percentages, 2009 and 2010

| Demographic Characteristic | Lifetime (2009) | Lifetime (2010) | Past Year (2009) | Past Year (2010) | Past Month (2009) | Past Month (2010) |
|---|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 20.2 | 20.1 | 4.7 | 4.8 | 2.1 | 2.2 |
| GENDER | | | | | | |
| Male | 22.0 | 22.8 | 5.2 | 5.5 | 2.5 | 2.5 |
| Female | 18.6 | 17.7 | 4.3 | 4.2 | 1.8 | 1.9 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 21.0 | 20.9 | 4.7 | 4.8 | 2.2 | 2.2 |
| White | 23.3 | 22.8 | 5.2 | 5.1 | 2.3 | 2.4 |
| Black or African American | 11.0 | 13.4 | 2.8 | 3.6 | 1.5 | 1.6 |
| American Indian or Alaska Native | * | 27.7 | * | 9.6 | * | 3.6 |
| Native Hawaiian or Other Pacific Islander | * | * | * | * | * | * |
| Asian | 6.6 | 9.3 | 1.9 | 3.5 | 1.2 | 0.7 |
| Two or More Races | 27.7 | 29.4 | 6.6 | 5.3 | 3.2 | 2.0 |
| Hispanic or Latino | 15.1 | 14.6 | 4.5 | 4.5 | 1.5 | 2.1 |
| EDUCATION | | | | | | |
| < High School | 17.3 | 16.6 | 5.4 | 4.9 | 2.9 | 2.5 |
| High School Graduate | 19.1 | 18.4 | 4.5 | 4.3 | 2.0 | 2.0 |
| Some College | 23.5 | 24.5 | 5.0 | 5.7 | 2.4 | 2.5 |
| College Graduate | 20.2 | 20.0 | 4.3 | 4.5 | 1.6 | 1.9 |
| CURRENT EMPLOYMENT | | | | | | |
| Full-Time | 23.1 | 22.8 | 5.0 | 5.3 | 2.0 | 2.4 |
| Part-Time | 21.8 | 22.2 | 5.4 | 4.7 | 2.2 | 2.0 |
| Unemployed | 29.7 | 27.6 | 8.2 | 8.6 | 4.1 | 4.1 |
| Other ¹ | 12.9 | 13.5 | 3.3 | 3.3 | 1.9 | 1.6 |

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.53A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Numbers in Thousands, 2009 and 2010

| Demographic Characteristic | Lifetime (2009) | Lifetime (2010) | Past Year (2009) | Past Year (2010) | Past Month (2009) | Past Month (2010) |
|---|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 48,975 | 49,058 | 14,120 | 14,219 | 6,182 | 6,242 |
| GENDER | | | | | | |
| Male | 25,634 | 26,607 | 7,569 | 7,799 | 3,488 | 3,399 |
| Female | 23,341 | 22,451 | 6,552 | 6,420 | 2,694 | 2,844 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 43,793 | 43,918 | 12,308 | 12,436 | 5,492 | 5,470 |
| White | 38,763 | 38,071 | 10,722 | 10,567 | 4,673 | 4,735 |
| Black or African American | 3,114 | 3,651 | 956 | 1,127 | 471 | 508 |
| American Indian or Alaska Native | 377 | 305 | 116 | 114 | 72 | 51 |
| Native Hawaiian or Other Pacific Islander | * | 64 | * | 27 | * | * |
| Asian | 737 | 1,026 | 257 | 405 | 136 | 92 |
| Two or More Races | 700 | 801 | 194 | 197 | 87 | 68 |
| Hispanic or Latino | 5,182 | 5,139 | 1,812 | 1,783 | 690 | 772 |
| EDUCATION | | | | | | |
| < High School | 6,871 | 6,524 | 2,532 | 2,286 | 1,307 | 1,148 |
| High School Graduate | 14,577 | 13,884 | 4,395 | 4,100 | 1,928 | 1,854 |
| Some College | 14,124 | 15,040 | 4,009 | 4,436 | 1,779 | 1,866 |
| College Graduate | 13,403 | 13,609 | 3,184 | 3,396 | 1,168 | 1,375 |
| CURRENT EMPLOYMENT | | | | | | |
| Full-Time | 27,428 | 26,950 | 6,907 | 7,181 | 2,811 | 3,182 |
| Part-Time | 7,494 | 7,983 | 2,612 | 2,528 | 1,098 | 974 |
| Unemployed | 4,512 | 4,174 | 1,644 | 1,564 | 766 | 766 |
| Other ¹ | 9,540 | 9,951 | 2,957 | 2,945 | 1,507 | 1,321 |

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.53B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Percentages, 2009 and 2010

| Demographic Characteristic | Lifetime (2009) | Lifetime (2010) | Past Year (2009) | Past Year (2010) | Past Month (2009) | Past Month (2010) |
|---|--------------------|--------------------|---------------------|---------------------|----------------------|----------------------|
| TOTAL | 21.6 | 21.4 | 6.2 | 6.2 | 2.7 | 2.7 |
| GENDER | | | | | | |
| Male | 23.4 | 24.0 | 6.9 | 7.0 | 3.2 | 3.1 |
| Female | 19.9 | 19.0 | 5.6 | 5.4 | 2.3 | 2.4 |
| HISPANIC ORIGIN AND RACE | | | | | | |
| Not Hispanic or Latino | 22.3 | 22.2 | 6.3 | 6.3 | 2.8 | 2.8 |
| White | 24.9 | 24.4 | 6.9 | 6.8 | 3.0 | 3.0 |
| Black or African American | 11.8 | 13.7 | 3.6 | 4.2 | 1.8 | 1.9 |
| American Indian or Alaska Native | 34.4 | 28.4 | 10.6 | 10.6 | 6.6 | 4.7 |
| Native Hawaiian or Other Pacific Islander | * | 9.6 | * | 4.0 | * | * |
| Asian | 7.3 | 9.9 | 2.6 | 3.9 | 1.4 | 0.9 |
| Two or More Races | 28.8 | 30.2 | 8.0 | 7.4 | 3.6 | 2.6 |
| Hispanic or Latino | 16.6 | 16.1 | 5.8 | 5.6 | 2.2 | 2.4 |
| EDUCATION | | | | | | |
| < High School | 19.8 | 18.6 | 7.3 | 6.5 | 3.8 | 3.3 |
| High School Graduate | 20.8 | 20.0 | 6.3 | 5.9 | 2.8 | 2.7 |
| Some College | 24.5 | 25.5 | 6.9 | 7.5 | 3.1 | 3.2 |
| College Graduate | 20.7 | 20.8 | 4.9 | 5.2 | 1.8 | 2.1 |
| CURRENT EMPLOYMENT | | | | | | |
| Full-Time | 23.9 | 23.6 | 6.0 | 6.3 | 2.4 | 2.8 |
| Part-Time | 23.6 | 24.1 | 8.2 | 7.6 | 3.5 | 2.9 |
| Unemployed | 30.6 | 28.2 | 11.1 | 10.6 | 5.2 | 5.2 |
| Other ¹ | 14.5 | 14.8 | 4.5 | 4.4 | 2.3 | 2.0 |

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.