

Table 1.59A (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	119,259	119,933	38,123	38,908	21,930	22,648
GEOGRAPHIC DIVISION						
Northeast	22,413	22,990	7,480	7,360	4,375	4,421
New England	6,625	6,615	2,370	2,210	1,452	1,413
Middle Atlantic	15,789	16,375	5,110	5,150	2,923	3,007
Midwest	26,718	26,248	7,666	7,873	4,353	4,533
East North Central	18,818	18,750	5,563	5,894	3,191	3,466
West North Central	7,900	7,497	2,104	1,978	1,162	1,068
South	41,016	41,393	12,731	12,637	7,209	7,228
South Atlantic	22,827	22,273	6,978	6,957	4,006	4,038
East South Central	6,445	6,726	1,815	1,907	1,041	1,076
West South Central	11,743	12,395	3,938	3,773	2,162	2,114
West	29,112	29,302	10,246	11,038	5,993	6,466
Mountain	8,921	9,208	2,971 ^a	3,469	1,696 ^a	2,037
Pacific	20,191	20,094	7,275	7,569	4,297	4,429
COUNTY TYPE						
Large Metro	64,271	66,195	21,025	21,977	12,133	12,801
Small Metro	37,003	36,587	11,839	11,858	6,783	6,827
250K – 1 Mil. Pop.	24,841	24,847	7,876	8,040	4,491	4,613
< 250K Pop.	12,162	11,739	3,963	3,818	2,291	2,214
Nonmetro	17,986	17,151	5,259	5,073	3,015	3,020
Urbanized	8,164	7,788	2,480	2,321	1,430	1,488
Less Urbanized	8,189	8,124	2,389	2,460	1,392	1,395
Completely Rural	1,633	1,239	390	293	193	137

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.59B (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	47.4	47.3	15.1	15.3	8.7	8.9
GEOGRAPHIC DIVISION						
Northeast	48.3	49.4	16.1	15.8	9.4	9.5
New England	54.4	54.0	19.5	18.0	11.9	11.5
Middle Atlantic	46.2	47.8	14.9	15.0	8.5	8.8
Midwest	48.4	47.4	13.9	14.2	7.9	8.2
East North Central	48.9	48.7	14.5	15.3	8.3	9.0
West North Central	47.3	44.6	12.6	11.8	7.0	6.4
South	44.6	44.5	13.8	13.6	7.8	7.8
South Atlantic	47.0	45.5	14.4	14.2	8.2	8.2
East South Central	43.0	44.6	12.1	12.6	6.9	7.1
West South Central	41.2	42.9	13.8	13.1	7.6	7.3
West	50.0	49.9	17.6	18.8	10.3	11.0
Mountain	50.1	51.1	16.7 ^a	19.2	9.5	11.3
Pacific	50.0	49.3	18.0	18.6	10.6	10.9
COUNTY TYPE						
Large Metro	48.0	48.8	15.7	16.2	9.1	9.4
Small Metro	48.2	47.2	15.4	15.3	8.8	8.8
250K – 1 Mil. Pop.	49.1	48.3	15.6	15.6	8.9	9.0
< 250K Pop.	46.6	45.2	15.2	14.7	8.8	8.5
Nonmetro	43.6	42.4	12.7	12.5	7.3	7.5
Urbanized	45.2	45.6	13.7	13.6	7.9	8.7
Less Urbanized	43.0	41.3	12.6	12.5	7.3	7.1
Completely Rural	39.0	33.3	9.3	7.9	4.6	3.7

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.60A (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
Numbers in Thousands, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	6,629 ^a	6,278	4,838	4,745	2,484	2,468
GEOGRAPHIC DIVISION						
Northeast	1,063	1,073	825	840	447	479
New England	302	307	241	242	141	142
Middle Atlantic	761	767	584	598	306	337
Midwest	1,422 ^a	1,320	1,020	986	557	506
East North Central	1,045 ^b	933	771 ^a	693	432 ^a	364
West North Central	377	387	249	293	125	141
South	2,385 ^a	2,200	1,686	1,657	796	792
South Atlantic	1,239	1,185	888	894	407	418
East South Central	390 ^a	330	286	251	129	111
West South Central	756	686	512	512	260	264
West	1,758	1,684	1,308	1,261	684	690
Mountain	515	520	386	394	191	198
Pacific	1,243	1,163	922	867	493	492
COUNTY TYPE						
Large Metro	3,521	3,353	2,574	2,571	1,322	1,405
Small Metro	2,120	2,011	1,552	1,496	819	757
250K – 1 Mil. Pop.	1,426	1,393	1,061	1,030	573	544
< 250K Pop.	694	617	491	466	246	213
Nonmetro	987	914	711	677	343	306
Urbanized	447 ^a	354	334	264	163 ^a	116
Less Urbanized	456	490	322	364	154	167
Completely Rural	84	70	55	50	25	23

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.60B (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	26.9 ^a	25.8	19.7	19.5	10.1	10.1
GEOGRAPHIC DIVISION						
Northeast	24.7	25.3	19.2	19.8	10.4	11.3
New England	27.1	28.1	21.6	22.1	12.7	13.0
Middle Atlantic	23.8	24.4	18.3	19.0	9.6	10.7
Midwest	26.3	24.7	18.8	18.5	10.3	9.5
East North Central	27.6 ^a	24.9	20.3	18.5	11.4 ^a	9.7
West North Central	23.2	24.3	15.4	18.4	7.7	8.9
South	26.5 ^a	24.6	18.7	18.5	8.8	8.8
South Atlantic	27.3	26.4	19.6	19.9	9.0	9.3
East South Central	26.8 ^a	22.8	19.6	17.4	8.8	7.7
West South Central	25.1	22.7	17.0	16.9	8.6	8.7
West	29.9	28.9	22.2	21.7	11.6	11.9
Mountain	28.6	29.1	21.4	22.0	10.6	11.1
Pacific	30.5	28.9	22.6	21.5	12.1	12.2
COUNTY TYPE						
Large Metro	27.0	25.7	19.8	19.7	10.1	10.8
Small Metro	27.6	26.7	20.2	19.9	10.7	10.1
250K – 1 Mil. Pop.	27.9	27.7	20.7	20.5	11.2	10.8
< 250K Pop.	27.2	24.8	19.2	18.7	9.6	8.5
Nonmetro	25.3	24.1	18.2	17.9	8.8	8.1
Urbanized	26.1	23.2	19.5	17.3	9.5	7.6
Less Urbanized	25.3	25.3	17.9	18.8	8.6	8.6
Completely Rural	21.6	22.1	14.2	15.7	6.5	7.3

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.61A (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:
Numbers in Thousands, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	19,658	19,538	12,186	11,990	7,176	7,345
GEOGRAPHIC DIVISION						
Northeast	3,762	3,686	2,554	2,373	1,555	1,465
New England	1,102	1,112	769	778	477	509
Middle Atlantic	2,659	2,573	1,785 ^a	1,596	1,078	956
Midwest	4,354	4,285	2,558	2,574	1,444	1,555
East North Central	3,093	2,974	1,850	1,828	1,063	1,115
West North Central	1,261	1,312	708	746	382	440
South	6,742	6,780	3,947	4,022	2,333	2,402
South Atlantic	3,497	3,497	2,112	2,166	1,215	1,325
East South Central	1,053	1,049	606	570	348	319
West South Central	2,193	2,234	1,229	1,285	770	758
West	4,799	4,787	3,128	3,020	1,845	1,923
Mountain	1,419	1,448	854	880	517	533
Pacific	3,380	3,339	2,274	2,140	1,328	1,390
COUNTY TYPE						
Large Metro	10,471	10,405	6,623	6,587	3,884	4,030
Small Metro	6,227	6,536	3,884	3,949	2,343	2,443
250K – 1 Mil. Pop.	3,966	4,233	2,443	2,541	1,441	1,554
< 250K Pop.	2,261	2,303	1,441	1,408	903	890
Nonmetro	2,960 ^a	2,597	1,678	1,454	949	872
Urbanized	1,486	1,276	855	750	494	458
Less Urbanized	1,288	1,153	720	616	393	363
Completely Rural	186	168	103	88	62	51

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.61B (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	58.5	57.3	36.3	35.2	21.4	21.6
GEOGRAPHIC DIVISION						
Northeast	61.5	61.0	41.7	39.3	25.4	24.2
New England	69.0	68.5	48.1	47.8	29.9	31.3
Middle Atlantic	58.8	58.2	39.5	36.1	23.8	21.6
Midwest	59.3	57.2	34.9	34.3	19.7	20.7
East North Central	60.9 ^a	57.9	36.4	35.6	20.9	21.7
West North Central	55.8	55.7	31.3	31.7	16.9	18.7
South	56.1	54.6	32.8	32.4	19.4	19.3
South Atlantic	57.2	55.4	34.5	34.3	19.9	21.0
East South Central	55.4	52.8	31.9	28.7	18.3	16.1
West South Central	54.7	54.3	30.7	31.2	19.2	18.4
West	59.3	59.0	38.6	37.2	22.8	23.7
Mountain	58.6	58.0	35.3	35.3	21.4	21.4
Pacific	59.5	59.5	40.0	38.1	23.4	24.8
COUNTY TYPE						
Large Metro	58.7	57.3	37.1	36.3	21.8	22.2
Small Metro	59.4	58.8	37.0	35.5	22.3	22.0
250K – 1 Mil. Pop.	59.3	60.3	36.5	36.2	21.5	22.1
< 250K Pop.	59.4	56.2	37.9	34.4	23.7	21.7
Nonmetro	56.5	54.2	32.0	30.3	18.1	18.2
Urbanized	59.1	60.4	34.0	35.5	19.7	21.7
Less Urbanized	53.6 ^a	49.5	30.0	26.4	16.3	15.6
Completely Rural	57.2	48.1	31.8	25.2	19.1	14.7

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.62A (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	92,973	94,117	21,099	22,174	12,270	12,835
GEOGRAPHIC DIVISION						
Northeast	17,588	18,231	4,102	4,147	2,373	2,476
New England	5,220	5,195	1,360	1,191	834	762
Middle Atlantic	12,368	13,036	2,741	2,956	1,540	1,714
Midwest	20,942	20,642	4,089	4,312	2,352	2,473
East North Central	14,679	14,843	2,942	3,373	1,696	1,986
West North Central	6,263	5,799	1,147	939	655	487
South	31,888	32,413	7,098	6,959	4,080	4,034
South Atlantic	18,092	17,591	3,978	3,897	2,384	2,296
East South Central	5,002	5,347	923	1,086	565	646
West South Central	8,794	9,475	2,197	1,976	1,131	1,092
West	22,555	22,831	5,811 ^a	6,756	3,465	3,852
Mountain	6,987	7,240	1,731 ^a	2,194	989 ^a	1,306
Pacific	15,568	15,591	4,080	4,562	2,476	2,546
COUNTY TYPE						
Large Metro	50,279	52,437	11,827	12,819	6,926	7,366
Small Metro	28,656	28,040	6,403	6,413	3,621	3,627
250K – 1 Mil. Pop.	19,450	19,221	4,372	4,469	2,478	2,515
< 250K Pop.	9,206	8,819	2,031	1,944	1,143	1,112
Nonmetro	14,038	13,641	2,870	2,942	1,723	1,842
Urbanized	6,231	6,158	1,291	1,307	773	914
Less Urbanized	6,445	6,481	1,347	1,480	845	866
Completely Rural	1,363	1,001	232	155	106	62

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.62B (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	48.0	48.2	10.9	11.4	6.3	6.6
GEOGRAPHIC DIVISION						
Northeast	48.9	50.3	11.4	11.4	6.6	6.8
New England	55.1	54.5	14.4	12.5	8.8	8.0
Middle Atlantic	46.7	48.8	10.3	11.1	5.8	6.4
Midwest	49.4	48.6	9.6	10.1	5.5	5.8
East North Central	49.6	50.1	9.9	11.4	5.7	6.7
West North Central	48.9	45.1	8.9	7.3	5.1	3.8
South	44.9	45.3	10.0	9.7	5.7	5.6
South Atlantic	47.7	46.1	10.5	10.2	6.3	6.0
East South Central	43.0	45.9	7.9	9.3	4.9	5.5
West South Central	41.0	43.6	10.2	9.1	5.3	5.0
West	51.0	50.9	13.1	15.1	7.8	8.6
Mountain	51.4	52.7	12.7 ^a	16.0	7.3 ^a	9.5
Pacific	50.8	50.1	13.3	14.7	8.1	8.2
COUNTY TYPE						
Large Metro	48.8	50.2	11.5	12.3	6.7	7.0
Small Metro	48.9	47.7	10.9	10.9	6.2	6.2
250K – 1 Mil. Pop.	50.1	48.8	11.3	11.3	6.4	6.4
< 250K Pop.	46.6	45.5	10.3	10.0	5.8	5.7
Nonmetro	43.7	42.8	8.9	9.2	5.4	5.8
Urbanized	45.1	45.8	9.3	9.7	5.6	6.8
Less Urbanized	43.5	42.1	9.1	9.6	5.7	5.6
Completely Rural	39.2	32.8	6.7	5.1	3.0	2.0

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.63A (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	112,631	113,655	33,285	34,163	19,446	20,180
GEOGRAPHIC DIVISION						
Northeast	21,350	21,917	6,655	6,520	3,928	3,941
New England	6,323	6,308	2,129	1,968	1,311	1,271
Middle Atlantic	15,027	15,609	4,526	4,552	2,617	2,670
Midwest	25,296	24,927	6,647	6,886	3,796	4,028
East North Central	17,773	17,817	4,792	5,201	2,759	3,101
West North Central	7,524	7,111	1,855	1,685	1,037	927
South	38,631	39,193	11,045	10,980	6,413	6,436
South Atlantic	21,588	21,088	6,089	6,063	3,599	3,621
East South Central	6,055	6,396	1,529	1,656	913	965
West South Central	10,987	11,709	3,426	3,262	1,902	1,850
West	27,354	27,618	8,938	9,776	5,310	5,775
Mountain	8,406	8,688	2,585 ^a	3,075	1,505 ^a	1,839
Pacific	18,948	18,930	6,354	6,702	3,804	3,936
COUNTY TYPE						
Large Metro	60,749	62,842	18,450	19,406	10,810	11,395
Small Metro	34,883	34,576	10,287	10,362	5,964	6,070
250K – 1 Mil. Pop.	23,415	23,454	6,815	7,011	3,919	4,068
< 250K Pop.	11,467	11,122	3,472	3,352	2,045	2,002
Nonmetro	16,999	16,238	4,548	4,396	2,672	2,714
Urbanized	7,717	7,434	2,145	2,057	1,267	1,372
Less Urbanized	7,733	7,634	2,067	2,096	1,238	1,228
Completely Rural	1,549	1,169	335	243	168	114

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.63B (REVISED) Illicit Drug Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	49.6	49.6	14.6	14.9	8.6	8.8
GEOGRAPHIC DIVISION						
Northeast	50.7	51.8	15.8	15.4	9.3	9.3
New England	57.1	56.5	19.2	17.6	11.8	11.4
Middle Atlantic	48.5	50.1	14.6	14.6	8.4	8.6
Midwest	50.8	49.8	13.4	13.8	7.6	8.1
East North Central	51.3	51.2	13.8	14.9	8.0	8.9
West North Central	49.9	46.7	12.3	11.1	6.9	6.1
South	46.5	46.7	13.3	13.1	7.7	7.7
South Atlantic	49.0	47.4	13.8	13.6	8.2	8.1
East South Central	44.7	46.9	11.3	12.1	6.7	7.1
West South Central	43.1	45.3	13.4	12.6	7.5	7.2
West	52.3	52.2	17.1	18.5	10.1	10.9
Mountain	52.5	53.5	16.1 ^a	18.9	9.4	11.3
Pacific	52.2	51.6	17.5	18.3	10.5	10.7
COUNTY TYPE						
Large Metro	50.3	51.2	15.3	15.8	8.9	9.3
Small Metro	50.5	49.4	14.9	14.8	8.6	8.7
250K – 1 Mil. Pop.	51.4	50.5	15.0	15.1	8.6	8.8
< 250K Pop.	48.7	47.4	14.7	14.3	8.7	8.5
Nonmetro	45.5	44.3	12.2	12.0	7.2	7.4
Urbanized	47.3	47.8	13.1	13.2	7.8	8.8
Less Urbanized	44.9	43.1	12.0	11.8	7.2	6.9
Completely Rural	40.8	34.3	8.8	7.1	4.4	3.3

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.