

**Table 1.74A (REVISED) Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Numbers in Thousands, 2009 and 2010**

| Geographic Characteristic | Lifetime (2009) | Lifetime (2010) | Past Year (2009) | Past Year (2010) | Past Month (2009) | Past Month (2010) |
|----------------------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 8,390 | 9,208 | 1,008 | 885 | 497 | 378 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 1,248 ^a | 1,643 | 247 | 213 | 145 | 101 |
| New England | 372 | 521 | 79 | 83 | 62 | 57 |
| Middle Atlantic | 877 | 1,122 | 167 | 130 | 83 | 44 |
| Midwest | 1,792 | 1,757 | 183 | 185 | 74 | 79 |
| East North Central | 1,142 | 1,126 | 128 | 160 | 57 | 64 |
| West North Central | 650 | 631 | 55 | 25 | 17 | 15 |
| South | 2,865 | 3,329 | 408 | 342 | 206 | 165 |
| South Atlantic | 1,481 | 1,975 | 239 | 153 | 150 | 62 |
| East South Central | 486 | 542 | 57 | 79 | 6 | 39 |
| West South Central | 898 | 813 | 111 | 110 | 50 | 64 |
| West | 2,485 | 2,479 | 169 | 144 | 72 | 33 |
| Mountain | 771 | 844 | 80 | 51 | 40 | 9 |
| Pacific | 1,714 | 1,635 | 89 | 93 | 32 | 24 |
| COUNTY TYPE | | | | | | |
| Large Metro | 4,098 | 4,619 | 495 | 469 | 226 | 202 |
| Small Metro | 2,750 | 3,145 | 342 | 349 | 221 | 155 |
| 250K – 1 Mil. Pop. | 1,772 | 2,095 | 258 | 249 | 179 | 135 |
| < 250K Pop. | 978 | 1,050 | 84 | 99 | 42 | 20 |
| Nonmetro | 1,543 | 1,444 | 171 ^b | 67 | 50 | 21 |
| Urbanized | 729 | 671 | 63 | 39 | 12 | 14 |
| Less Urbanized | 680 | 709 | 103 ^a | 26 | 37 | 7 |
| Completely Rural | 133 ^a | 64 | 4 | 1 | * | * |

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.74B (REVISED) Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2009 and 2010

| Geographic Characteristic | Lifetime (2009) | Lifetime (2010) | Past Year (2009) | Past Year (2010) | Past Month (2009) | Past Month (2010) |
|----------------------------------|----------------------------|----------------------------|-----------------------------|-----------------------------|------------------------------|------------------------------|
| TOTAL | 3.3 | 3.6 | 0.4 | 0.3 | 0.2 | 0.1 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 2.7 ^a | 3.5 | 0.5 | 0.5 | 0.3 | 0.2 |
| New England | 3.1 | 4.3 | 0.7 | 0.7 | 0.5 | 0.5 |
| Middle Atlantic | 2.6 | 3.3 | 0.5 | 0.4 | 0.2 | 0.1 |
| Midwest | 3.2 | 3.2 | 0.3 | 0.3 | 0.1 | 0.1 |
| East North Central | 3.0 | 2.9 | 0.3 | 0.4 | 0.1 | 0.2 |
| West North Central | 3.9 | 3.8 | 0.3 | 0.1 | 0.1 | 0.1 |
| South | 3.1 | 3.6 | 0.4 | 0.4 | 0.2 | 0.2 |
| South Atlantic | 3.0 | 4.0 | 0.5 | 0.3 | 0.3 | 0.1 |
| East South Central | 3.2 | 3.6 | 0.4 | 0.5 | 0.0 | 0.3 |
| West South Central | 3.2 | 2.8 | 0.4 | 0.4 | 0.2 | 0.2 |
| West | 4.3 | 4.2 | 0.3 | 0.2 | 0.1 | 0.1 |
| Mountain | 4.3 | 4.7 | 0.4 | 0.3 | 0.2 | 0.0 |
| Pacific | 4.2 | 4.0 | 0.2 | 0.2 | 0.1 | 0.1 |
| COUNTY TYPE | | | | | | |
| Large Metro | 3.1 | 3.4 | 0.4 | 0.3 | 0.2 | 0.1 |
| Small Metro | 3.6 | 4.1 | 0.4 | 0.4 | 0.3 | 0.2 |
| 250K – 1 Mil. Pop. | 3.5 | 4.1 | 0.5 | 0.5 | 0.4 | 0.3 |
| < 250K Pop. | 3.7 | 4.0 | 0.3 | 0.4 | 0.2 | 0.1 |
| Nonmetro | 3.7 | 3.6 | 0.4 ^b | 0.2 | 0.1 | 0.1 |
| Urbanized | 4.0 | 3.9 | 0.4 | 0.2 | 0.1 | 0.1 |
| Less Urbanized | 3.6 | 3.6 | 0.5 ^a | 0.1 | 0.2 | 0.0 |
| Completely Rural | 3.2 | 1.7 | 0.1 | 0.0 | * | * |

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.