

Table 1.78A (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	51,991	51,832	16,064	16,051	6,980	6,957
GEOGRAPHIC DIVISION						
Northeast	8,553	8,572	2,646	2,697	1,124	1,298
New England	2,357	2,346	774	758	303	411
Middle Atlantic	6,196	6,227	1,873	1,938	821	887
Midwest	10,639	10,678	3,289	3,268	1,368	1,383
East North Central	7,433	7,674	2,474	2,423	1,058	1,080
West North Central	3,206	3,004	815	845	311	303
South	18,725	18,348	5,960	5,602	2,675	2,524
South Atlantic	9,878	9,698	2,996	2,951	1,334	1,378
East South Central	2,991	3,241	948	874	495	404
West South Central	5,856	5,410	2,016	1,777	847	742
West	14,075	14,234	4,169	4,484	1,812	1,751
Mountain	4,362	4,523	1,389	1,460	635	603
Pacific	9,713	9,710	2,781	3,024	1,177	1,149
COUNTY TYPE						
Large Metro	27,343	28,305	8,297	8,711	3,590	3,671
Small Metro	16,609	16,190	5,235	5,100	2,187	2,214
250K – 1 Mil. Pop.	11,176	10,740	3,504	3,480	1,336	1,517
< 250K Pop.	5,433	5,450	1,731	1,620	851	697
Nonmetro	8,039	7,337	2,532	2,239	1,202	1,072
Urbanized	3,791	3,415	1,289	1,061	575	533
Less Urbanized	3,602	3,409	1,060	1,061	543	498
Completely Rural	646	513	183	117	83	41

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.78B (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	20.6	20.4	6.4	6.3	2.8	2.7
GEOGRAPHIC DIVISION						
Northeast	18.4	18.4	5.7	5.8	2.4	2.8
New England	19.3	19.2	6.4	6.2	2.5	3.4
Middle Atlantic	18.1	18.2	5.5	5.7	2.4	2.6
Midwest	19.3	19.3	6.0	5.9	2.5	2.5
East North Central	19.3	19.9	6.4	6.3	2.7	2.8
West North Central	19.2	17.9	4.9	5.0	1.9	1.8
South	20.3	19.7	6.5	6.0	2.9	2.7
South Atlantic	20.3	19.8	6.2	6.0	2.7	2.8
East South Central	20.0	21.5	6.3	5.8	3.3	2.7
West South Central	20.6	18.7	7.1	6.2	3.0	2.6
West	24.2	24.2	7.2	7.6	3.1	3.0
Mountain	24.5	25.1	7.8	8.1	3.6	3.3
Pacific	24.0	23.8	6.9	7.4	2.9	2.8
COUNTY TYPE						
Large Metro	20.4	20.9	6.2	6.4	2.7	2.7
Small Metro	21.6	20.9	6.8	6.6	2.9	2.9
250K – 1 Mil. Pop.	22.1	20.9	6.9	6.8	2.6	2.9
< 250K Pop.	20.8	21.0	6.6	6.2	3.3	2.7
Nonmetro	19.5	18.1	6.1	5.5	2.9	2.6
Urbanized	21.0	20.0	7.1	6.2	3.2	3.1
Less Urbanized	18.9	17.3	5.6	5.4	2.9	2.5
Completely Rural	15.4	13.8	4.4	3.1	2.0	1.1

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.79A (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	2,810 ^a	2,591	1,895	1,812	774	721
GEOGRAPHIC DIVISION						
Northeast	383	336	266	247	106	94
New England	106	88	76	65	35	23
Middle Atlantic	277	248	190	182	71	71
Midwest	612	564	434	414	189	159
East North Central	453 ^a	390	327	286	143	114
West North Central	158	174	107	128	46	45
South	1,085	975	717	651	296	266
South Atlantic	514	488	351	326	123	120
East South Central	220 ^b	161	153 ^b	103	76 ^a	48
West South Central	351	326	212	222	96	98
West	731	715	479	500	183	202
Mountain	198 ^a	252	140 ^a	189	50	86
Pacific	533	463	339	311	133	116
COUNTY TYPE						
Large Metro	1,422	1,313	931	928	377	360
Small Metro	943	853	665	594	263	248
250K – 1 Mil. Pop.	571	577	393	405	153	166
< 250K Pop.	373 ^a	276	271 ^a	189	110	82
Nonmetro	445	425	299	291	134	113
Urbanized	217	179	157	133	65	53
Less Urbanized	197	219	124	143	59	51
Completely Rural	30	27	18	16	9	9

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.79B (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	11.4 ^a	10.6	7.7	7.4	3.1	3.0
GEOGRAPHIC DIVISION						
Northeast	8.9	7.9	6.2	5.8	2.5	2.2
New England	9.6	8.1	6.9	6.0	3.2	2.1
Middle Atlantic	8.7	7.9	6.0	5.8	2.2	2.3
Midwest	11.3	10.6	8.0	7.8	3.5	3.0
East North Central	12.0 ^a	10.4	8.6	7.6	3.8	3.0
West North Central	9.8	11.0	6.6	8.0	2.8	2.8
South	12.0	10.9	8.0	7.3	3.3	3.0
South Atlantic	11.3	10.9	7.7	7.3	2.7	2.7
East South Central	15.1 ^b	11.1	10.5 ^b	7.1	5.2 ^a	3.3
West South Central	11.6	10.8	7.1	7.4	3.2	3.2
West	12.4	12.3	8.1	8.6	3.1	3.5
Mountain	11.0 ^a	14.1	7.7 ^a	10.6	2.8	4.8
Pacific	13.1	11.5	8.3	7.7	3.3	2.9
COUNTY TYPE						
Large Metro	10.9	10.1	7.1	7.1	2.9	2.8
Small Metro	12.3	11.3	8.7	7.9	3.4	3.3
250K – 1 Mil. Pop.	11.2	11.5	7.7	8.0	3.0	3.3
< 250K Pop.	14.6 ^b	11.1	10.6 ^b	7.6	4.3	3.3
Nonmetro	11.4	11.2	7.7	7.7	3.4	3.0
Urbanized	12.7	11.7	9.1	8.7	3.8	3.5
Less Urbanized	11.0	11.3	6.9	7.4	3.3	2.6
Completely Rural	7.8	8.4	4.7	5.0	2.3	2.9

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.80A (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	9,896	9,832	5,061	4,876	2,133	1,997
GEOGRAPHIC DIVISION						
Northeast	1,746	1,666	992	893	434 ^a	351
New England	523	517	306	263	147	116
Middle Atlantic	1,223	1,149	687	629	287	235
Midwest	2,198	2,107	1,090	1,107	450	472
East North Central	1,565	1,470	803	781	342	349
West North Central	634	637	287	326	107	123
South	3,512	3,664	1,806	1,763	759	712
South Atlantic	1,681	1,802	884	880	346	345
East South Central	630	605	310	277	136	126
West South Central	1,201	1,257	613	607	277	241
West	2,440	2,395	1,172	1,113	490	462
Mountain	785	749	382	347	162	154
Pacific	1,655	1,646	791	766	328	309
COUNTY TYPE						
Large Metro	4,992	5,049	2,559	2,520	1,014	983
Small Metro	3,327	3,411	1,707	1,703	767	736
250K – 1 Mil. Pop.	2,085	2,193	1,090	1,082	497	486
< 250K Pop.	1,242	1,218	617	621	270	250
Nonmetro	1,577	1,372	795	652	351	277
Urbanized	779	670	400	322	178	140
Less Urbanized	704	606	353	288	157	130
Completely Rural	95	96	42	43	17	8

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.80B (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	29.5	28.9	15.1	14.3	6.4	5.9
GEOGRAPHIC DIVISION						
Northeast	28.5	27.6	16.2	14.8	7.1	5.8
New England	32.7	31.8	19.2	16.2	9.2	7.1
Middle Atlantic	27.0	26.0	15.2	14.2	6.3	5.3
Midwest	30.0	28.1	14.9	14.8	6.1	6.3
East North Central	30.8	28.6	15.8	15.2	6.7	6.8
West North Central	28.0	27.0	12.7	13.8	4.7	5.2
South	29.2	29.5	15.0	14.2	6.3	5.7
South Atlantic	27.5	28.5	14.5	13.9	5.7	5.5
East South Central	33.1	30.4	16.3	13.9	7.1	6.4
West South Central	30.0	30.6	15.3	14.7	6.9	5.9
West	30.1	29.5	14.5	13.7	6.1	5.7
Mountain	32.4	30.0	15.8	13.9	6.7	6.1
Pacific	29.1	29.3	13.9	13.6	5.8	5.5
COUNTY TYPE						
Large Metro	28.0	27.8	14.3	13.9	5.7	5.4
Small Metro	31.7	30.7	16.3	15.3	7.3	6.6
250K – 1 Mil. Pop.	31.2	31.2	16.3	15.4	7.4	6.9
< 250K Pop.	32.7	29.7	16.2	15.2	7.1	6.1
Nonmetro	30.1	28.6	15.2	13.6	6.7	5.8
Urbanized	31.0	31.7	15.9	15.2	7.1	6.6
Less Urbanized	29.3	26.0	14.7	12.4	6.5	5.6
Completely Rural	29.1	27.6	12.8	12.2	5.1 ^a	2.2

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.81A (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	39,285	39,409	9,108	9,362	4,072	4,239
GEOGRAPHIC DIVISION						
Northeast	6,424	6,570	1,387	1,557	584	853
New England	1,727	1,741	391	430	120 ^b	272
Middle Atlantic	4,697	4,829	996	1,128	463	581
Midwest	7,829	8,007	1,765	1,746	730	753
East North Central	5,415	5,815	1,344	1,356	572	618
West North Central	2,414	2,192	421	391	158	135
South	14,128	13,708	3,437	3,188	1,621	1,546
South Atlantic	7,683	7,407	1,761	1,746	864	913
East South Central	2,141	2,475	485	494	283	230
West South Central	4,304	3,826	1,191	948	473	403
West	10,904	11,124	2,519	2,871	1,139	1,087
Mountain	3,379	3,522	867	924	423	363
Pacific	7,525	7,602	1,651	1,947	716	724
COUNTY TYPE						
Large Metro	20,929	21,943	4,807	5,263	2,198	2,328
Small Metro	12,339	11,926	2,863	2,804	1,157	1,230
250K – 1 Mil. Pop.	8,520	7,970	2,021	1,993	685	865
< 250K Pop.	3,818	3,956	843	810	472	365
Nonmetro	6,017	5,540	1,438	1,296	717	681
Urbanized	2,796	2,566	732	607	332	340
Less Urbanized	2,701	2,584	583	630	327	317
Completely Rural	521	390	123	59	58	24

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.81B (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	20.3	20.2	4.7	4.8	2.1	2.2
GEOGRAPHIC DIVISION						
Northeast	17.9	18.1	3.9	4.3	1.6	2.4
New England	18.2	18.3	4.1	4.5	1.3 ^b	2.9
Middle Atlantic	17.7	18.1	3.8	4.2	1.7	2.2
Midwest	18.5	18.8	4.2	4.1	1.7	1.8
East North Central	18.3	19.6	4.5	4.6	1.9	2.1
West North Central	18.8	17.0	3.3	3.0	1.2	1.0
South	19.9	19.1	4.8	4.5	2.3	2.2
South Atlantic	20.3	19.4	4.6	4.6	2.3	2.4
East South Central	18.4	21.2	4.2	4.2	2.4	2.0
West South Central	20.0	17.6	5.5	4.4	2.2	1.9
West	24.7	24.8	5.7	6.4	2.6	2.4
Mountain	24.9	25.6	6.4	6.7	3.1	2.6
Pacific	24.6	24.4	5.4	6.3	2.3	2.3
COUNTY TYPE						
Large Metro	20.3	21.0	4.7	5.0	2.1	2.2
Small Metro	21.1	20.3	4.9	4.8	2.0	2.1
250K – 1 Mil. Pop.	21.9	20.2	5.2	5.1	1.8	2.2
< 250K Pop.	19.3	20.4	4.3	4.2	2.4	1.9
Nonmetro	18.7	17.4	4.5	4.1	2.2	2.1
Urbanized	20.2	19.1	5.3	4.5	2.4	2.5
Less Urbanized	18.2	16.8	3.9	4.1	2.2	2.1
Completely Rural	15.0	12.7	3.5	1.9	1.7	0.8

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.82A (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	49,181	49,241	14,169	14,238	6,205	6,236
GEOGRAPHIC DIVISION						
Northeast	8,170	8,236	2,380	2,450	1,017	1,204
New England	2,250	2,258	697	693	267 ^a	388
Middle Atlantic	5,920	5,978	1,683	1,757	750	816
Midwest	10,028	10,114	2,855	2,853	1,179	1,224
East North Central	6,979	7,284	2,147	2,137	914	967
West North Central	3,048	2,829	708	717	265	258
South	17,640	17,373	5,243	4,951	2,379	2,258
South Atlantic	9,364	9,209	2,645	2,625	1,210	1,258
East South Central	2,771	3,080	795	771	419	356
West South Central	5,505	5,084	1,804	1,555	750	644
West	13,344	13,519	3,691	3,984	1,629	1,550
Mountain	4,164	4,271	1,249	1,271	584	517
Pacific	9,180	9,248	2,442	2,713	1,044	1,033
COUNTY TYPE						
Large Metro	25,921	26,992	7,366	7,784	3,213	3,311
Small Metro	15,666	15,337	4,570	4,507	1,924	1,966
250K – 1 Mil. Pop.	10,605	10,164	3,111	3,075	1,183	1,351
< 250K Pop.	5,061	5,174	1,459	1,432	742	615
Nonmetro	7,594	6,912	2,233	1,948	1,068	958
Urbanized	3,574	3,236	1,132	928	510	480
Less Urbanized	3,405	3,190	935	918	484	447
Completely Rural	615	486	165	102	74	32

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.82B (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	21.6	21.5	6.2	6.2	2.7	2.7
GEOGRAPHIC DIVISION						
Northeast	19.4	19.5	5.7	5.8	2.4	2.8
New England	20.3	20.2	6.3	6.2	2.4 ^a	3.5
Middle Atlantic	19.1	19.2	5.4	5.6	2.4	2.6
Midwest	20.2	20.2	5.7	5.7	2.4	2.4
East North Central	20.1	20.9	6.2	6.1	2.6	2.8
West North Central	20.2	18.6	4.7	4.7	1.8	1.7
South	21.2	20.7	6.3	5.9	2.9	2.7
South Atlantic	21.3	20.7	6.0	5.9	2.7	2.8
East South Central	20.5	22.6	5.9	5.7	3.1	2.6
West South Central	21.6	19.7	7.1	6.0	2.9	2.5
West	25.5	25.5	7.1	7.5	3.1	2.9
Mountain	26.0	26.3	7.8	7.8	3.6	3.2
Pacific	25.3	25.2	6.7	7.4	2.9	2.8
COUNTY TYPE						
Large Metro	21.5	22.0	6.1	6.3	2.7	2.7
Small Metro	22.7	21.9	6.6	6.4	2.8	2.8
250K – 1 Mil. Pop.	23.3	21.9	6.8	6.6	2.6	2.9
< 250K Pop.	21.5	22.0	6.2	6.1	3.1	2.6
Nonmetro	20.3	18.8	6.0	5.3	2.9	2.6
Urbanized	21.9	20.8	6.9	6.0	3.1	3.1
Less Urbanized	19.8	18.0	5.4	5.2	2.8	2.5
Completely Rural	16.2	14.3	4.3	3.0	2.0	0.9

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.