

Table 1.83A (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	35,197	34,908	12,450	12,242	5,282	5,093
GEOGRAPHIC DIVISION						
Northeast	5,851	5,656	2,088	1,987	893	904
New England	1,637	1,702	594	559	229	301
Middle Atlantic	4,215	3,954	1,494	1,428	663	604
Midwest	7,442	7,257	2,534	2,601	1,002	1,101
East North Central	5,370	5,391	1,918	1,912	779	860
West North Central	2,072	1,866	617	689	223	241
South	12,635	12,357	4,699 ^a	4,092	2,069	1,759
South Atlantic	6,548	6,231	2,346	2,088	1,044	889
East South Central	2,096	2,334	706	676	351	318
West South Central	3,992	3,792	1,647 ^a	1,327	674	552
West	9,268	9,639	3,129	3,562	1,319	1,328
Mountain	2,971	3,235	1,115	1,183	482	432
Pacific	6,297	6,404	2,014	2,380	837	896
COUNTY TYPE						
Large Metro	18,441	18,773	6,358	6,565	2,641	2,600
Small Metro	11,256	11,082	4,123	3,885	1,736	1,623
250K – 1 Mil. Pop.	7,533	7,263	2,737	2,609	1,051	1,135
< 250K Pop.	3,723	3,819	1,386	1,276	685 ^a	488
Nonmetro	5,499	5,054	1,969	1,792	905	870
Urbanized	2,703	2,376	1,017	825	446	418
Less Urbanized	2,401	2,325	852	862	401	413
Completely Rural	396	353	101	106	59	39

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.83B (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	14.0	13.8	4.9	4.8	2.1	2.0
GEOGRAPHIC DIVISION						
Northeast	12.6	12.2	4.5	4.3	1.9	1.9
New England	13.4	13.9	4.9	4.6	1.9	2.5
Middle Atlantic	12.3	11.5	4.4	4.2	1.9	1.8
Midwest	13.5	13.1	4.6	4.7	1.8	2.0
East North Central	14.0	14.0	5.0	5.0	2.0	2.2
West North Central	12.4	11.1	3.7	4.1	1.3	1.4
South	13.7	13.3	5.1 ^a	4.4	2.2	1.9
South Atlantic	13.5	12.7	4.8	4.3	2.1	1.8
East South Central	14.0	15.5	4.7	4.5	2.3	2.1
West South Central	14.0	13.1	5.8 ^a	4.6	2.4	1.9
West	15.9	16.4	5.4	6.1	2.3	2.3
Mountain	16.7	17.9	6.3	6.6	2.7	2.4
Pacific	15.6	15.7	5.0	5.8	2.1	2.2
COUNTY TYPE						
Large Metro	13.8	13.8	4.8	4.8	2.0	1.9
Small Metro	14.7	14.3	5.4	5.0	2.3	2.1
250K – 1 Mil. Pop.	14.9	14.1	5.4	5.1	2.1	2.2
< 250K Pop.	14.3	14.7	5.3	4.9	2.6 ^a	1.9
Nonmetro	13.3	12.5	4.8	4.4	2.2	2.1
Urbanized	15.0	13.9	5.6	4.8	2.5	2.4
Less Urbanized	12.6	11.8	4.5	4.4	2.1	2.1
Completely Rural	9.5	9.5	2.4	2.8	1.4	1.0

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.84A (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	2,393	2,245	1,628	1,522	658	607
GEOGRAPHIC DIVISION						
Northeast	309	290	220	216	83	86
New England	81	72	62	55	29	21
Middle Atlantic	229	217	158	161	55	65
Midwest	530	493	374	344	155	130
East North Central	392 ^a	341	281	237	119	96
West North Central	138	153	93	107	36	34
South	930	840	615	540	250	219
South Atlantic	448	412	302	266	99	94
East South Central	186 ^a	144	129 ^a	89	68 ^a	41
West South Central	296	284	183	184	83	84
West	623	621	419	422	169	172
Mountain	168 ^a	218	112 ^a	157	45	73
Pacific	455	403	307	265	124	99
COUNTY TYPE						
Large Metro	1,199	1,103	793	753	321	296
Small Metro	800	762	574	515	227	214
250K – 1 Mil. Pop.	489	515	350	359	135	150
< 250K Pop.	311	246	224 ^a	156	92	64
Nonmetro	394	379	261	253	109	97
Urbanized	189	163	141	122	53	48
Less Urbanized	179	192	104	117	48	42
Completely Rural	27	25	16	14	8	8

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.84B (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	9.7	9.2	6.6	6.3	2.7	2.5
GEOGRAPHIC DIVISION						
Northeast	7.2	6.8	5.1	5.1	1.9	2.0
New England	7.3	6.6	5.5	5.1	2.6	2.0
Middle Atlantic	7.2	6.9	5.0	5.1	1.7	2.1
Midwest	9.8	9.2	6.9	6.4	2.9	2.4
East North Central	10.3	9.1	7.4	6.3	3.1	2.6
West North Central	8.5	9.6	5.8	6.7	2.2	2.1
South	10.3	9.4	6.8	6.0	2.8	2.4
South Atlantic	9.9	9.2	6.7	5.9	2.2	2.1
East South Central	12.8 ^a	10.0	8.9 ^a	6.2	4.7 ^a	2.9
West South Central	9.8	9.4	6.1	6.1	2.7	2.8
West	10.6	10.7	7.1	7.2	2.9	3.0
Mountain	9.3 ^a	12.2	6.2 ^a	8.8	2.5	4.1
Pacific	11.2	10.0	7.5	6.6	3.0	2.4
COUNTY TYPE						
Large Metro	9.2	8.5	6.1	5.8	2.5	2.3
Small Metro	10.4	10.1	7.5	6.9	3.0	2.8
250K – 1 Mil. Pop.	9.5	10.2	6.8	7.1	2.6	3.0
< 250K Pop.	12.2 ^a	9.9	8.7 ^b	6.3	3.6	2.6
Nonmetro	10.1	10.0	6.7	6.7	2.8	2.6
Urbanized	11.0	10.7	8.2	8.0	3.1	3.1
Less Urbanized	9.9	9.9	5.8	6.0	2.7	2.2
Completely Rural	6.9	7.9	4.2	4.5	2.1	2.5

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.85A (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	8,221	8,143	4,027	3,779	1,610	1,510
GEOGRAPHIC DIVISION						
Northeast	1,468	1,369	800 ^a	662	337 ^a	255
New England	420	411	239	182	105	79
Middle Atlantic	1,048	958	561	481	232	176
Midwest	1,857	1,765	856	905	337	388
East North Central	1,325	1,244	634	641	256	285
West North Central	532	521	222	264	81	103
South	2,882	2,958	1,482 ^a	1,316	581	506
South Atlantic	1,331	1,419	707	623	260	232
East South Central	542	512	251	226	112	87
West South Central	1,009	1,028	524	466	209	187
West	2,014	2,050	889	896	355	362
Mountain	662	636	322	278	132	113
Pacific	1,352	1,414	567	618	223	249
COUNTY TYPE						
Large Metro	4,094	4,140	2,023	1,913	747	730
Small Metro	2,791	2,840	1,355	1,351	587	570
250K – 1 Mil. Pop.	1,735	1,830	859	854	372	382
< 250K Pop.	1,057	1,010	496	496	214	188
Nonmetro	1,336	1,163	650 ^a	515	277	210
Urbanized	667	578	330	254	136	110
Less Urbanized	587	501	285	221	125	93
Completely Rural	82	85	35	40	16	7

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.85B (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	24.5	23.9	12.0 ^a	11.1	4.8	4.4
GEOGRAPHIC DIVISION						
Northeast	24.0	22.6	13.1 ^a	11.0	5.5 ^a	4.2
New England	26.3	25.3	15.0 ^a	11.2	6.6	4.8
Middle Atlantic	23.2	21.7	12.4	10.9	5.1	4.0
Midwest	25.3	23.5	11.7	12.1	4.6	5.2
East North Central	26.1	24.2	12.5	12.5	5.0	5.5
West North Central	23.5	22.1	9.8	11.2	3.6	4.4
South	24.0	23.8	12.3 ^a	10.6	4.8	4.1
South Atlantic	21.8	22.5	11.6	9.9	4.3	3.7
East South Central	28.5	25.7	13.2	11.4	5.9	4.4
West South Central	25.2	25.0	13.1	11.3	5.2	4.5
West	24.9	25.3	11.0	11.0	4.4	4.5
Mountain	27.3	25.5	13.3	11.1	5.5	4.5
Pacific	23.8	25.2	10.0	11.0	3.9	4.4
COUNTY TYPE						
Large Metro	22.9	22.8	11.3	10.5	4.2	4.0
Small Metro	26.6	25.5	12.9	12.1	5.6	5.1
250K – 1 Mil. Pop.	25.9	26.0	12.8	12.2	5.6	5.4
< 250K Pop.	27.8	24.7	13.0	12.1	5.6	4.6
Nonmetro	25.5	24.3	12.4	10.8	5.3	4.4
Urbanized	26.5	27.4	13.1	12.0	5.4	5.2
Less Urbanized	24.4	21.5	11.9	9.5	5.2	4.0
Completely Rural	25.3	24.2	10.7	11.5	5.0 ^a	2.0

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.86A (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	24,583	24,521	6,795	6,941	3,014	2,975
GEOGRAPHIC DIVISION						
Northeast	4,074	3,997	1,067	1,108	473	563
New England	1,136	1,218	293	322	96 ^a	201
Middle Atlantic	2,938	2,779	774	786	377	363
Midwest	5,055	4,998	1,304	1,352	509	583
East North Central	3,653	3,806	1,002	1,033	403	479
West North Central	1,402	1,192	302	319	106	104
South	8,823	8,558	2,602	2,236	1,238	1,034
South Atlantic	4,768	4,401	1,337	1,199	685	564
East South Central	1,368	1,678	325	361	171	190
West South Central	2,687	2,480	939	677	382	281
West	6,632	6,968	1,822	2,244	795	794
Mountain	2,141	2,381	681	747	304	246
Pacific	4,490	4,587	1,140	1,497	490	549
COUNTY TYPE						
Large Metro	13,148	13,529	3,542	3,898	1,573	1,574
Small Metro	7,665	7,480	2,194	2,019	922	838
250K – 1 Mil. Pop.	5,310	4,918	1,528	1,396	544	603
< 250K Pop.	2,355	2,562	666	623	378	236
Nonmetro	3,770	3,512	1,058	1,024	519	562
Urbanized	1,848	1,635	546	449	257	261
Less Urbanized	1,635	1,633	463	524	228	277
Completely Rural	286	243	50	51	34	24

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.86B (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	12.7	12.6	3.5	3.6	1.6	1.5
GEOGRAPHIC DIVISION						
Northeast	11.3	11.0	3.0	3.1	1.3	1.6
New England	12.0	12.8	3.1	3.4	1.0 ^a	2.1
Middle Atlantic	11.1	10.4	2.9	2.9	1.4	1.4
Midwest	11.9	11.8	3.1	3.2	1.2	1.4
East North Central	12.3	12.8	3.4	3.5	1.4	1.6
West North Central	10.9	9.3	2.4	2.5	0.8	0.8
South	12.4	12.0	3.7	3.1	1.7	1.4
South Atlantic	12.6	11.5	3.5	3.1	1.8	1.5
East South Central	11.8	14.4	2.8	3.1	1.5	1.6
West South Central	12.5	11.4	4.4 ^a	3.1	1.8	1.3
West	15.0	15.5	4.1	5.0	1.8	1.8
Mountain	15.8	17.3	5.0	5.4	2.2	1.8
Pacific	14.7	14.8	3.7	4.8	1.6	1.8
COUNTY TYPE						
Large Metro	12.8	12.9	3.4	3.7	1.5	1.5
Small Metro	13.1	12.7	3.7	3.4	1.6	1.4
250K – 1 Mil. Pop.	13.7	12.5	3.9	3.5	1.4	1.5
< 250K Pop.	11.9	13.2	3.4	3.2	1.9	1.2
Nonmetro	11.7	11.0	3.3	3.2	1.6	1.8
Urbanized	13.4	12.2	4.0	3.3	1.9	1.9
Less Urbanized	11.0	10.6	3.1	3.4	1.5	1.8
Completely Rural	8.2	8.0	1.4	1.7	1.0	0.8

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.87A (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	32,804	32,664	10,822	10,720	4,624	4,485
GEOGRAPHIC DIVISION						
Northeast	5,542	5,366	1,868	1,771	809	818
New England	1,556	1,630	532	504	201	279
Middle Atlantic	3,986	3,736	1,336	1,267	609	539
Midwest	6,912	6,763	2,160	2,257	846	971
East North Central	4,978	5,051	1,637	1,674	660	764
West North Central	1,934	1,713	524	582	187	207
South	11,705	11,516	4,084 ^a	3,552	1,819	1,540
South Atlantic	6,100	5,820	2,044	1,822	945	795
East South Central	1,910	2,190	577	587	282	277
West South Central	3,695	3,507	1,463 ^a	1,143	592	468
West	8,645	9,018	2,710	3,140	1,149	1,156
Mountain	2,803	3,017	1,003	1,025	436	358
Pacific	5,842	6,001	1,707	2,115	713	798
COUNTY TYPE						
Large Metro	17,242	17,669	5,565	5,811	2,320	2,304
Small Metro	10,456	10,320	3,549	3,369	1,509	1,408
250K – 1 Mil. Pop.	7,044	6,748	2,387	2,250	916	985
< 250K Pop.	3,412	3,572	1,162	1,119	593 ^a	424
Nonmetro	5,105	4,674	1,708	1,539	796	772
Urbanized	2,514	2,213	876	704	393	371
Less Urbanized	2,222	2,133	748	745	353	371
Completely Rural	369	328	84	91	50	31

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.87B (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	14.4	14.2	4.8	4.7	2.0	2.0
GEOGRAPHIC DIVISION						
Northeast	13.2	12.7	4.4	4.2	1.9	1.9
New England	14.1	14.6	4.8	4.5	1.8	2.5
Middle Atlantic	12.9	12.0	4.3	4.1	2.0	1.7
Midwest	13.9	13.5	4.3	4.5	1.7	1.9
East North Central	14.4	14.5	4.7	4.8	1.9	2.2
West North Central	12.8	11.3	3.5	3.8	1.2	1.4
South	14.1	13.7	4.9 ^a	4.2	2.2	1.8
South Atlantic	13.9	13.1	4.6	4.1	2.1	1.8
East South Central	14.1	16.1	4.3	4.3	2.1	2.0
West South Central	14.5	13.6	5.7 ^a	4.4	2.3	1.8
West	16.5	17.0	5.2	5.9	2.2	2.2
Mountain	17.5	18.6	6.3	6.3	2.7	2.2
Pacific	16.1	16.3	4.7	5.8	2.0	2.2
COUNTY TYPE						
Large Metro	14.3	14.4	4.6	4.7	1.9	1.9
Small Metro	15.1	14.8	5.1	4.8	2.2	2.0
250K – 1 Mil. Pop.	15.5	14.5	5.2	4.8	2.0	2.1
< 250K Pop.	14.5	15.2	4.9	4.8	2.5	1.8
Nonmetro	13.7	12.7	4.6	4.2	2.1	2.1
Urbanized	15.4	14.2	5.4	4.5	2.4	2.4
Less Urbanized	12.9	12.0	4.3	4.2	2.0	2.1
Completely Rural	9.7	9.6	2.2	2.7	1.3	0.9

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.