

**Table 1.88A Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Numbers in Thousands, 2009 and 2010**

<b>Drug or Method of Administration</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
<b>HALLUCINOGENS</b>	37,256	37,476	1,012	999	6,365	6,261	29,879	30,216
PCP (Angel Dust, Phencyclidine)	6,239	6,258	99	76	468	437	5,672	5,744
LSD (Acid)	23,635	23,320	249	231	2,294	2,176	21,092	20,913
Peyote	5,196	5,776	63	42	345	331	4,788	5,403
Mescaline	8,252	8,663	26	40	356	380	7,870	8,242
Psilocybin (Mushrooms)	21,170	21,126	406	366	3,810	3,776	16,954	16,984
Ecstasy (MDMA)	14,234 <sup>b</sup>	15,901	572	619	4,168	4,216	9,494 <sup>b</sup>	11,066
<b>INHALANTS</b>	22,448	21,660	2,267 <sup>b</sup>	2,005	3,581	3,363	16,599	16,292
Amyl Nitrite, Poppers, Locker Room Odorizers, or Rush	7,407	7,176	370	336	565	600	6,472	6,240
Correction Fluid, Degreaser, or Cleaning Fluid	1,818	1,956	460	411	308	327	1,050	1,218
Gasoline or Lighter Fluid	3,743	3,385	773 <sup>b</sup>	633	521	496	2,449	2,256
Glue, Shoe Polish, or Toluene	3,558	4,019	784	770	463	481	2,311	2,768
Halothane, Ether, or Other Anesthetics	1,158	1,042	74	79	215	195	868	767
Lacquer Thinner or Other Paint Solvents	1,710	1,826	353	287	309	316	1,048	1,224
Lighter Gases (Butane, Propane)	926	831	227 <sup>a</sup>	158	162	161	538	512
Nitrous Oxide or Whippits	12,401	12,222	258	259	2,260	2,070	9,883	9,894
Spray Paints	1,998	2,098	713 <sup>b</sup>	554	381	438	904	1,106
Other Aerosol Sprays	2,381	2,147	501	423	629	579	1,252	1,145
<b>NEEDLE USE<sup>1,2,3</sup></b>	3,782	3,635	59	46	427	372	3,296	3,217
Heroin Needle Use <sup>1,2</sup>	1,499	1,757	27	24	216	227	1,256	1,506
Cocaine Needle Use <sup>1,2</sup>	2,119	2,265	16	*	152	154	1,951	2,111
Stimulant Needle Use <sup>1,2,3</sup>	1,920	1,812	26	17	184	129	1,710	1,665
Methamphetamine Needle Use <sup>1,2,3</sup>	1,589	1,557	18	11	147	104	1,424	1,441
<b>HEROIN</b>	3,683	4,126	56	46	573	607	3,054	3,473
Smoke Heroin <sup>2</sup>	1,358	1,465	30	22	226	244	1,102	1,199
Sniff or Snort Heroin <sup>2</sup>	2,516 <sup>a</sup>	3,064	27	17	442	425	2,047 <sup>a</sup>	2,622

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

<sup>2</sup> These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

<sup>3</sup> Estimates of Stimulant Needle Use, Methamphetamine Needle Use, and the Needle Use summary variable include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.88B Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Percentages, 2009 and 2010**

<b>Drug or Method of Administration</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
<b>HALLUCINOGENS</b>	14.8	14.8	4.1	4.1	19.0	18.4	15.4	15.5
PCP (Angel Dust, Phencyclidine)	2.5	2.5	0.4	0.3	1.4	1.3	2.9	2.9
LSD (Acid)	9.4	9.2	1.0	0.9	6.8	6.4	10.9	10.7
Peyote	2.1	2.3	0.3	0.2	1.0	1.0	2.5	2.8
Mescaline	3.3	3.4	0.1	0.2	1.1	1.1	4.1	4.2
Psilocybin (Mushrooms)	8.4	8.3	1.7	1.5	11.3	11.1	8.8	8.7
Ecstasy (MDMA)	5.7 <sup>b</sup>	6.3	2.3	2.5	12.4	12.4	4.9 <sup>b</sup>	5.7
<b>INHALANTS</b>	8.9	8.5	9.2 <sup>b</sup>	8.2	10.7 <sup>a</sup>	9.9	8.6	8.3
Amyl Nitrite, Poppers, Locker Room Odorizers, or Rush	2.9	2.8	1.5	1.4	1.7	1.8	3.3	3.2
Correction Fluid, Degreaser, or Cleaning Fluid	0.7	0.8	1.9	1.7	0.9	1.0	0.5	0.6
Gasoline or Lighter Fluid	1.5	1.3	3.1 <sup>a</sup>	2.6	1.6	1.5	1.3	1.2
Glue, Shoe Polish, or Toluene	1.4	1.6	3.2	3.2	1.4	1.4	1.2	1.4
Halothane, Ether, or Other Anesthetics	0.5	0.4	0.3	0.3	0.6	0.6	0.4	0.4
Lacquer Thinner or Other Paint Solvents	0.7	0.7	1.4	1.2	0.9	0.9	0.5	0.6
Lighter Gases (Butane, Propane)	0.4	0.3	0.9 <sup>a</sup>	0.6	0.5	0.5	0.3	0.3
Nitrous Oxide or Whippits	4.9	4.8	1.0	1.1	6.7 <sup>a</sup>	6.1	5.1	5.1
Spray Paints	0.8	0.8	2.9 <sup>b</sup>	2.3	1.1	1.3	0.5	0.6
Other Aerosol Sprays	0.9	0.8	2.0	1.7	1.9	1.7	0.6	0.6
<b>NEEDLE USE<sup>1,2,3</sup></b>	1.5	1.4	0.2	0.2	1.3	1.1	1.7	1.6
Heroin Needle Use <sup>1,2</sup>	0.6	0.7	0.1	0.1	0.6	0.7	0.6	0.8
Cocaine Needle Use <sup>1,2</sup>	0.8	0.9	0.1	*	0.5	0.5	1.0	1.1
Stimulant Needle Use <sup>1,2,3</sup>	0.8	0.7	0.1	0.1	0.5	0.4	0.9	0.9
Methamphetamine Needle Use <sup>1,2,3</sup>	0.6	0.6	0.1	0.0	0.4	0.3	0.7	0.7
<b>HEROIN</b>	1.5	1.6	0.2	0.2	1.7	1.8	1.6	1.8
Smoke Heroin <sup>2</sup>	0.5	0.6	0.1	0.1	0.7	0.7	0.6	0.6
Sniff or Snort Heroin <sup>2</sup>	1.0	1.2	0.1	0.1	1.3	1.2	1.1 <sup>a</sup>	1.3

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

<sup>2</sup> These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

<sup>3</sup> Estimates of Stimulant Needle Use, Methamphetamine Needle Use, and the Needle Use summary variable include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.89A Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Numbers in Thousands, 2009 and 2010**

<b>Pain Reliever</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
Darvocet <sup>®</sup> , Darvon <sup>®</sup> , or Tylenol <sup>®</sup> with Codeine	18,889	17,839	1,051	964	3,475 <sup>a</sup>	3,216	14,363	13,659
Percocet <sup>®</sup> , Percodan <sup>®</sup> , or Tylox <sup>®</sup>	13,570	13,753	469	418	3,219	3,171	9,883	10,164
Vicodin <sup>®</sup> , Lortab <sup>®</sup> , or Lorcet <sup>®</sup>	22,065	22,479	1,104	988	6,088	5,956	14,873	15,535
Codeine	7,120	6,790	349	378	1,953	1,854	4,818	4,558
Demerol <sup>®</sup>	2,219	2,394	61	54	445	434	1,712	1,907
Dilaudid <sup>®</sup>	1,092	1,095	20	14	193 <sup>b</sup>	284	879	797
Fioricet <sup>®</sup>	430	373	16	15	82	62	332	296
Fiorinal <sup>®</sup>	342	257	8	11	25	16	309	230
Hydrocodone	9,626	10,208	429	490	2,950	3,109	6,248	6,609
Methadone	2,026	2,189	115	139	688 <sup>a</sup>	821	1,223	1,228
Morphine	2,748	3,018	182	159	985	1,009	1,580	1,850
OxyContin <sup>®</sup>	5,829	6,121	378	366	2,084	2,129	3,368	3,626
Phenaphen <sup>®</sup> with Codeine	602	598	44	50	159	160	399	388
Propoxyphene	318 <sup>a</sup>	190	22	35	100	63	196 <sup>a</sup>	92
SK-65 <sup>®</sup>	67	64	6	15	18	29	43	19
Stadol <sup>®</sup>	81	94	2	3	26	18	53	73
Talacen <sup>®</sup>	77	38	9	7	7	17	60	14
Talwin <sup>®</sup>	234	278	11	10	10	12	213	255
Talwin <sup>®</sup> NX	78	63	5	6	15	22	58	35
Tramadol	1,443	1,711	90	86	567	605	786	1,020
Ultram <sup>®</sup>	1,359	1,229	54	42	362 <sup>a</sup>	282	943	905
<b>SELECTED GROUPS OF DRUGS<sup>1</sup></b>								
Propoxyphene or Codeine Products <sup>2</sup>	20,985 <sup>a</sup>	19,663	1,201	1,141	4,135	3,880	15,649	14,643
Oxycodone Products <sup>3</sup>	15,281	15,553	691	639	3,893	3,933	10,697	10,981
Hydrocodone Products <sup>4</sup>	23,543	24,093	1,238	1,148	6,512	6,479	15,794	16,467
Tramadol Products <sup>5</sup>	2,181	2,269	129	113	762	692	1,289	1,464

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Includes other-specify drug responses that are not asked about explicitly in the Pain Relievers module but fall into this category.

<sup>2</sup> Includes Darvocet<sup>®</sup>, Darvon<sup>®</sup> or Tylenol<sup>®</sup> with Codeine, codeine, Phenaphen<sup>®</sup> with Codeine, propoxyphene, and SK-65<sup>®</sup>.

<sup>3</sup> Includes Percocet<sup>®</sup>, Percodan<sup>®</sup> or Tylox<sup>®</sup>, and OxyContin<sup>®</sup>.

<sup>4</sup> Includes Vicodin<sup>®</sup>, Lortab<sup>®</sup>, or Lorcet<sup>®</sup>, and hydrocodone.

<sup>5</sup> Includes tramadol and Ultram<sup>®</sup>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.89B Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Percentages, 2009 and 2010**

<b>Pain Reliever</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
Darvocet <sup>®</sup> , Darvon <sup>®</sup> , or Tylenol <sup>®</sup> with Codeine	7.5	7.0	4.3	4.0	10.3 <sup>a</sup>	9.4	7.4	7.0
Percocet <sup>®</sup> , Percodan <sup>®</sup> , or Tylox <sup>®</sup>	5.4	5.4	1.9	1.7	9.6	9.3	5.1	5.2
Vicodin <sup>®</sup> , Lortab <sup>®</sup> , or Lorcet <sup>®</sup>	8.8	8.9	4.5	4.1	18.1	17.5	7.7	8.0
Codeine	2.8	2.7	1.4	1.6	5.8	5.4	2.5	2.3
Demerol <sup>®</sup>	0.9	0.9	0.2	0.2	1.3	1.3	0.9	1.0
Dilaudid <sup>®</sup>	0.4	0.4	0.1	0.1	0.6 <sup>a</sup>	0.8	0.5	0.4
Fioricet <sup>®</sup>	0.2	0.1	0.1	0.1	0.2	0.2	0.2	0.2
Fiorinal <sup>®</sup>	0.1	0.1	0.0	0.0	0.1	0.0	0.2	0.1
Hydrocodone	3.8	4.0	1.7	2.0	8.8	9.1	3.2	3.4
Methadone	0.8	0.9	0.5	0.6	2.0	2.4	0.6	0.6
Morphine	1.1	1.2	0.7	0.7	2.9	3.0	0.8	0.9
OxyContin <sup>®</sup>	2.3	2.4	1.5	1.5	6.2	6.2	1.7	1.9
Phenaphen <sup>®</sup> with Codeine	0.2	0.2	0.2	0.2	0.5	0.5	0.2	0.2
Propoxyphene	0.1 <sup>a</sup>	0.1	0.1	0.1	0.3	0.2	0.1 <sup>a</sup>	0.0
SK-65 <sup>®</sup>	0.0	0.0	0.0	0.1	0.1	0.1	0.0	0.0
Stadol <sup>®</sup>	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0
Talacen <sup>®</sup>	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Talwin <sup>®</sup>	0.1	0.1	0.0	0.0	0.0	0.0	0.1	0.1
Talwin <sup>®</sup> NX	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0
Tramadol	0.6	0.7	0.4	0.4	1.7	1.8	0.4	0.5
Ultram <sup>®</sup>	0.5	0.5	0.2	0.2	1.1 <sup>a</sup>	0.8	0.5	0.5
<b>SELECTED GROUPS OF DRUGS<sup>1</sup></b>								
Propoxyphene or Codeine Products <sup>2</sup>	8.3 <sup>a</sup>	7.8	4.9	4.7	12.3 <sup>a</sup>	11.4	8.1	7.5
Oxycodone Products <sup>3</sup>	6.1	6.1	2.8	2.6	11.6	11.5	5.5	5.6
Hydrocodone Products <sup>4</sup>	9.3	9.5	5.0	4.7	19.4	19.0	8.2	8.4
Tramadol Products <sup>5</sup>	0.9	0.9	0.5	0.5	2.3	2.0	0.7	0.8

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Includes other-specify drug responses that are not asked about explicitly in the Pain Relievers module but fall into this category.

<sup>2</sup> Includes Darvocet<sup>®</sup>, Darvon<sup>®</sup> or Tylenol<sup>®</sup> with Codeine, codeine, Phenaphen<sup>®</sup> with Codeine, propoxyphene, and SK-65<sup>®</sup>.

<sup>3</sup> Includes Percocet<sup>®</sup>, Percodan<sup>®</sup> or Tylox<sup>®</sup>, and OxyContin<sup>®</sup>.

<sup>4</sup> Includes Vicodin<sup>®</sup>, Lortab<sup>®</sup>, or Lorcet<sup>®</sup>, and hydrocodone.

<sup>5</sup> Includes tramadol and Ultram<sup>®</sup>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.90A Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Numbers in Thousands, 2009 and 2010**

<b>Tranquilizer</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
Klonopin® or Clonazepam	4,725	4,729	162	193	1,400	1,393	3,163	3,144
Xanax®, Alprazolam, Ativan®, or Lorazepam	12,846	13,645	487	541	3,346	3,582	9,014	9,522
Valium® or Diazepam	13,338	13,468	210	215	1,992	1,986	11,136	11,267
Atarax®	307	182	20	23	39	46	248 <sup>a</sup>	114
BuSpar®	489	532	26	13	148	154	314	365
Equanil®	73	58	5	8	12	8	56	42
Flexeril®	2,399	2,569	46	56	539	500	1,814	2,013
Librium®	874	795	14	16	47	59	812	720
Limbitrol®	87	99	14	8	34	15	40	76
Meprobamate	106	64	12	9	21	26	73	29
Miltown®	76	59	4	10	5	7	66	42
Rohypnol®	295	336	22	11	66	60	207	265
Serax®	74	64	11	9	21	25	42	30
Soma®	2,949	3,240	104	117	928	916	1,917	2,207
Tranxene®	157 <sup>a</sup>	66	16	*	20	19	121	47
Vistaril®	200	277	14	18	46	56	140	203
<b>SELECTED GROUPS OF DRUGS<sup>1</sup></b>								
Benzodiazepines <sup>2</sup>	20,449	20,708	625	642	4,086	4,261	15,739	15,805
Meprobamate Products <sup>3</sup>	228	158	19	22	27	33	182	104
Muscle Relaxants <sup>4</sup>	4,453	4,761	140	162	1,205	1,155	3,108	3,444

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Includes other-specify drug responses that are not asked about explicitly in the Tranquilizers module but fall into this category.

<sup>2</sup> Includes Klonopin® or clonazepam, Xanax®, alprazolam, Ativan® or lorazepam, Valium® or diazepam, Librium®, Limbitrol®, Rohypnol®, Serax®, and Tranxene®.

<sup>3</sup> Includes Equanil®, meprobamate, and Miltown®.

<sup>4</sup> Includes Flexeril® and Soma®.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.90B Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Percentages, 2009 and 2010**

<b>Tranquilizer</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
Klonopin <sup>®</sup> or Clonazepam	1.9	1.9	0.7	0.8	4.2	4.1	1.6	1.6
Xanax <sup>®</sup> , Alprazolam, Ativan <sup>®</sup> , or Lorazepam	5.1	5.4	2.0	2.2	10.0	10.5	4.7	4.9
Valium <sup>®</sup> or Diazepam	5.3	5.3	0.9	0.9	5.9	5.8	5.8	5.8
Atarax <sup>®</sup>	0.1	0.1	0.1	0.1	0.1	0.1	0.1 <sup>a</sup>	0.1
BuSpar <sup>®</sup>	0.2	0.2	0.1	0.1	0.4	0.5	0.2	0.2
Equanil <sup>®</sup>	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Flexeril <sup>®</sup>	1.0	1.0	0.2	0.2	1.6	1.5	0.9	1.0
Librium <sup>®</sup>	0.3	0.3	0.1	0.1	0.1	0.2	0.4	0.4
Limbitrol <sup>®</sup>	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.0
Meprobamate	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0
Miltown <sup>®</sup>	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Rohypnol <sup>®</sup>	0.1	0.1	0.1	0.0	0.2	0.2	0.1	0.1
Serax <sup>®</sup>	0.0	0.0	0.0	0.0	0.1	0.1	0.0	0.0
Soma <sup>®</sup>	1.2	1.3	0.4	0.5	2.8	2.7	1.0	1.1
Tranxene <sup>®</sup>	0.1 <sup>a</sup>	0.0	0.1	*	0.1	0.1	0.1	0.0
Vistaril <sup>®</sup>	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1
<b>SELECTED GROUPS OF DRUGS<sup>1</sup></b>								
Benzodiazepines <sup>2</sup>	8.1	8.2	2.5	2.6	12.2	12.5	8.1	8.1
Meprobamate Products <sup>3</sup>	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Muscle Relaxants <sup>4</sup>	1.8	1.9	0.6	0.7	3.6	3.4	1.6	1.8

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Includes other-specify drug responses that are not asked about explicitly in the Tranquilizers module but fall into this category.

<sup>2</sup> Includes Klonopin<sup>®</sup> or clonazepam, Xanax<sup>®</sup>, alprazolam, Ativan<sup>®</sup> or lorazepam, Valium<sup>®</sup> or diazepam, Librium<sup>®</sup>, Limbitrol<sup>®</sup>, Rohypnol<sup>®</sup>, Serax<sup>®</sup>, and Tranxene<sup>®</sup>.

<sup>3</sup> Includes Equanil<sup>®</sup>, meprobamate, and Miltown<sup>®</sup>.

<sup>4</sup> Includes Flexeril<sup>®</sup> and Soma<sup>®</sup>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.91A Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Numbers in Thousands, 2009 and 2010**

<b>Stimulant</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
Methamphetamine, Desoxyn <sup>®</sup> , or Methedrine <sup>1</sup>	12,837	13,012	185	157	1,511	1,353	11,141	11,503
Prescription Diet Pills <sup>2</sup>	7,861	7,342	158	127	693	694	7,010	6,522
Ritalin <sup>®</sup> or Methylphenidate	5,363	5,422	269	250	1,754	1,818	3,340	3,354
Cylert <sup>®</sup>	181	105	13	9	33	41	135	55
Dexedrine <sup>®</sup>	2,357	1,829	21	14	164	145	2,172	1,670
Dextroamphetamine	668	791	28	27	183	190	457	573
Didrex <sup>®</sup>	203	95	9	7	22	18	171	69
Eskatrol <sup>®</sup>	150	97	9	6	13	26	128	64
Ionamin <sup>®</sup>	362	236	15	5	8	6	339	225
Mazanor <sup>®</sup>	21	26	5	3	6	13	10	10
Obedrin-LA <sup>®</sup>	*	*	6	*	4	3	*	*
Plegine <sup>®</sup>	20	*	6	4	8	8	*	*
Preludin <sup>®</sup>	392	431	9	5	16	26	367	400
Sanorex <sup>®</sup>	67	50	9	8	19 <sup>a</sup>	5	38	37
Tenuate <sup>®</sup>	118	63	6	8	11	4	100	50
<b>SELECTED GROUPS OF DRUGS<sup>3</sup></b>								
Amphetamine, Dextroamphetamine, and Phentermine Products <sup>4</sup>	9,794	9,525	239	201	1,424	1,535	8,131	7,789
Mazindol Products <sup>5</sup>	84	73	13	9	22	17	48	47
Methylphenidate or Dexmethylphenidate Products <sup>6</sup>	5,376	5,445	270	258	1,766	1,830	3,340	3,357

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Also known as crank, crystal, ice, or speed. Estimates of Use of Methamphetamine, Desoxyn<sup>®</sup>, or Methedrine include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Respondents were asked about their use of prescription diet pills and were given the following as examples: amphetamines, Benzedrine<sup>®</sup>, Biphetamine<sup>®</sup>, Fastin<sup>®</sup>, or phentermine. However, respondents were not given an exhaustive list of examples of prescription diet pills.

<sup>3</sup> Includes other-specify drug responses that are not asked about explicitly in the Stimulants module but fall into this category.

<sup>4</sup> Includes prescription diet pills, Dexedrine<sup>®</sup>, dextroamphetamine, and Ionamin<sup>®</sup>.

<sup>5</sup> Includes Mazanor<sup>®</sup> and Sanorex<sup>®</sup>.

<sup>6</sup> Includes Ritalin<sup>®</sup>, other products containing methylphenidate, and products containing dexmethylphenidate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.91B Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Percentages, 2009 and 2010**

<b>Stimulant</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
Methamphetamine, Desoxyn <sup>®</sup> , or Methedrine <sup>1</sup>	5.1	5.1	0.8	0.6	4.5 <sup>a</sup>	4.0	5.8	5.9
Prescription Diet Pills <sup>2</sup>	3.1	2.9	0.6	0.5	2.1	2.0	3.6	3.3
Ritalin <sup>®</sup> or Methylphenidate	2.1	2.1	1.1	1.0	5.2	5.3	1.7	1.7
Cylert <sup>®</sup>	0.1	0.0	0.1	0.0	0.1	0.1	0.1	0.0
Dexedrine <sup>®</sup>	0.9	0.7	0.1	0.1	0.5	0.4	1.1	0.9
Dextroamphetamine	0.3	0.3	0.1	0.1	0.5	0.6	0.2	0.3
Didrex <sup>®</sup>	0.1	0.0	0.0	0.0	0.1	0.1	0.1	0.0
Eskatrol <sup>®</sup>	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.0
Ionamin <sup>®</sup>	0.1	0.1	0.1	0.0	0.0	0.0	0.2	0.1
Mazanor <sup>®</sup>	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Obedrin-LA <sup>®</sup>	*	*	0.0	*	0.0	0.0	*	*
Plegine <sup>®</sup>	0.0	*	0.0	0.0	0.0	0.0	*	*
Preludin <sup>®</sup>	0.2	0.2	0.0	0.0	0.0	0.1	0.2	0.2
Sanorex <sup>®</sup>	0.0	0.0	0.0	0.0	0.1 <sup>a</sup>	0.0	0.0	0.0
Tenuate <sup>®</sup>	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0
<b>SELECTED GROUPS OF DRUGS<sup>3</sup></b>								
Amphetamine, Dextroamphetamine, and Phentermine Products <sup>4</sup>	3.9	3.8	1.0	0.8	4.2	4.5	4.2	4.0
Mazindol Products <sup>5</sup>	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.0
Methylphenidate or Dexmethylphenidate Products <sup>6</sup>	2.1	2.1	1.1	1.1	5.3	5.4	1.7	1.7

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Also known as crank, crystal, ice, or speed. Estimates of Use of Methamphetamine, Desoxyn<sup>®</sup>, or Methedrine include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

<sup>2</sup> Respondents were asked about their use of prescription diet pills and were given the following as examples: amphetamines, Benzedrine<sup>®</sup>, Biphetamine<sup>®</sup>, Fastin<sup>®</sup>, or phentermine. However, respondents were not given an exhaustive list of examples of prescription diet pills.

<sup>3</sup> Includes other-specify drug responses that are not asked about explicitly in the Stimulants module but fall into this category.

<sup>4</sup> Includes prescription diet pills, Dexedrine<sup>®</sup>, dextroamphetamine, and Ionamin<sup>®</sup>.

<sup>5</sup> Includes Mazanor<sup>®</sup> and Sanorex<sup>®</sup>.

<sup>6</sup> Includes Ritalin<sup>®</sup>, other products containing methylphenidate, and products containing dexmethylphenidate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.92A Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Numbers in Thousands, 2009 and 2010**

<b>Sedative</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
Methaqualone, Sopor <sup>®</sup> , or Quaalude <sup>®</sup>	5,728	5,059	25	41	154	118	5,549	4,899
Barbiturates <sup>1</sup>	2,449	2,023	27	20	85	76	2,337	1,927
Restoril <sup>®</sup> or Temazepam	1,187	1,033	46	43	147	135	994	854
Amytal <sup>®</sup>	210	231	9	5	24	30	177	196
Butisol <sup>®</sup>	84	104	6	17	7	15	71	71
Chloral Hydrate	153	132	2	6	19	10	132	116
Dalmane <sup>®</sup>	281	219	16	8	18	19	247	193
Halcion <sup>®</sup>	526	571	15	23	25	39	486	510
Phenobarbital	982	856	16	24	62	48	904	785
Placidyl <sup>®</sup>	493	406	2	5	5	8	486	394
Tuinal <sup>®</sup>	729	516	10	2	6	9	713	505
<b>SELECTED GROUPS OF DRUGS<sup>2</sup></b>								
Temazepam, Flurazepam, or Triazolam <sup>3</sup>	1,667	1,558	70	68	164	158	1,433	1,332
Any Barbiturates <sup>4</sup>	3,036	2,642	54	55	149	120	2,833	2,466

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Respondents were asked about their use of barbiturates and were given the following as examples: Nembutal<sup>®</sup>, pentobarbital, Seconal<sup>®</sup>, secobarbital, or butalbital. However, respondents were not given an exhaustive list of examples of barbiturates.

<sup>2</sup> Includes other-specify drug responses that are not asked about explicitly in the Sedatives module but fall into this category.

<sup>3</sup> Includes Restoril<sup>®</sup>, temazepam, Dalmane<sup>®</sup>, or Halcion<sup>®</sup>.

<sup>4</sup> Includes barbiturates, Amytal<sup>®</sup>, Butisol<sup>®</sup>, phenobarbital, and Tuinal<sup>®</sup>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.92B Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Percentages, 2009 and 2010**

<b>Sedative</b>	<b>Total (2009)</b>	<b>Total (2010)</b>	<b>Aged 12-17 (2009)</b>	<b>Aged 12-17 (2010)</b>	<b>Aged 18-25 (2009)</b>	<b>Aged 18-25 (2010)</b>	<b>Aged 26+ (2009)</b>	<b>Aged 26+ (2010)</b>
Methaqualone, Sopor <sup>®</sup> , or Quaalude <sup>®</sup>	2.3	2.0	0.1	0.2	0.5	0.3	2.9	2.5
Barbiturates <sup>1</sup>	1.0	0.8	0.1	0.1	0.3	0.2	1.2	1.0
Restoril <sup>®</sup> or Temazepam	0.5	0.4	0.2	0.2	0.4	0.4	0.5	0.4
Amytal <sup>®</sup>	0.1	0.1	0.0	0.0	0.1	0.1	0.1	0.1
Butisol <sup>®</sup>	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0
Chloral Hydrate	0.1	0.1	0.0	0.0	0.1	0.0	0.1	0.1
Dalmane <sup>®</sup>	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.1
Halcion <sup>®</sup>	0.2	0.2	0.1	0.1	0.1	0.1	0.3	0.3
Phenobarbital	0.4	0.3	0.1	0.1	0.2	0.1	0.5	0.4
Placidyl <sup>®</sup>	0.2	0.2	0.0	0.0	0.0	0.0	0.3	0.2
Tuinal <sup>®</sup>	0.3	0.2	0.0	0.0	0.0	0.0	0.4	0.3
<b>SELECTED GROUPS OF DRUGS<sup>2</sup></b>								
Temazepam, Flurazepam, or Triazolam <sup>3</sup>	0.7	0.6	0.3	0.3	0.5	0.5	0.7	0.7
Any Barbiturates <sup>4</sup>	1.2	1.0	0.2	0.2	0.4	0.4	1.5	1.3

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Respondents were asked about their use of barbiturates and were given the following as examples: Nembutal<sup>®</sup>, pentobarbital, Seconal<sup>®</sup>, secobarbital, or butalbital. However, respondents were not given an exhaustive list of examples of barbiturates.

<sup>2</sup> Includes other-specify drug responses that are not asked about explicitly in the Sedatives module but fall into this category.

<sup>3</sup> Includes Restoril<sup>®</sup>, temazepam, Dalmane<sup>®</sup>, or Halcion<sup>®</sup>.

<sup>4</sup> Includes barbiturates, Amytal<sup>®</sup>, Butisol<sup>®</sup>, phenobarbital, and Tuinal<sup>®</sup>.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.