

Table 1.49C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	738	785	367	398	250	268
AGE						
12-17	69	68	58	58	36	38
18-25	152	143	118	114	73	72
26 or Older	702	752	329	364	229	252
GENDER						
Male	504	576	271	314	184	200
Female	506	501	229	242	160	167
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	674	730	339	365	236	253
White	627	650	318	335	216	234
Black or African American	181	210	86	107	66	75
American Indian or Alaska Native	83	41	38	26	35	18
Native Hawaiian or Other Pacific Islander	*	19	*	12	*	11
Asian	109	144	50	107	39	30
Two or More Races	71	104	39	29	33	17
Hispanic or Latino	240	250	147	150	92	89

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.49D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.29	0.31	0.15	0.16	0.10	0.11
AGE						
12-17	0.28	0.28	0.24	0.24	0.15	0.16
18-25	0.45	0.42	0.35	0.33	0.22	0.21
26 or Older	0.36	0.39	0.17	0.19	0.12	0.13
GENDER						
Male	0.41	0.47	0.22	0.25	0.15	0.16
Female	0.39	0.38	0.18	0.19	0.12	0.13
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.31	0.34	0.16	0.17	0.11	0.12
White	0.37	0.38	0.19	0.20	0.13	0.14
Black or African American	0.60	0.69	0.29	0.35	0.22	0.25
American Indian or Alaska Native	4.85	3.05	2.43	2.12	2.50	1.48
Native Hawaiian or Other Pacific Islander	*	2.56	*	1.69	*	1.52
Asian	0.97	1.19	0.46	0.88	0.35	0.27
Two or More Races	2.12	2.71	1.28	0.97	1.11	0.54
Hispanic or Latino	0.67	0.68	0.41	0.41	0.26	0.24

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.49P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.6509		0.8698		0.9220	
AGE						
12-17	0.0460		0.5114		0.4574	
18-25	0.3790		0.1416		0.1456	
26 or Older	0.8522		0.7014		0.6778	
GENDER						
Male	0.4501		0.8332		0.4901	
Female	0.1365		0.5985		0.5095	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.7775		0.9440		0.7644	
White	0.3388		0.6098		0.9548	
Black or African American	0.0977		0.3956		0.9918	
American Indian or Alaska Native	0.3978		0.9923		0.5938	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.1877		0.2642		0.3341	
Two or More Races	0.5561		0.7949		0.8086	
Hispanic or Latino	0.6063		0.8078		0.5793	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.50C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	69	68	58	58	36	38
GENDER						
Male	50	48	41	41	24	24
Female	48	48	41	42	27	30
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	60	58	50	50	32	33
White	56	57	48	49	28	29
Black or African American	32	28	24	20	16	13
American Indian or Alaska Native	4	5	4	4	1	2
Native Hawaiian or Other Pacific Islander	*	2	*	*	*	1
Asian	15	13	11	10	6	8
Two or More Races	8	12	6	11	4	8
Hispanic or Latino	38	36	29	32	19	20
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	38	36	32	31	18	19
Female, White, Not Hispanic	39	41	33	35	20	22
Male, Black, Not Hispanic	22	20	18	13	11	7
Female, Black, Not Hispanic	21	19	16	16	12	11
Male, Hispanic	28	29	20	25	12	14
Female, Hispanic	29	25	22	20	15	15

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.50D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.28	0.28	0.24	0.24	0.15	0.16
GENDER						
Male	0.40	0.39	0.33	0.33	0.19	0.19
Female	0.39	0.40	0.34	0.35	0.23	0.25
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.30	0.30	0.25	0.26	0.16	0.17
White	0.35	0.35	0.30	0.31	0.18	0.19
Black or African American	0.75	0.70	0.59	0.51	0.42	0.35
American Indian or Alaska Native	3.14	3.39	2.63	2.91	1.13	1.29
Native Hawaiian or Other Pacific Islander	*	3.51	*	*	*	1.01
Asian	1.41	1.14	1.05	0.88	0.61	0.76
Two or More Races	1.55	1.96	1.11	1.83	0.76	1.52
Hispanic or Latino	0.79	0.74	0.60	0.65	0.40	0.41
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.47	0.45	0.40	0.41	0.24	0.25
Female, White, Not Hispanic	0.51	0.53	0.44	0.47	0.27	0.32
Male, Black, Not Hispanic	1.05	1.03	0.88	0.67	0.56	0.40
Female, Black, Not Hispanic	1.05	0.99	0.81	0.85	0.65	0.59
Male, Hispanic	1.10	1.01	0.80	0.92	0.50	0.54
Female, Hispanic	1.09	0.97	0.86	0.80	0.60	0.62

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.50P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.0460		0.5114		0.4574	
GENDER						
Male	0.0480		0.3148		0.0694	
Female	0.4565		0.9247		0.5772	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0340		0.2953		0.3329	
White	0.1229		0.5192		0.4917	
Black or African American	0.1824		0.1253		0.1236	
American Indian or Alaska Native	0.9901		0.9264		0.8629	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.1697		0.4027		0.9118	
Two or More Races	0.0978		0.0192		0.0666	
Hispanic or Latino	0.7041		0.6548		0.8666	
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	0.0461		0.1981		0.1096	
Female, White, Not Hispanic	0.8425		0.7427		0.6262	
Male, Black, Not Hispanic	0.0678		0.0442		0.1685	
Female, Black, Not Hispanic	0.9969		0.8907		0.3787	
Male, Hispanic	0.7062		0.4162		0.7183	
Female, Hispanic	0.3310		0.8528		0.5776	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.51C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	152	143	118	114	73	72
GENDER						
Male	108	105	85	87	56	55
Female	98	94	71	71	45	47
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	136	130	107	104	63	64
White	180	165	123	117	67	64
Black or African American	48	50	31	34	18	23
American Indian or Alaska Native	13	10	10	7	10	7
Native Hawaiian or Other Pacific Islander	*	*	*	5	*	*
Asian	21	25	16	18	9	10
Two or More Races	17	21	10	13	5	5
Hispanic or Latino	57	63	48	45	36	28
EDUCATION						
< High School	68	67	47	46	34	32
High School Graduate	104	100	73	66	45	45
Some College	119	114	81	81	44	44
College Graduate	72	70	48	52	27	26
CURRENT EMPLOYMENT						
Full-Time	106	112	71	81	44	52
Part-Time	92	96	69	64	43	35
Unemployed	66	61	46	42	29	32
Other ¹	93	80	67	56	35	33

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.51D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.45	0.42	0.35	0.33	0.22	0.21
GENDER						
Male	0.64	0.61	0.51	0.51	0.33	0.32
Female	0.59	0.56	0.43	0.42	0.27	0.28
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.50	0.47	0.39	0.38	0.23	0.23
White	0.58	0.55	0.47	0.45	0.29	0.28
Black or African American	0.89	0.90	0.59	0.68	0.36	0.48
American Indian or Alaska Native	4.35	4.35	3.97	3.58	3.96	3.36
Native Hawaiian or Other Pacific Islander	*	*	*	3.12	*	*
Asian	1.28	1.62	1.02	1.17	0.61	0.72
Two or More Races	3.04	3.31	2.15	2.30	1.17	0.99
Hispanic or Latino	0.94	0.96	0.79	0.68	0.58	0.43
EDUCATION						
< High School	0.92	0.95	0.71	0.73	0.53	0.53
High School Graduate	0.68	0.69	0.53	0.51	0.36	0.36
Some College	0.77	0.71	0.60	0.58	0.37	0.35
College Graduate	1.21	1.12	0.90	0.98	0.55	0.52
CURRENT EMPLOYMENT						
Full-Time	0.70	0.68	0.52	0.58	0.34	0.39
Part-Time	0.80	0.81	0.65	0.59	0.43	0.36
Unemployed	1.13	1.15	0.87	0.86	0.60	0.67
Other ¹	0.92	0.83	0.75	0.63	0.43	0.41

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.51P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.3790		0.1416		0.1456	
GENDER						
Male	0.5968		0.1475		0.0831	
Female	0.4150		0.4981		0.8517	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.6113		0.4131		0.5236	
White	0.5619		0.4654		0.5419	
Black or African American	0.8174		0.7522		0.6454	
American Indian or Alaska Native	0.4730		0.3713		0.6387	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.2417		0.8804		0.7133	
Two or More Races	0.8352		0.6943		0.5696	
Hispanic or Latino	0.4883		0.1347		0.0760	
EDUCATION						
< High School	0.0255		0.1024		0.2671	
High School Graduate	0.0717		0.0452		0.6319	
Some College	0.5355		0.7620		0.6240	
College Graduate	0.1105		0.2861		0.3365	
CURRENT EMPLOYMENT						
Full-Time	0.9824		0.7757		0.9297	
Part-Time	0.4002		0.8459		0.0175	
Unemployed	0.0577		0.0278		0.8878	
Other ¹	0.1480		0.3505		0.5245	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.52C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	702	752	329	364	229	252
GENDER						
Male	485	559	245	290	170	189
Female	481	479	210	222	147	156
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	647	704	304	336	220	240
White	716	752	294	319	204	224
Black or African American	181	222	81	103	64	70
American Indian or Alaska Native	*	38	*	24	*	16
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	106	138	46	105	37	27
Two or More Races	69	103	37	23	33	13
Hispanic or Latino	226	231	133	135	80	81
EDUCATION						
< High School	255	265	149	135	102	84
High School Graduate	388	418	171	188	113	116
Some College	397	443	166	194	121	133
College Graduate	487	466	191	212	123	146
CURRENT EMPLOYMENT						
Full-Time	584	623	253	271	159	188
Part-Time	260	273	121	114	81	74
Unemployed	198	205	95	131	60	69
Other ¹	362	390	163	183	133	110

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.52D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.36	0.39	0.17	0.19	0.12	0.13
GENDER						
Male	0.52	0.60	0.26	0.31	0.18	0.20
Female	0.48	0.47	0.21	0.22	0.15	0.15
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.38	0.41	0.18	0.20	0.13	0.14
White	0.45	0.46	0.21	0.23	0.15	0.16
Black or African American	0.80	0.94	0.37	0.47	0.29	0.32
American Indian or Alaska Native	*	3.97	*	2.64	*	1.84
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.22	1.47	0.54	1.11	0.44	0.31
Two or More Races	3.05	3.97	1.81	1.11	1.61	0.63
Hispanic or Latino	0.90	0.91	0.53	0.53	0.32	0.32
EDUCATION						
< High School	0.85	0.89	0.52	0.47	0.36	0.29
High School Graduate	0.58	0.64	0.28	0.32	0.19	0.20
Some College	0.75	0.82	0.35	0.39	0.26	0.28
College Graduate	0.69	0.67	0.31	0.34	0.20	0.24
CURRENT EMPLOYMENT						
Full-Time	0.49	0.52	0.24	0.26	0.15	0.19
Part-Time	1.04	1.03	0.54	0.48	0.36	0.31
Unemployed	1.65	1.64	0.92	1.18	0.58	0.65
Other ¹	0.61	0.62	0.27	0.30	0.22	0.18

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.52P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.8522		0.7014		0.6778	
GENDER						
Male	0.3136		0.4314		1.0000	
Female	0.1760		0.7279		0.5008	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.9464		0.6980		0.9922	
White	0.4184		0.8399		0.7425	
Black or African American	0.0542		0.1665		0.8383	
American Indian or Alaska Native	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.1828		0.2085		0.3798	
Two or More Races	0.7374		0.5429		0.5159	
Hispanic or Latino	0.7043		0.9468		0.2040	
EDUCATION						
< High School	0.5451		0.4682		0.4135	
High School Graduate	0.4230		0.6824		0.8135	
Some College	0.3664		0.1822		0.7242	
College Graduate	0.8929		0.7070		0.2384	
CURRENT EMPLOYMENT						
Full-Time	0.6457		0.3477		0.1193	
Part-Time	0.7982		0.3512		0.7779	
Unemployed	0.3591		0.7923		0.9318	
Other ¹	0.5423		0.9358		0.2771	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.53C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	731	777	360	390	246	265
GENDER						
Male	501	572	268	308	182	199
Female	501	495	223	236	157	164
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	671	726	333	359	233	251
White	757	789	327	346	218	236
Black or African American	190	227	86	106	66	73
American Indian or Alaska Native	82	40	37	25	35	18
Native Hawaiian or Other Pacific Islander	*	19	*	12	*	*
Asian	108	144	48	106	39	29
Two or More Races	71	105	38	26	33	14
Hispanic or Latino	234	246	145	145	88	87
EDUCATION						
< High School	262	274	156	144	108	90
High School Graduate	413	437	191	200	122	125
Some College	415	461	185	212	131	142
College Graduate	499	478	199	219	126	148
CURRENT EMPLOYMENT						
Full-Time	595	641	262	284	165	197
Part-Time	278	287	137	125	90	79
Unemployed	205	219	106	141	67	77
Other ¹	378	400	178	192	137	116

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.53D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.32	0.34	0.16	0.17	0.11	0.12
GENDER						
Male	0.46	0.52	0.24	0.28	0.17	0.18
Female	0.43	0.42	0.19	0.20	0.13	0.14
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.34	0.37	0.17	0.18	0.12	0.13
White	0.40	0.42	0.20	0.21	0.14	0.15
Black or African American	0.68	0.78	0.32	0.40	0.24	0.28
American Indian or Alaska Native	5.35	3.43	2.71	2.33	2.79	1.66
Native Hawaiian or Other Pacific Islander	*	2.81	*	1.86	*	*
Asian	1.05	1.31	0.49	0.97	0.38	0.28
Two or More Races	2.57	3.23	1.54	1.07	1.34	0.55
Hispanic or Latino	0.75	0.77	0.47	0.45	0.28	0.27
EDUCATION						
< High School	0.72	0.77	0.45	0.42	0.31	0.26
High School Graduate	0.51	0.55	0.26	0.28	0.17	0.18
Some College	0.62	0.67	0.31	0.34	0.22	0.24
College Graduate	0.64	0.63	0.30	0.33	0.19	0.22
CURRENT EMPLOYMENT						
Full-Time	0.44	0.47	0.22	0.25	0.14	0.17
Part-Time	0.78	0.76	0.43	0.38	0.28	0.24
Unemployed	1.16	1.21	0.71	0.87	0.45	0.51
Other ¹	0.56	0.57	0.27	0.28	0.21	0.17

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.53P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Demographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.7397		0.9553		0.9902	
GENDER						
Male	0.3699		0.7202		0.6371	
Female	0.1411		0.5903		0.5629	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.8634		0.9293		0.8634	
White	0.3706		0.6705		0.8891	
Black or African American	0.0646		0.2169		0.7327	
American Indian or Alaska Native	0.3979		1.0000		0.5864	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.1400		0.2124		0.3321	
Two or More Races	0.7458		0.7673		0.4874	
Hispanic or Latino	0.6394		0.7344		0.5863	
EDUCATION						
< High School	0.2540		0.2011		0.2265	
High School Graduate	0.2477		0.3156		0.7205	
Some College	0.3019		0.2138		0.8120	
College Graduate	0.9384		0.5525		0.3271	
CURRENT EMPLOYMENT						
Full-Time	0.6612		0.3933		0.1268	
Part-Time	0.6402		0.2900		0.1725	
Unemployed	0.1507		0.6075		0.9732	
Other ¹	0.6920		0.7975		0.2427	

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

¹ The Other Employment category includes retired persons, disabled persons, homemakers, students, or other persons not in the labor force.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.