

**Table 1.69C (REVISED) Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	655	682	209	205	136	108
GEOGRAPHIC DIVISION						
Northeast	299	283	95	100	62	51
New England	154	134	53	44	42	29
Middle Atlantic	256	248	79	90	45	42
Midwest	226	242	72	64	45	34
East North Central	179	200	56	58	39	31
West North Central	144	135	43	27	23	14
South	362	411	126	110	98	72
South Atlantic	291	317	105	89	83	58
East South Central	125	126	28	37	19	23
West South Central	192	215	63	53	47	37
West	390	413	112	120	56	57
Mountain	187	177	58	67	34	16
Pacific	338	375	98	99	46	55
COUNTY TYPE						
Large Metro	584	596	152	176	92	92
Small Metro	426	443	125	95	92	59
250K – 1 Mil. Pop.	362	354	111	76	89	55
< 250K Pop.	241	266	59	57	24	23
Nonmetro	244	325	67	55	43	21
Urbanized	172	211	39	33	9	18
Less Urbanized	165	261	52	43	42	12
Completely Rural	61	57	17	9	1	1

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.69D (REVISED) Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.26	0.27	0.08	0.08	0.05	0.04
GEOGRAPHIC DIVISION						
Northeast	0.65	0.61	0.21	0.21	0.13	0.11
New England	1.26	1.09	0.44	0.36	0.35	0.24
Middle Atlantic	0.75	0.72	0.23	0.26	0.13	0.12
Midwest	0.41	0.44	0.13	0.12	0.08	0.06
East North Central	0.46	0.52	0.15	0.15	0.10	0.08
West North Central	0.86	0.80	0.26	0.16	0.14	0.08
South	0.39	0.44	0.14	0.12	0.11	0.08
South Atlantic	0.60	0.65	0.22	0.18	0.17	0.12
East South Central	0.84	0.83	0.18	0.24	0.13	0.16
West South Central	0.67	0.74	0.22	0.18	0.16	0.13
West	0.67	0.70	0.19	0.20	0.10	0.10
Mountain	1.05	0.98	0.33	0.37	0.19	0.09
Pacific	0.84	0.92	0.24	0.24	0.11	0.13
COUNTY TYPE						
Large Metro	0.38	0.38	0.11	0.13	0.07	0.07
Small Metro	0.47	0.46	0.16	0.12	0.12	0.08
250K – 1 Mil. Pop.	0.58	0.55	0.21	0.15	0.17	0.10
< 250K Pop.	0.84	0.81	0.22	0.21	0.09	0.09
Nonmetro	0.51	0.69	0.16	0.13	0.10	0.05
Urbanized	0.82	0.96	0.21	0.18	0.05	0.10
Less Urbanized	0.76	1.11	0.26	0.21	0.22	0.06
Completely Rural	1.08	1.34	0.38	0.23	0.04	0.02

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.69P (REVISED) Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.6992		0.2939		0.3028	
GEOGRAPHIC DIVISION						
Northeast	0.8603		0.3292		0.5644	
New England	0.0353		0.2992		0.5819	
Middle Atlantic	0.2989		0.6102		0.7808	
Midwest	0.5044		0.3863		0.3019	
East North Central	0.7295		0.6860		0.4617	
West North Central	0.5127		0.3162		0.4204	
South	0.2477		0.6122		0.3148	
South Atlantic	0.0680		0.8362		0.4235	
East South Central	0.3291		0.3044		0.5799	
West South Central	0.9588		0.1682		0.3147	
West	0.9418		0.9781		0.5580	
Mountain	0.4567		0.6495		0.1110	
Pacific	0.7693		0.8197		0.1449	
COUNTY TYPE						
Large Metro	0.5622		0.7771		0.9351	
Small Metro	0.6910		0.1143		0.1727	
250K – 1 Mil. Pop.	0.6533		0.1130		0.1617	
< 250K Pop.	0.9530		0.6484		0.9137	
Nonmetro	0.7454		0.1252		0.3601	
Urbanized	0.4533		0.2101		0.1989	
Less Urbanized	0.9442		0.4002		0.1185	
Completely Rural	0.4791		0.4071		0.3246	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.70C (REVISED) Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	28	26	22	22	13	11
GEOGRAPHIC DIVISION						
Northeast	9	8	7	7	5	4
New England	4	5	4	3	2	1
Middle Atlantic	8	7	7	6	4	4
Midwest	9	7	8	6	4	3
East North Central	8	6	7	4	3	2
West North Central	5	5	4	5	3	2
South	16	17	12	14	5	8
South Atlantic	10	9	7	8	3	5
East South Central	5	6	5	5	2	4
West South Central	11	12	8	10	4	5
West	18	18	15	14	10	6
Mountain	8	6	6	4	3	1
Pacific	16	17	13	13	9	6
COUNTY TYPE						
Large Metro	21	21	17	18	9	8
Small Metro	17	14	13	9	7	4
250K – 1 Mil. Pop.	13	12	10	8	6	4
< 250K Pop.	10	7	8	4	5	2
Nonmetro	9	10	6	10	4	7
Urbanized	6	6	5	6	2	5
Less Urbanized	6	8	4	7	3	4
Completely Rural	1	2	1	1	1	*

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.70D (REVISED) Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.11	0.11	0.09	0.09	0.05	0.05
GEOGRAPHIC DIVISION						
Northeast	0.22	0.20	0.17	0.17	0.11	0.09
New England	0.37	0.47	0.32	0.30	0.14	0.07
Middle Atlantic	0.26	0.21	0.21	0.20	0.14	0.11
Midwest	0.17	0.14	0.14	0.11	0.08	0.05
East North Central	0.20	0.15	0.17	0.12	0.09	0.05
West North Central	0.28	0.33	0.23	0.29	0.16	0.13
South	0.18	0.18	0.13	0.16	0.06	0.09
South Atlantic	0.22	0.20	0.15	0.18	0.07	0.12
East South Central	0.37	0.38	0.33	0.36	0.14	0.27
West South Central	0.37	0.41	0.27	0.35	0.12	0.16
West	0.31	0.31	0.25	0.24	0.16	0.10
Mountain	0.43	0.32	0.33	0.24	0.18	0.07
Pacific	0.40	0.42	0.33	0.33	0.22	0.14
COUNTY TYPE						
Large Metro	0.16	0.16	0.13	0.13	0.07	0.06
Small Metro	0.22	0.17	0.17	0.12	0.10	0.05
250K – 1 Mil. Pop.	0.25	0.22	0.19	0.15	0.11	0.07
< 250K Pop.	0.37	0.27	0.31	0.18	0.19	0.06
Nonmetro	0.21	0.25	0.16	0.24	0.09	0.17
Urbanized	0.36	0.38	0.30	0.38	0.14	0.32
Less Urbanized	0.31	0.41	0.20	0.37	0.14	0.22
Completely Rural	0.31	0.51	0.23	0.30	0.21	*

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.70P (REVISED) Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.3150		0.9964		0.5900	
GEOGRAPHIC DIVISION						
Northeast	0.2670		0.6307		0.4267	
New England	0.9430		0.5216		0.3445	
Middle Atlantic	0.2004		0.7736		0.5886	
Midwest	0.4540		0.6511		0.5725	
East North Central	0.0432		0.0743		0.2356	
West North Central	0.1488		0.0706		0.6091	
South	0.6511		0.5886		0.3994	
South Atlantic	0.2698		0.9932		0.3507	
East South Central	0.6194		0.7769		0.8803	
West South Central	0.6105		0.3786		0.7850	
West	0.8012		0.9707		0.3919	
Mountain	0.0465		0.1817		0.4069	
Pacific	0.6094		0.6269		0.5286	
COUNTY TYPE						
Large Metro	0.6903		0.7486		0.8011	
Small Metro	0.3168		0.4000		0.2165	
250K – 1 Mil. Pop.	0.8860		0.7600		0.5166	
< 250K Pop.	0.0415		0.0582		0.2372	
Nonmetro	0.6863		0.5978		0.5400	
Urbanized	0.3083		0.7764		0.5576	
Less Urbanized	0.7769		0.3227		0.6748	
Completely Rural	0.7035		0.5920		*	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.71C (REVISED) Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	113	112	69	69	36	39
GEOGRAPHIC DIVISION						
Northeast	51	45	34	33	20	23
New England	25	19	17	13	12	9
Middle Atlantic	44	41	29	31	16	21
Midwest	43	43	27	24	15	12
East North Central	36	35	24	21	14	10
West North Central	24	25	14	14	6	6
South	64	70	39	36	19	18
South Atlantic	49	51	29	29	15	15
East South Central	19	21	10	9	4	5
West South Central	39	43	23	19	11	8
West	69	63	39	43	18	24
Mountain	34	26	17	16	9	8
Pacific	61	58	35	40	15	23
COUNTY TYPE						
Large Metro	95	99	55	58	28	32
Small Metro	79	76	46	44	21	21
250K – 1 Mil. Pop.	61	56	36	35	18	18
< 250K Pop.	48	52	29	25	10	11
Nonmetro	49	43	26	22	11	12
Urbanized	39	31	19	18	6	9
Less Urbanized	30	31	17	13	10	7
Completely Rural	9	12	3	1	1	1

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.71D (REVISED) Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.34	0.33	0.20	0.20	0.11	0.11
GEOGRAPHIC DIVISION						
Northeast	0.83	0.75	0.55	0.55	0.32	0.38
New England	1.59	1.20	1.09	0.82	0.74	0.57
Middle Atlantic	0.97	0.92	0.64	0.69	0.34	0.48
Midwest	0.59	0.57	0.37	0.33	0.21	0.15
East North Central	0.70	0.68	0.48	0.41	0.28	0.19
West North Central	1.06	1.06	0.60	0.59	0.26	0.26
South	0.54	0.56	0.33	0.29	0.16	0.14
South Atlantic	0.80	0.82	0.48	0.46	0.24	0.24
East South Central	1.02	1.05	0.53	0.47	0.24	0.24
West South Central	0.97	1.05	0.58	0.46	0.28	0.20
West	0.86	0.78	0.48	0.53	0.22	0.30
Mountain	1.39	1.06	0.71	0.66	0.38	0.31
Pacific	1.07	1.03	0.62	0.71	0.27	0.41
COUNTY TYPE						
Large Metro	0.48	0.48	0.29	0.30	0.16	0.17
Small Metro	0.62	0.59	0.40	0.36	0.19	0.18
250K – 1 Mil. Pop.	0.81	0.70	0.51	0.47	0.26	0.25
< 250K Pop.	0.96	1.08	0.66	0.58	0.25	0.26
Nonmetro	0.75	0.78	0.40	0.43	0.21	0.24
Urbanized	1.17	1.16	0.55	0.73	0.24	0.41
Less Urbanized	1.04	1.17	0.64	0.55	0.40	0.31
Completely Rural	2.50	2.65	1.06	0.28	0.39	0.18

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.71P (REVISED) Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.0014		0.0247		0.5804	
GEOGRAPHIC DIVISION						
Northeast	0.0096		0.0151		0.9104	
New England	0.0637		0.0316		0.1186	
Middle Atlantic	0.0466		0.0987		0.5464	
Midwest	0.2454		0.2181		0.5267	
East North Central	0.1681		0.3764		0.4240	
West North Central	0.9426		0.3960		0.8252	
South	0.0721		0.0545		0.9924	
South Atlantic	0.3896		0.3204		0.8466	
East South Central	0.1498		0.4367		0.5801	
West South Central	0.2405		0.0913		0.5743	
West	0.3590		0.5025		0.1384	
Mountain	0.0050		0.4065		0.4157	
Pacific	0.8986		0.3168		0.0549	
COUNTY TYPE						
Large Metro	0.0139		0.0943		0.5538	
Small Metro	0.1512		0.2214		0.7927	
250K – 1 Mil. Pop.	0.0841		0.2588		0.6467	
< 250K Pop.	0.8505		0.5820		0.7755	
Nonmetro	0.1267		0.2556		0.5433	
Urbanized	0.7738		0.6517		0.2406	
Less Urbanized	0.0515		0.5985		0.9487	
Completely Rural	0.8635		0.0069		0.2438	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.72C (REVISED) Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	646	663	189	183	127	99
GEOGRAPHIC DIVISION						
Northeast	290	277	84	94	55	46
New England	150	132	48	41	39	28
Middle Atlantic	247	242	69	84	39	36
Midwest	218	236	64	59	39	31
East North Central	171	196	49	53	33	28
West North Central	140	131	39	24	20	12
South	356	405	117	103	96	69
South Atlantic	284	309	99	80	83	55
East South Central	123	127	26	36	19	23
West South Central	190	214	56	48	45	36
West	390	403	101	103	49	49
Mountain	189	180	54	60	32	15
Pacific	336	363	87	84	40	46
COUNTY TYPE						
Large Metro	566	573	134	159	84	83
Small Metro	414	420	113	82	87	56
250K – 1 Mil. Pop.	347	339	103	67	85	51
< 250K Pop.	240	250	48	47	22	21
Nonmetro	236	312	62	45	42	14
Urbanized	165	197	34	20	7	12
Less Urbanized	158	248	49	40	41	8
Completely Rural	59	53	16	8	*	1

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.72D (REVISED) Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.33	0.34	0.10	0.09	0.07	0.05
GEOGRAPHIC DIVISION						
Northeast	0.81	0.76	0.23	0.26	0.15	0.13
New England	1.59	1.38	0.51	0.44	0.41	0.29
Middle Atlantic	0.93	0.91	0.26	0.31	0.15	0.14
Midwest	0.51	0.56	0.15	0.14	0.09	0.07
East North Central	0.58	0.66	0.17	0.18	0.11	0.10
West North Central	1.09	1.02	0.30	0.19	0.16	0.10
South	0.50	0.57	0.16	0.14	0.14	0.10
South Atlantic	0.75	0.81	0.26	0.21	0.22	0.14
East South Central	1.05	1.09	0.22	0.31	0.16	0.19
West South Central	0.88	0.98	0.26	0.22	0.21	0.16
West	0.88	0.90	0.23	0.23	0.11	0.11
Mountain	1.39	1.31	0.40	0.44	0.24	0.11
Pacific	1.10	1.17	0.28	0.27	0.13	0.15
COUNTY TYPE						
Large Metro	0.48	0.48	0.13	0.15	0.08	0.08
Small Metro	0.61	0.57	0.19	0.14	0.15	0.09
250K – 1 Mil. Pop.	0.73	0.70	0.26	0.17	0.22	0.13
< 250K Pop.	1.15	1.02	0.25	0.24	0.11	0.11
Nonmetro	0.65	0.86	0.19	0.14	0.13	0.04
Urbanized	1.06	1.18	0.24	0.15	0.05	0.09
Less Urbanized	0.95	1.37	0.33	0.26	0.27	0.05
Completely Rural	1.29	1.55	0.45	0.28	*	0.02

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.72P (REVISED) Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.3554		0.7191		0.2070	
GEOGRAPHIC DIVISION						
Northeast	0.8240		0.9273		0.5980	
New England	0.0550		0.6736		0.8329	
Middle Atlantic	0.1632		0.8694		0.5886	
Midwest	0.6367		0.6449		0.3833	
East North Central	0.9592		0.9273		0.6756	
West North Central	0.4709		0.3191		0.2894	
South	0.1360		0.9860		0.2623	
South Atlantic	0.0404		0.9593		0.3506	
East South Central	0.4762		0.1863		0.6709	
West South Central	0.8058		0.3279		0.3486	
West	0.8441		0.7694		0.8871	
Mountain	0.2323		0.8682		0.1753	
Pacific	0.6919		0.8098		0.3537	
COUNTY TYPE						
Large Metro	0.3294		0.3830		0.9152	
Small Metro	0.8885		0.1862		0.1865	
250K – 1 Mil. Pop.	0.8271		0.1612		0.1817	
< 250K Pop.	0.9548		0.8416		0.8762	
Nonmetro	0.5687		0.2410		0.2066	
Urbanized	0.4278		0.3331		0.3655	
Less Urbanized	0.8094		0.4451		0.1008	
Completely Rural	0.3892		0.7020		*	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.73C (REVISED) Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	654	680	208	202	135	106
GEOGRAPHIC DIVISION						
Northeast	298	283	95	100	62	51
New England	151	134	53	43	42	29
Middle Atlantic	255	248	79	90	45	42
Midwest	226	243	72	64	44	34
East North Central	178	200	56	57	38	31
West North Central	144	136	43	27	22	14
South	364	411	126	109	98	71
South Atlantic	292	315	105	88	83	57
East South Central	126	126	28	36	19	23
West South Central	192	217	62	52	47	36
West	390	411	111	119	55	57
Mountain	192	179	58	67	34	16
Pacific	336	373	97	98	46	54
COUNTY TYPE						
Large Metro	581	595	151	175	91	92
Small Metro	425	441	125	94	91	59
250K – 1 Mil. Pop.	360	351	110	75	88	54
< 250K Pop.	241	266	57	57	24	23
Nonmetro	244	324	67	52	43	18
Urbanized	172	209	39	30	9	15
Less Urbanized	165	260	52	42	42	11
Completely Rural	61	56	17	9	1	1

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.73D (REVISED) Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.29	0.30	0.09	0.09	0.06	0.05
GEOGRAPHIC DIVISION						
Northeast	0.71	0.67	0.23	0.24	0.15	0.12
New England	1.36	1.20	0.48	0.39	0.38	0.26
Middle Atlantic	0.82	0.80	0.25	0.29	0.14	0.13
Midwest	0.45	0.49	0.14	0.13	0.09	0.07
East North Central	0.51	0.57	0.16	0.17	0.11	0.09
West North Central	0.95	0.90	0.29	0.18	0.15	0.09
South	0.44	0.49	0.15	0.13	0.12	0.08
South Atlantic	0.66	0.71	0.24	0.20	0.19	0.13
East South Central	0.93	0.92	0.20	0.27	0.14	0.17
West South Central	0.76	0.84	0.24	0.20	0.18	0.14
West	0.75	0.78	0.21	0.22	0.11	0.11
Mountain	1.20	1.10	0.36	0.41	0.21	0.10
Pacific	0.93	1.01	0.27	0.27	0.13	0.15
COUNTY TYPE						
Large Metro	0.42	0.42	0.12	0.14	0.08	0.07
Small Metro	0.53	0.50	0.18	0.13	0.13	0.08
250K – 1 Mil. Pop.	0.64	0.61	0.24	0.16	0.19	0.12
< 250K Pop.	0.94	0.89	0.24	0.24	0.10	0.10
Nonmetro	0.57	0.76	0.17	0.14	0.11	0.05
Urbanized	0.90	1.05	0.22	0.18	0.06	0.09
Less Urbanized	0.83	1.23	0.29	0.23	0.24	0.06
Completely Rural	1.22	1.46	0.42	0.25	0.03	0.02

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.73P (REVISED) Cocaine Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.7190		0.2844		0.3200	
GEOGRAPHIC DIVISION						
Northeast	0.8594		0.3414		0.5967	
New England	0.0310		0.3079		0.5989	
Middle Atlantic	0.2952		0.6213		0.8134	
Midwest	0.4998		0.4087		0.3204	
East North Central	0.7630		0.8226		0.5166	
West North Central	0.4689		0.2232		0.3713	
South	0.2553		0.5664		0.2791	
South Atlantic	0.0666		0.8291		0.3862	
East South Central	0.3378		0.2793		0.5915	
West South Central	0.9920		0.1254		0.2972	
West	0.9632		0.9766		0.4816	
Mountain	0.4319		0.7333		0.1273	
Pacific	0.7256		0.8697		0.1202	
COUNTY TYPE						
Large Metro	0.5763		0.8092		0.9208	
Small Metro	0.6745		0.1306		0.1983	
250K – 1 Mil. Pop.	0.6020		0.1026		0.1689	
< 250K Pop.	1.0000		0.8035		0.9426	
Nonmetro	0.7445		0.1075		0.2980	
Urbanized	0.4633		0.2048		0.2027	
Less Urbanized	0.9689		0.3420		0.1047	
Completely Rural	0.4501		0.3835		0.5998	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.