

**Table 1.74C (REVISED) Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	316	379	105	92	77	60
GEOGRAPHIC DIVISION						
Northeast	111	136	53	45	46	30
New England	48	79	37	29	36	24
Middle Atlantic	100	112	39	35	29	18
Midwest	121	111	44	35	29	24
East North Central	88	84	31	32	24	20
West North Central	82	70	31	13	17	13
South	187	256	64	65	49	45
South Atlantic	142	205	53	42	45	29
East South Central	70	81	20	26	4	18
West South Central	102	109	31	35	20	30
West	191	193	38	32	25	12
Mountain	84	89	24	13	21	5
Pacific	171	171	29	29	13	11
COUNTY TYPE						
Large Metro	232	257	73	70	48	40
Small Metro	178	227	63	60	56	45
250K – 1 Mil. Pop.	139	176	57	51	53	44
< 250K Pop.	108	144	25	32	20	10
Nonmetro	135	145	35	15	21	9
Urbanized	92	105	16	12	7	9
Less Urbanized	92	101	31	8	20	4
Completely Rural	33	20	2	1	*	*

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.74D (REVISED) Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.13	0.15	0.04	0.04	0.03	0.02
GEOGRAPHIC DIVISION						
Northeast	0.24	0.29	0.12	0.10	0.10	0.06
New England	0.39	0.64	0.30	0.23	0.30	0.19
Middle Atlantic	0.29	0.33	0.11	0.10	0.08	0.05
Midwest	0.22	0.20	0.08	0.06	0.05	0.04
East North Central	0.23	0.22	0.08	0.08	0.06	0.05
West North Central	0.49	0.42	0.19	0.08	0.10	0.07
South	0.20	0.28	0.07	0.07	0.05	0.05
South Atlantic	0.29	0.42	0.11	0.09	0.09	0.06
East South Central	0.47	0.54	0.14	0.17	0.03	0.12
West South Central	0.36	0.38	0.11	0.12	0.07	0.10
West	0.33	0.33	0.07	0.05	0.04	0.02
Mountain	0.47	0.49	0.13	0.07	0.12	0.03
Pacific	0.42	0.42	0.07	0.07	0.03	0.03
COUNTY TYPE						
Large Metro	0.17	0.19	0.05	0.05	0.04	0.03
Small Metro	0.22	0.28	0.08	0.08	0.07	0.06
250K – 1 Mil. Pop.	0.26	0.33	0.11	0.10	0.10	0.08
< 250K Pop.	0.40	0.53	0.09	0.12	0.08	0.04
Nonmetro	0.31	0.35	0.08	0.04	0.05	0.02
Urbanized	0.48	0.61	0.09	0.07	0.04	0.05
Less Urbanized	0.45	0.46	0.16	0.04	0.11	0.02
Completely Rural	0.66	0.51	0.05	0.02	*	*

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.74P (REVISED) Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.1350		0.3530		0.2224	
GEOGRAPHIC DIVISION						
Northeast	0.0259		0.6477		0.4759	
New England	0.1251		0.9587		0.9096	
Middle Atlantic	0.0940		0.4743		0.2554	
Midwest	0.8077		0.9828		0.8991	
East North Central	0.8812		0.4795		0.8149	
West North Central	0.8366		0.3712		0.9010	
South	0.1812		0.4442		0.5252	
South Atlantic	0.0638		0.1916		0.0977	
East South Central	0.6234		0.5012		0.0801	
West South Central	0.5177		0.9539		0.7101	
West	0.9180		0.5819		0.1488	
Mountain	0.6172		0.2679		0.1416	
Pacific	0.7118		0.9463		0.6341	
COUNTY TYPE						
Large Metro	0.1956		0.7420		0.6623	
Small Metro	0.1956		0.9691		0.3845	
250K – 1 Mil. Pop.	0.1824		0.8739		0.5308	
< 250K Pop.	0.6663		0.6895		0.3349	
Nonmetro	0.7302		0.0069		0.2261	
Urbanized	0.8955		0.2916		0.8627	
Less Urbanized	0.9608		0.0138		0.1417	
Completely Rural	0.0594		0.2208		*	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.