

Table 1.76C (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	473	473	111	115	58	83
GEOGRAPHIC DIVISION						
Northeast	217	205	43	55	15	34
New England	101	114	19	18	11	12
Middle Atlantic	191	169	39	52	10	32
Midwest	177	157	26	33	12	20
East North Central	138	134	22	30	10	19
West North Central	111	90	12	15	7	5
South	257	273	52	77	18	69
South Atlantic	200	213	39	73	16	68
East South Central	96	97	29	17	5	8
West South Central	141	140	19	24	7	10
West	283	318	86	55	51	26
Mountain	124	145	17	20	11	13
Pacific	255	285	84	52	50	22
COUNTY TYPE						
Large Metro	408	414	101	103	53	77
Small Metro	256	298	44	47	19	30
250K – 1 Mil. Pop.	216	250	39	41	16	26
< 250K Pop.	161	165	20	24	10	16
Nonmetro	181	195	26	23	12	10
Urbanized	128	133	15	13	7	7
Less Urbanized	111	146	20	16	9	7
Completely Rural	73	42	6	9	3	3

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.76D (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.19	0.19	0.04	0.05	0.02	0.03
GEOGRAPHIC DIVISION						
Northeast	0.47	0.44	0.09	0.12	0.03	0.07
New England	0.83	0.93	0.16	0.15	0.09	0.10
Middle Atlantic	0.56	0.49	0.11	0.15	0.03	0.09
Midwest	0.32	0.28	0.05	0.06	0.02	0.04
East North Central	0.36	0.35	0.06	0.08	0.03	0.05
West North Central	0.66	0.53	0.07	0.09	0.04	0.03
South	0.28	0.29	0.06	0.08	0.02	0.07
South Atlantic	0.41	0.43	0.08	0.15	0.03	0.14
East South Central	0.64	0.65	0.19	0.11	0.03	0.05
West South Central	0.50	0.49	0.07	0.08	0.03	0.03
West	0.49	0.54	0.15	0.09	0.09	0.04
Mountain	0.70	0.81	0.10	0.11	0.06	0.07
Pacific	0.63	0.70	0.21	0.13	0.12	0.05
COUNTY TYPE						
Large Metro	0.28	0.27	0.07	0.08	0.04	0.06
Small Metro	0.30	0.33	0.06	0.06	0.03	0.04
250K – 1 Mil. Pop.	0.37	0.43	0.08	0.08	0.03	0.05
< 250K Pop.	0.55	0.52	0.08	0.09	0.04	0.06
Nonmetro	0.40	0.44	0.06	0.06	0.03	0.03
Urbanized	0.62	0.66	0.08	0.07	0.04	0.04
Less Urbanized	0.54	0.66	0.10	0.08	0.05	0.04
Completely Rural	1.46	1.05	0.15	0.25	0.07	0.07

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.76P (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2010 Versus 2009**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.1696		0.6336		0.2254	
GEOGRAPHIC DIVISION						
Northeast	0.1945		0.2768		0.0901	
New England	0.2201		0.6482		0.6857	
Middle Atlantic	0.4479		0.1756		0.0341	
Midwest	0.1721		0.4906		0.4658	
East North Central	0.6695		0.6283		0.2158	
West North Central	0.0968		0.5682		0.2598	
South	0.9804		0.6564		0.0167	
South Atlantic	0.9311		0.5805		0.0501	
East South Central	0.5779		0.6239		0.1880	
West South Central	0.5808		0.6882		0.0836	
West	0.6311		0.0321		0.0318	
Mountain	0.9884		0.3637		0.2128	
Pacific	0.6056		0.0145		0.0096	
COUNTY TYPE						
Large Metro	0.2348		0.7879		0.3686	
Small Metro	0.5094		0.8542		0.2063	
250K – 1 Mil. Pop.	0.6525		0.9165		0.2496	
< 250K Pop.	0.6317		0.5767		0.5772	
Nonmetro	0.0387		0.4130		0.5988	
Urbanized	0.2072		0.4356		0.5114	
Less Urbanized	0.2166		0.3549		0.8936	
Completely Rural	0.0556		0.3740		0.9731	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.77C (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	66	61	42	42	22	23
GEOGRAPHIC DIVISION						
Northeast	25	23	15	16	7	9
New England	11	11	7	8	4	5
Middle Atlantic	22	20	13	13	6	7
Midwest	23	23	15	16	8	9
East North Central	18	20	13	14	8	9
West North Central	14	12	8	8	4	3
South	37	36	25	25	11	14
South Atlantic	30	26	19	18	8	10
East South Central	13	13	9	10	4	6
West South Central	20	19	13	12	5	8
West	40	38	27	26	17	13
Mountain	17	16	9	10	6	4
Pacific	37	35	25	24	16	13
COUNTY TYPE						
Large Metro	57	52	34	35	17	19
Small Metro	40	37	24	22	14	13
250K – 1 Mil. Pop.	32	31	21	18	12	10
< 250K Pop.	25	21	13	13	6	9
Nonmetro	24	21	15	15	7	6
Urbanized	16	13	11	10	5	3
Less Urbanized	16	15	9	11	4	5
Completely Rural	9	7	6	5	3	2

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 1.77D (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
Standard Errors of Percentages, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.27	0.25	0.17	0.17	0.09	0.10
GEOGRAPHIC DIVISION						
Northeast	0.58	0.55	0.35	0.37	0.16	0.21
New England	0.98	1.03	0.62	0.74	0.37	0.43
Middle Atlantic	0.70	0.64	0.42	0.43	0.18	0.24
Midwest	0.42	0.42	0.28	0.29	0.16	0.17
East North Central	0.48	0.53	0.34	0.36	0.20	0.24
West North Central	0.88	0.74	0.49	0.50	0.22	0.18
South	0.41	0.40	0.28	0.27	0.12	0.16
South Atlantic	0.66	0.58	0.42	0.40	0.18	0.23
East South Central	0.90	0.93	0.64	0.70	0.29	0.43
West South Central	0.65	0.62	0.45	0.41	0.18	0.25
West	0.68	0.66	0.46	0.44	0.29	0.23
Mountain	0.92	0.90	0.52	0.58	0.34	0.24
Pacific	0.90	0.87	0.62	0.59	0.38	0.31
COUNTY TYPE						
Large Metro	0.40	0.37	0.25	0.26	0.13	0.14
Small Metro	0.45	0.44	0.30	0.28	0.17	0.17
250K – 1 Mil. Pop.	0.53	0.54	0.37	0.33	0.23	0.19
< 250K Pop.	0.82	0.70	0.49	0.52	0.23	0.34
Nonmetro	0.56	0.50	0.37	0.38	0.18	0.15
Urbanized	0.81	0.75	0.60	0.58	0.30	0.19
Less Urbanized	0.77	0.72	0.47	0.54	0.22	0.23
Completely Rural	1.86	1.98	1.40	1.61	0.71	0.64

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.77P (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.0043		0.2555		0.5039	
GEOGRAPHIC DIVISION						
Northeast	0.4360		0.2287		0.2024	
New England	0.1084		0.2795		0.3194	
Middle Atlantic	0.9832		0.4445		0.3887	
Midwest	0.3133		0.4334		0.7826	
East North Central	0.3635		0.2576		0.9812	
West North Central	0.6472		0.7395		0.4036	
South	0.0013		0.2556		0.0470	
South Atlantic	0.0288		0.4570		0.5047	
East South Central	0.5916		0.9299		0.2543	
West South Central	0.0058		0.2501		0.0537	
West	0.1576		0.3232		0.1989	
Mountain	0.7137		0.7809		0.7354	
Pacific	0.0813		0.3447		0.2212	
COUNTY TYPE						
Large Metro	0.0464		0.6048		0.4301	
Small Metro	0.2610		0.1128		0.7636	
250K – 1 Mil. Pop.	0.6654		0.2838		0.9294	
< 250K Pop.	0.0078		0.2125		0.4944	
Nonmetro	0.0342		0.7410		0.7381	
Urbanized	0.0995		0.8639		0.3969	
Less Urbanized	0.1827		0.6805		0.6655	
Completely Rural	0.6648		0.5167		0.8863	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.