

Table 1.78C (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	740	784	366	397	250	268
GEOGRAPHIC DIVISION						
Northeast	282	285	143	149	98	117
New England	116	146	67	61	34	49
Middle Atlantic	257	245	126	135	92	106
Midwest	266	275	132	133	81	94
East North Central	205	231	106	118	68	88
West North Central	164	149	73	63	43	32
South	461	463	247	251	166	167
South Atlantic	370	349	193	197	131	136
East South Central	143	188	83	86	66	54
West South Central	237	223	126	122	76	76
West	455	488	198	240	137	135
Mountain	200	228	91	112	66	69
Pacific	408	430	176	216	120	117
COUNTY TYPE						
Large Metro	712	740	297	322	200	206
Small Metro	464	508	209	224	126	129
250K – 1 Mil. Pop.	407	415	177	191	97	111
< 250K Pop.	258	300	126	123	80	72
Nonmetro	325	329	165	158	115	110
Urbanized	245	241	128	99	81	70
Less Urbanized	216	268	100	129	80	84
Completely Rural	97	75	42	27	28	14

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.78D (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.29	0.31	0.15	0.16	0.10	0.11
GEOGRAPHIC DIVISION						
Northeast	0.61	0.61	0.31	0.32	0.21	0.25
New England	0.95	1.19	0.55	0.50	0.28	0.40
Middle Atlantic	0.75	0.71	0.37	0.39	0.27	0.31
Midwest	0.48	0.50	0.24	0.24	0.15	0.17
East North Central	0.53	0.60	0.28	0.31	0.18	0.23
West North Central	0.98	0.89	0.43	0.38	0.25	0.19
South	0.50	0.50	0.27	0.27	0.18	0.18
South Atlantic	0.76	0.71	0.40	0.40	0.27	0.28
East South Central	0.96	1.25	0.56	0.57	0.44	0.36
West South Central	0.83	0.77	0.44	0.42	0.27	0.26
West	0.78	0.83	0.34	0.41	0.24	0.23
Mountain	1.12	1.27	0.51	0.62	0.37	0.38
Pacific	1.01	1.06	0.44	0.53	0.30	0.29
COUNTY TYPE						
Large Metro	0.43	0.45	0.21	0.23	0.14	0.15
Small Metro	0.53	0.52	0.26	0.27	0.16	0.16
250K – 1 Mil. Pop.	0.66	0.62	0.33	0.34	0.19	0.21
< 250K Pop.	0.91	0.92	0.47	0.43	0.30	0.27
Nonmetro	0.61	0.69	0.37	0.36	0.27	0.26
Urbanized	0.92	1.06	0.61	0.52	0.41	0.39
Less Urbanized	0.91	1.06	0.49	0.58	0.40	0.40
Completely Rural	1.66	1.60	0.94	0.71	0.65	0.36

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.78P (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.6301		0.8124		0.8440	
GEOGRAPHIC DIVISION						
Northeast	0.9837		0.8362		0.2656	
New England	0.8957		0.8307		0.0718	
Middle Atlantic	0.9643		0.7363		0.6487	
Midwest	0.9912		0.8659		0.9339	
East North Central	0.4751		0.7332		0.8593	
West North Central	0.3578		0.7845		0.8418	
South	0.3983		0.2106		0.4361	
South Atlantic	0.6081		0.7838		0.8590	
East South Central	0.2956		0.5110		0.2562	
West South Central	0.1169		0.1099		0.2732	
West	0.9735		0.4166		0.7042	
Mountain	0.7181		0.7082		0.6691	
Pacific	0.8918		0.4775		0.8345	
COUNTY TYPE						
Large Metro	0.5112		0.4807		0.9131	
Small Metro	0.3189		0.5160		0.9712	
250K – 1 Mil. Pop.	0.1887		0.7294		0.2809	
< 250K Pop.	0.8905		0.5331		0.1456	
Nonmetro	0.1558		0.2745		0.4968	
Urbanized	0.4892		0.2484		0.9059	
Less Urbanized	0.2496		0.8204		0.5758	
Completely Rural	0.4063		0.2923		0.2341	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.79C (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	70	68	59	58	36	38
GEOGRAPHIC DIVISION						
Northeast	22	20	19	18	12	11
New England	13	11	10	10	7	6
Middle Atlantic	18	17	16	14	10	10
Midwest	25	24	22	21	15	14
East North Central	21	20	19	17	13	11
West North Central	15	13	12	12	7	8
South	42	47	37	39	23	24
South Atlantic	32	36	28	30	16	18
East South Central	17	15	14	13	9	9
West South Central	23	25	19	21	13	14
West	43	41	34	35	20	25
Mountain	16	21	13	20	8	17
Pacific	40	34	31	29	19	17
COUNTY TYPE						
Large Metro	58	58	46	48	27	31
Small Metro	47	41	40	33	22	20
250K – 1 Mil. Pop.	34	34	27	28	16	17
< 250K Pop.	32	24	28	17	15	11
Nonmetro	27	34	21	29	13	16
Urbanized	19	27	17	22	10	14
Less Urbanized	18	21	13	17	8	8
Completely Rural	6	6	4	4	3	3

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.79D (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.28	0.28	0.24	0.24	0.15	0.16
GEOGRAPHIC DIVISION						
Northeast	0.52	0.48	0.44	0.42	0.29	0.27
New England	1.14	0.98	0.90	0.94	0.66	0.54
Middle Atlantic	0.57	0.55	0.51	0.46	0.31	0.31
Midwest	0.47	0.45	0.42	0.39	0.28	0.26
East North Central	0.55	0.54	0.49	0.46	0.36	0.29
West North Central	0.95	0.84	0.74	0.75	0.42	0.52
South	0.46	0.52	0.41	0.44	0.25	0.27
South Atlantic	0.70	0.80	0.61	0.67	0.36	0.39
East South Central	1.15	1.06	0.99	0.93	0.63	0.62
West South Central	0.76	0.83	0.64	0.71	0.44	0.46
West	0.73	0.70	0.57	0.61	0.35	0.42
Mountain	0.86	1.18	0.70	1.09	0.43	0.97
Pacific	0.97	0.85	0.76	0.72	0.46	0.42
COUNTY TYPE						
Large Metro	0.40	0.40	0.33	0.34	0.20	0.23
Small Metro	0.53	0.47	0.46	0.39	0.27	0.26
250K – 1 Mil. Pop.	0.59	0.58	0.50	0.49	0.30	0.33
< 250K Pop.	0.95	0.80	0.87	0.62	0.53	0.44
Nonmetro	0.63	0.74	0.51	0.65	0.33	0.39
Urbanized	0.96	1.25	0.88	1.10	0.51	0.76
Less Urbanized	0.89	0.95	0.68	0.83	0.46	0.42
Completely Rural	1.35	1.76	0.97	1.22	0.76	0.95

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.79P (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.0392		0.4377		0.3911	
GEOGRAPHIC DIVISION						
Northeast	0.1479		0.5447		0.5450	
New England	0.2466		0.4785		0.2240	
Middle Atlantic	0.3201		0.7927		0.9222	
Midwest	0.2579		0.6606		0.1874	
East North Central	0.0372		0.1451		0.1046	
West North Central	0.3722		0.1864		0.9710	
South	0.0916		0.2656		0.3851	
South Atlantic	0.6623		0.6221		0.9277	
East South Central	0.0086		0.0085		0.0262	
West South Central	0.4364		0.7497		0.9449	
West	0.8891		0.5732		0.5189	
Mountain	0.0280		0.0302		0.0666	
Pacific	0.1940		0.5569		0.5394	
COUNTY TYPE						
Large Metro	0.1200		0.9440		0.6621	
Small Metro	0.1684		0.2087		0.7127	
250K – 1 Mil. Pop.	0.7036		0.6038		0.5137	
< 250K Pop.	0.0064		0.0084		0.1353	
Nonmetro	0.8518		0.9808		0.4043	
Urbanized	0.5448		0.7411		0.7101	
Less Urbanized	0.7992		0.6924		0.2970	
Completely Rural	0.7786		0.8498		0.6485	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.80C (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	153	143	119	114	74	72
GEOGRAPHIC DIVISION						
Northeast	61	59	48	45	32	27
New England	27	27	24	20	17	14
Middle Atlantic	54	53	42	40	26	23
Midwest	60	62	43	45	27	28
East North Central	48	51	34	38	21	24
West North Central	36	33	26	25	17	16
South	88	93	76	71	44	46
South Atlantic	69	71	62	56	32	34
East South Central	30	34	26	27	16	18
West South Central	47	49	37	34	27	25
West	88	80	62	60	41	39
Mountain	35	34	32	25	20	16
Pacific	80	72	53	55	36	36
COUNTY TYPE						
Large Metro	146	153	97	99	56	56
Small Metro	115	127	79	85	48	46
250K – 1 Mil. Pop.	85	97	56	60	38	36
< 250K Pop.	80	80	55	57	28	31
Nonmetro	107	69	77	48	37	27
Urbanized	95	52	67	37	31	21
Less Urbanized	53	43	38	28	21	17
Completely Rural	17	20	9	13	4	2

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.80D (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.46	0.42	0.35	0.33	0.22	0.21
GEOGRAPHIC DIVISION						
Northeast	0.99	0.98	0.79	0.74	0.52	0.44
New England	1.72	1.65	1.52	1.24	1.09	0.88
Middle Atlantic	1.20	1.19	0.93	0.91	0.58	0.52
Midwest	0.81	0.83	0.59	0.60	0.36	0.38
East North Central	0.94	1.00	0.67	0.74	0.42	0.47
West North Central	1.57	1.40	1.13	1.05	0.74	0.67
South	0.73	0.75	0.63	0.57	0.37	0.37
South Atlantic	1.14	1.13	1.01	0.89	0.53	0.54
East South Central	1.60	1.69	1.36	1.37	0.85	0.93
West South Central	1.18	1.18	0.91	0.83	0.67	0.61
West	1.08	0.99	0.77	0.74	0.51	0.49
Mountain	1.47	1.37	1.31	0.99	0.81	0.63
Pacific	1.40	1.29	0.94	0.97	0.63	0.64
COUNTY TYPE						
Large Metro	0.66	0.63	0.48	0.46	0.30	0.29
Small Metro	0.77	0.72	0.63	0.58	0.42	0.37
250K – 1 Mil. Pop.	0.94	0.85	0.72	0.68	0.53	0.47
< 250K Pop.	1.35	1.27	1.15	1.01	0.65	0.62
Nonmetro	0.99	0.98	0.93	0.83	0.51	0.51
Urbanized	1.49	1.47	1.44	1.33	0.71	0.85
Less Urbanized	1.39	1.26	1.32	0.99	0.80	0.67
Completely Rural	3.65	4.05	2.36	3.22	1.22	0.65

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.80P (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.3163		0.1123		0.1020	
GEOGRAPHIC DIVISION						
Northeast	0.4593		0.1594		0.0534	
New England	0.6723		0.1394		0.1575	
Middle Atlantic	0.5060		0.4323		0.1611	
Midwest	0.1192		0.9166		0.7570	
East North Central	0.1083		0.5276		0.9477	
West North Central	0.6549		0.4716		0.6345	
South	0.7744		0.3118		0.2578	
South Atlantic	0.5160		0.6694		0.7766	
East South Central	0.2670		0.2287		0.5170	
West South Central	0.7075		0.6526		0.2431	
West	0.6797		0.4934		0.6097	
Mountain	0.1518		0.2088		0.5644	
Pacific	0.9359		0.8471		0.7543	
COUNTY TYPE						
Large Metro	0.8560		0.4934		0.5047	
Small Metro	0.3123		0.2458		0.2037	
250K – 1 Mil. Pop.	0.9728		0.3675		0.4697	
< 250K Pop.	0.0982		0.4282		0.2514	
Nonmetro	0.2900		0.2180		0.2082	
Urbanized	0.7064		0.7215		0.6862	
Less Urbanized	0.0735		0.1663		0.3653	
Completely Rural	0.7678		0.8744		0.0376	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.81C (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	705	752	329	363	228	252
GEOGRAPHIC DIVISION						
Northeast	271	280	129	142	91	114
New England	112	139	62	57	27	46
Middle Atlantic	247	242	113	129	86	104
Midwest	250	263	118	122	74	87
East North Central	194	222	96	108	63	82
West North Central	155	142	65	56	37	28
South	451	441	225	225	154	158
South Atlantic	363	333	172	180	124	131
East South Central	140	180	74	77	63	48
West South Central	227	210	117	105	69	66
West	425	467	176	224	125	121
Mountain	196	219	80	100	61	62
Pacific	376	410	158	202	109	104
COUNTY TYPE						
Large Metro	659	688	265	294	183	194
Small Metro	420	462	180	199	110	118
250K – 1 Mil. Pop.	372	378	158	173	84	103
< 250K Pop.	227	270	98	101	71	62
Nonmetro	292	299	137	137	107	98
Urbanized	213	208	100	80	73	61
Less Urbanized	194	249	87	114	74	76
Completely Rural	89	66	40	22	28	13

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.81D (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.36	0.39	0.17	0.19	0.12	0.13
GEOGRAPHIC DIVISION						
Northeast	0.75	0.77	0.36	0.39	0.25	0.31
New England	1.18	1.46	0.65	0.60	0.28	0.48
Middle Atlantic	0.93	0.91	0.43	0.48	0.33	0.39
Midwest	0.59	0.62	0.28	0.29	0.18	0.20
East North Central	0.65	0.75	0.32	0.36	0.21	0.28
West North Central	1.21	1.11	0.50	0.43	0.29	0.22
South	0.64	0.62	0.32	0.31	0.22	0.22
South Atlantic	0.96	0.87	0.45	0.47	0.33	0.34
East South Central	1.20	1.55	0.64	0.66	0.54	0.41
West South Central	1.06	0.96	0.55	0.48	0.32	0.30
West	0.96	1.04	0.40	0.50	0.28	0.27
Mountain	1.44	1.59	0.59	0.73	0.45	0.45
Pacific	1.23	1.32	0.51	0.65	0.36	0.33
COUNTY TYPE						
Large Metro	0.53	0.57	0.24	0.27	0.17	0.18
Small Metro	0.64	0.65	0.30	0.32	0.19	0.20
250K – 1 Mil. Pop.	0.81	0.77	0.39	0.42	0.22	0.26
< 250K Pop.	1.11	1.16	0.50	0.50	0.35	0.32
Nonmetro	0.77	0.83	0.42	0.41	0.33	0.30
Urbanized	1.20	1.26	0.69	0.59	0.51	0.46
Less Urbanized	1.11	1.32	0.57	0.68	0.49	0.47
Completely Rural	1.92	1.76	1.09	0.72	0.77	0.42

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.81P (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.8533		0.7143		0.6992	
GEOGRAPHIC DIVISION						
Northeast	0.8038		0.4093		0.0678	
New England	0.9854		0.6779		0.0040	
Middle Atlantic	0.7850		0.4750		0.3975	
Midwest	0.6748		0.8945		0.8592	
East North Central	0.2051		0.9489		0.6866	
West North Central	0.3161		0.6897		0.5839	
South	0.4130		0.3670		0.6846	
South Atlantic	0.5179		0.9047		0.8078	
East South Central	0.1156		0.9362		0.4805	
West South Central	0.1022		0.0886		0.4135	
West	0.9163		0.2978		0.7199	
Mountain	0.7186		0.7225		0.4673	
Pacific	0.9530		0.3308		0.9868	
COUNTY TYPE						
Large Metro	0.4095		0.3241		0.7145	
Small Metro	0.3913		0.7802		0.6725	
250K – 1 Mil. Pop.	0.1258		0.7941		0.2111	
< 250K Pop.	0.5060		0.9023		0.2897	
Nonmetro	0.2442		0.5179		0.8375	
Urbanized	0.5364		0.4036		0.8676	
Less Urbanized	0.4094		0.8539		0.8314	
Completely Rural	0.3269		0.2200		0.3243	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.82C (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	733	777	360	389	246	265
GEOGRAPHIC DIVISION						
Northeast	281	283	142	147	97	117
New England	117	144	67	60	32	48
Middle Atlantic	255	243	125	134	91	105
Midwest	261	273	128	131	79	93
East North Central	203	230	104	116	66	87
West North Central	162	149	71	62	42	33
South	459	462	244	243	163	165
South Atlantic	369	349	192	192	131	135
East South Central	144	187	80	84	64	52
West South Central	235	221	123	117	74	73
West	452	481	193	237	136	132
Mountain	207	224	89	108	65	66
Pacific	401	425	172	214	119	115
COUNTY TYPE						
Large Metro	697	731	288	315	195	203
Small Metro	456	497	206	217	126	127
250K – 1 Mil. Pop.	402	407	175	185	99	109
< 250K Pop.	250	292	120	118	77	71
Nonmetro	319	324	162	153	113	107
Urbanized	239	232	125	92	79	66
Less Urbanized	213	266	98	126	79	83
Completely Rural	94	73	42	26	28	13

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.82D (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.32	0.34	0.16	0.17	0.11	0.12
GEOGRAPHIC DIVISION						
Northeast	0.67	0.67	0.34	0.35	0.23	0.28
New England	1.05	1.29	0.61	0.54	0.29	0.43
Middle Atlantic	0.82	0.78	0.40	0.43	0.29	0.34
Midwest	0.52	0.55	0.26	0.26	0.16	0.19
East North Central	0.58	0.66	0.30	0.33	0.19	0.25
West North Central	1.07	0.98	0.47	0.40	0.28	0.22
South	0.55	0.55	0.29	0.29	0.20	0.20
South Atlantic	0.84	0.78	0.44	0.43	0.30	0.30
East South Central	1.06	1.37	0.59	0.62	0.47	0.38
West South Central	0.92	0.85	0.48	0.45	0.29	0.28
West	0.86	0.91	0.37	0.45	0.26	0.25
Mountain	1.29	1.38	0.56	0.66	0.41	0.41
Pacific	1.11	1.16	0.47	0.58	0.33	0.31
COUNTY TYPE						
Large Metro	0.47	0.50	0.22	0.25	0.15	0.16
Small Metro	0.58	0.57	0.29	0.29	0.18	0.18
250K – 1 Mil. Pop.	0.74	0.68	0.37	0.37	0.22	0.23
< 250K Pop.	1.00	1.00	0.51	0.47	0.32	0.30
Nonmetro	0.68	0.76	0.40	0.39	0.29	0.28
Urbanized	1.00	1.16	0.67	0.55	0.44	0.42
Less Urbanized	1.01	1.18	0.53	0.64	0.44	0.44
Completely Rural	1.84	1.72	1.04	0.76	0.71	0.39

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.82P (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.7222		0.9109		0.9438	
GEOGRAPHIC DIVISION						
Northeast	0.9515		0.7746		0.2329	
New England	0.9554		0.9134		0.0397	
Middle Atlantic	0.9227		0.7107		0.6537	
Midwest	0.9302		0.9307		0.7595	
East North Central	0.3772		0.9137		0.6720	
West North Central	0.3109		0.9801		0.8461	
South	0.4764		0.2790		0.5088	
South Atlantic	0.6224		0.8487		0.8493	
East South Central	0.1904		0.8009		0.4166	
West South Central	0.1320		0.0840		0.2500	
West	0.9824		0.4541		0.6289	
Mountain	0.8760		0.9769		0.4055	
Pacific	0.9591		0.4227		0.9001	
COUNTY TYPE						
Large Metro	0.4402		0.4600		0.8603	
Small Metro	0.3597		0.6653		0.9178	
250K – 1 Mil. Pop.	0.1609		0.6767		0.3298	
< 250K Pop.	0.7009		0.8815		0.2180	
Nonmetro	0.1542		0.2725		0.5625	
Urbanized	0.4994		0.2737		0.9511	
Less Urbanized	0.2504		0.7677		0.6491	
Completely Rural	0.3722		0.2861		0.2099	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.