

Table 1.83C (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	584	610	322	336	214	215
GEOGRAPHIC DIVISION						
Northeast	230	225	126	120	86	85
New England	108	125	65	52	28	40
Middle Atlantic	202	187	108	107	81	75
Midwest	215	222	106	117	67	86
East North Central	173	184	84	103	55	81
West North Central	126	114	60	54	35	29
South	365	355	218	195	142	128
South Atlantic	296	265	172	148	115	103
East South Central	111	145	58	67	43	46
West South Central	194	173	117	103	70	61
West	346	372	175	217	115	125
Mountain	150	188	86	103	63	56
Pacific	310	319	151	193	97	112
COUNTY TYPE						
Large Metro	532	530	253	271	164	168
Small Metro	340	381	184	188	114	98
250K – 1 Mil. Pop.	290	294	159	160	88	86
< 250K Pop.	197	240	110	101	74	52
Nonmetro	246	226	132	128	88	101
Urbanized	190	161	104	83	65	61
Less Urbanized	157	176	83	99	61	79
Completely Rural	67	60	24	27	20	14

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.83D (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.23	0.24	0.13	0.13	0.09	0.08
GEOGRAPHIC DIVISION						
Northeast	0.50	0.48	0.27	0.26	0.19	0.18
New England	0.89	1.02	0.53	0.43	0.23	0.33
Middle Atlantic	0.59	0.54	0.31	0.31	0.24	0.22
Midwest	0.39	0.40	0.19	0.21	0.12	0.16
East North Central	0.45	0.48	0.22	0.27	0.14	0.21
West North Central	0.75	0.68	0.36	0.32	0.21	0.17
South	0.40	0.38	0.24	0.21	0.15	0.14
South Atlantic	0.61	0.54	0.36	0.30	0.24	0.21
East South Central	0.74	0.96	0.39	0.45	0.29	0.31
West South Central	0.68	0.60	0.41	0.36	0.24	0.21
West	0.59	0.63	0.30	0.37	0.20	0.21
Mountain	0.84	1.04	0.48	0.57	0.35	0.31
Pacific	0.77	0.78	0.37	0.47	0.24	0.27
COUNTY TYPE						
Large Metro	0.35	0.34	0.18	0.20	0.12	0.12
Small Metro	0.41	0.42	0.23	0.23	0.15	0.13
250K – 1 Mil. Pop.	0.51	0.49	0.30	0.29	0.17	0.17
< 250K Pop.	0.71	0.80	0.40	0.37	0.27	0.20
Nonmetro	0.50	0.50	0.30	0.30	0.21	0.24
Urbanized	0.76	0.80	0.48	0.45	0.33	0.35
Less Urbanized	0.68	0.71	0.41	0.46	0.31	0.38
Completely Rural	1.40	1.42	0.59	0.69	0.48	0.36

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.83P (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.5212		0.5145		0.4595	
GEOGRAPHIC DIVISION						
Northeast	0.4836		0.5186		0.9441	
New England	0.7301		0.6636		0.1615	
Middle Atlantic	0.2917		0.6228		0.5867	
Midwest	0.5032		0.7164		0.4015	
East North Central	0.9641		0.9459		0.4474	
West North Central	0.1879		0.3779		0.7012	
South	0.4201		0.0181		0.0790	
South Atlantic	0.3424		0.1852		0.2713	
East South Central	0.2015		0.7068		0.5806	
West South Central	0.3267		0.0236		0.1594	
West	0.5915		0.1675		0.9835	
Mountain	0.3251		0.6860		0.4875	
Pacific	0.9101		0.1836		0.7467	
COUNTY TYPE						
Large Metro	0.9165		0.7435		0.7376	
Small Metro	0.5359		0.2512		0.3926	
250K – 1 Mil. Pop.	0.2799		0.4046		0.5941	
< 250K Pop.	0.6719		0.4541		0.0293	
Nonmetro	0.2304		0.4138		0.8889	
Urbanized	0.3138		0.2117		0.9648	
Less Urbanized	0.4250		0.8783		0.9900	
Completely Rural	0.9836		0.6188		0.5455	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.84C (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	64	64	54	54	34	35
GEOGRAPHIC DIVISION						
Northeast	20	19	18	17	11	11
New England	11	10	10	10	7	6
Middle Atlantic	17	16	15	13	9	9
Midwest	24	22	21	18	14	12
East North Central	19	18	17	15	13	10
West North Central	15	13	12	11	6	7
South	38	44	34	36	20	23
South Atlantic	29	34	25	27	14	17
East South Central	15	15	13	13	9	8
West South Central	21	24	18	20	12	13
West	39	37	31	32	20	21
Mountain	13	20	12	18	7	14
Pacific	36	31	28	25	18	16
COUNTY TYPE						
Large Metro	53	53	43	42	26	27
Small Metro	42	39	35	31	19	19
250K – 1 Mil. Pop.	30	32	25	26	15	17
< 250K Pop.	28	23	24	15	12	10
Nonmetro	25	33	20	27	11	16
Urbanized	18	26	15	22	8	14
Less Urbanized	17	20	12	16	7	8
Completely Rural	5	6	4	4	3	3

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.84D (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.26	0.26	0.22	0.22	0.14	0.14
GEOGRAPHIC DIVISION						
Northeast	0.47	0.45	0.42	0.39	0.25	0.26
New England	0.96	0.94	0.86	0.91	0.59	0.53
Middle Atlantic	0.54	0.51	0.48	0.42	0.28	0.30
Midwest	0.44	0.42	0.38	0.35	0.26	0.23
East North Central	0.51	0.49	0.44	0.39	0.33	0.27
West North Central	0.91	0.81	0.71	0.70	0.38	0.43
South	0.43	0.49	0.37	0.40	0.22	0.25
South Atlantic	0.64	0.76	0.55	0.61	0.30	0.38
East South Central	1.06	1.02	0.90	0.87	0.61	0.58
West South Central	0.71	0.80	0.60	0.66	0.40	0.42
West	0.66	0.64	0.52	0.54	0.33	0.36
Mountain	0.73	1.14	0.67	1.03	0.40	0.78
Pacific	0.89	0.77	0.68	0.63	0.45	0.39
COUNTY TYPE						
Large Metro	0.37	0.37	0.31	0.31	0.19	0.20
Small Metro	0.46	0.45	0.40	0.37	0.24	0.25
250K – 1 Mil. Pop.	0.53	0.55	0.46	0.46	0.29	0.32
< 250K Pop.	0.80	0.77	0.71	0.58	0.42	0.40
Nonmetro	0.58	0.73	0.47	0.63	0.29	0.39
Urbanized	0.90	1.26	0.81	1.09	0.47	0.76
Less Urbanized	0.86	0.93	0.62	0.78	0.39	0.40
Completely Rural	1.28	1.75	0.91	1.19	0.75	0.91

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.84P (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.1500		0.2292		0.3732	
GEOGRAPHIC DIVISION						
Northeast	0.5830		0.9932		0.7813	
New England	0.6117		0.6780		0.4583	
Middle Atlantic	0.7394		0.7885		0.3939	
Midwest	0.3625		0.3782		0.2237	
East North Central	0.0659		0.0706		0.1754	
West North Central	0.3972		0.3666		0.8909	
South	0.1403		0.1491		0.3371	
South Atlantic	0.4731		0.3900		0.8686	
East South Central	0.0439		0.0234		0.0229	
West South Central	0.6943		0.9847		0.9485	
West	0.9231		0.8621		0.8813	
Mountain	0.0221		0.0286		0.0708	
Pacific	0.3080		0.2973		0.3154	
COUNTY TYPE						
Large Metro	0.1485		0.4820		0.4826	
Small Metro	0.6352		0.2456		0.7463	
250K – 1 Mil. Pop.	0.3429		0.6405		0.4211	
< 250K Pop.	0.0388		0.0088		0.0753	
Nonmetro	0.9316		0.9911		0.6446	
Urbanized	0.8190		0.8497		0.9757	
Less Urbanized	0.9593		0.8074		0.3816	
Completely Rural	0.6475		0.8189		0.7852	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.85C (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	139	134	103	100	62	64
GEOGRAPHIC DIVISION						
Northeast	57	54	43	37	27	24
New England	25	26	21	17	14	11
Middle Atlantic	51	47	38	32	24	21
Midwest	55	57	37	42	23	28
East North Central	45	47	30	35	19	23
West North Central	33	30	21	23	14	16
South	80	83	65	60	36	36
South Atlantic	60	63	50	46	25	26
East South Central	27	31	21	24	14	15
West South Central	45	46	35	32	23	21
West	81	77	55	53	35	36
Mountain	34	31	31	22	18	14
Pacific	73	71	45	48	30	33
COUNTY TYPE						
Large Metro	130	134	85	84	47	50
Small Metro	103	107	68	70	40	38
250K – 1 Mil. Pop.	74	82	49	52	32	32
< 250K Pop.	72	70	46	47	24	23
Nonmetro	86	61	60	39	28	22
Urbanized	74	48	51	28	22	16
Less Urbanized	44	38	31	23	17	15
Completely Rural	16	18	7	13	4	2

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.85D (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.42	0.39	0.31	0.29	0.19	0.19
GEOGRAPHIC DIVISION						
Northeast	0.93	0.89	0.70	0.61	0.45	0.39
New England	1.54	1.60	1.29	1.07	0.85	0.69
Middle Atlantic	1.12	1.07	0.83	0.73	0.52	0.48
Midwest	0.75	0.77	0.50	0.56	0.31	0.37
East North Central	0.88	0.92	0.59	0.68	0.37	0.45
West North Central	1.44	1.26	0.95	0.96	0.60	0.66
South	0.67	0.67	0.54	0.49	0.30	0.29
South Atlantic	0.99	1.00	0.81	0.73	0.41	0.41
East South Central	1.44	1.58	1.13	1.20	0.72	0.75
West South Central	1.11	1.13	0.88	0.77	0.58	0.50
West	1.00	0.95	0.67	0.65	0.43	0.44
Mountain	1.39	1.24	1.28	0.88	0.75	0.55
Pacific	1.29	1.26	0.78	0.86	0.52	0.59
COUNTY TYPE						
Large Metro	0.61	0.58	0.43	0.41	0.26	0.27
Small Metro	0.73	0.67	0.56	0.51	0.36	0.31
250K – 1 Mil. Pop.	0.85	0.81	0.65	0.63	0.45	0.41
< 250K Pop.	1.31	1.22	1.00	0.85	0.58	0.48
Nonmetro	0.92	0.95	0.73	0.71	0.41	0.44
Urbanized	1.29	1.46	1.05	1.13	0.56	0.71
Less Urbanized	1.36	1.28	1.11	0.85	0.67	0.61
Completely Rural	3.53	3.89	2.01	3.20	1.22	0.62

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.85P (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.2944		0.0313		0.1568	
GEOGRAPHIC DIVISION						
Northeast	0.2719		0.0177		0.0260	
New England	0.6622		0.0382		0.1090	
Middle Atlantic	0.2901		0.1308		0.0943	
Midwest	0.1004		0.5846		0.2180	
East North Central	0.1325		0.9849		0.3886	
West North Central	0.4738		0.3111		0.3584	
South	0.8689		0.0126		0.0627	
South Atlantic	0.5996		0.0871		0.2991	
East South Central	0.2207		0.2737		0.1143	
West South Central	0.8949		0.1258		0.3823	
West	0.7693		0.9411		0.9016	
Mountain	0.2608		0.1568		0.2774	
Pacific	0.4516		0.4060		0.5215	
COUNTY TYPE						
Large Metro	0.8667		0.1796		0.6547	
Small Metro	0.2543		0.2931		0.3048	
250K – 1 Mil. Pop.	0.9225		0.4573		0.8275	
< 250K Pop.	0.0646		0.4319		0.1601	
Nonmetro	0.3470		0.0949		0.1347	
Urbanized	0.6563		0.4739		0.8138	
Less Urbanized	0.0969		0.0813		0.1775	
Completely Rural	0.8297		0.8205		0.0341	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.86C (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	544	579	290	305	198	196
GEOGRAPHIC DIVISION						
Northeast	217	217	114	112	80	80
New England	102	118	59	50	23	38
Middle Atlantic	191	181	97	100	76	71
Midwest	200	209	96	106	60	78
East North Central	161	175	76	93	50	74
West North Central	117	106	55	50	32	25
South	350	339	198	173	134	121
South Atlantic	285	250	154	132	111	99
East South Central	104	138	51	59	40	42
West South Central	183	161	109	88	65	53
West	314	346	155	205	106	112
Mountain	140	176	77	94	58	52
Pacific	279	297	134	184	88	99
COUNTY TYPE						
Large Metro	478	491	223	249	150	156
Small Metro	302	338	157	162	102	86
250K – 1 Mil. Pop.	261	261	140	140	79	77
< 250K Pop.	166	211	86	84	65	46
Nonmetro	218	198	110	114	84	92
Urbanized	164	132	83	69	60	56
Less Urbanized	136	157	71	88	56	73
Completely Rural	62	51	21	22	19	13

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.86D (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.28	0.30	0.15	0.16	0.10	0.10
GEOGRAPHIC DIVISION						
Northeast	0.60	0.60	0.32	0.31	0.22	0.22
New England	1.08	1.24	0.62	0.53	0.24	0.40
Middle Atlantic	0.72	0.68	0.37	0.38	0.29	0.26
Midwest	0.47	0.49	0.23	0.25	0.14	0.18
East North Central	0.54	0.59	0.26	0.31	0.17	0.25
West North Central	0.91	0.83	0.43	0.39	0.25	0.19
South	0.49	0.47	0.28	0.24	0.19	0.17
South Atlantic	0.75	0.65	0.41	0.35	0.29	0.26
East South Central	0.90	1.18	0.44	0.50	0.34	0.36
West South Central	0.85	0.74	0.51	0.41	0.30	0.24
West	0.71	0.77	0.35	0.46	0.24	0.25
Mountain	1.03	1.28	0.57	0.69	0.43	0.38
Pacific	0.91	0.95	0.44	0.59	0.29	0.32
COUNTY TYPE						
Large Metro	0.42	0.43	0.21	0.24	0.14	0.15
Small Metro	0.49	0.52	0.26	0.27	0.17	0.15
250K – 1 Mil. Pop.	0.62	0.60	0.35	0.34	0.21	0.19
< 250K Pop.	0.83	0.99	0.43	0.42	0.32	0.24
Nonmetro	0.61	0.59	0.33	0.35	0.26	0.28
Urbanized	0.98	0.94	0.55	0.51	0.42	0.42
Less Urbanized	0.81	0.86	0.47	0.54	0.37	0.45
Completely Rural	1.61	1.50	0.61	0.70	0.55	0.42

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.86P (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.7444		0.8263		0.8211	
GEOGRAPHIC DIVISION						
Northeast	0.7071		0.8345		0.4412	
New England	0.6267		0.7295		0.0184	
Middle Atlantic	0.4524		0.9715		0.8638	
Midwest	0.8181		0.7577		0.4909	
East North Central	0.5483		0.8167		0.4426	
West North Central	0.1632		0.8198		0.9522	
South	0.4927		0.1278		0.2280	
South Atlantic	0.2940		0.4450		0.3838	
East South Central	0.0614		0.6534		0.7422	
West South Central	0.3369		0.0460		0.2003	
West	0.6157		0.1392		0.9446	
Mountain	0.3282		0.6363		0.4276	
Pacific	0.9454		0.1600		0.7194	
COUNTY TYPE						
Large Metro	0.7778		0.3615		0.9155	
Small Metro	0.6081		0.3829		0.5213	
250K – 1 Mil. Pop.	0.1660		0.4068		0.6525	
< 250K Pop.	0.3205		0.7892		0.0887	
Nonmetro	0.3935		0.8688		0.7043	
Urbanized	0.3568		0.4315		0.8908	
Less Urbanized	0.7302		0.6965		0.6548	
Completely Rural	0.8844		0.7947		0.7709	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.87C (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	577	606	318	329	212	212
GEOGRAPHIC DIVISION						
Northeast	229	223	126	118	85	84
New England	108	123	64	51	27	40
Middle Atlantic	202	186	108	106	80	74
Midwest	212	220	104	116	65	86
East North Central	172	183	83	102	53	80
West North Central	123	114	58	54	35	30
South	363	355	215	189	140	127
South Atlantic	294	265	171	142	115	102
East South Central	110	145	56	65	41	45
West South Central	191	172	114	99	68	59
West	339	367	171	215	115	122
Mountain	152	182	85	99	62	54
Pacific	302	317	148	192	97	110
COUNTY TYPE						
Large Metro	519	523	245	264	159	166
Small Metro	332	370	181	181	114	95
250K – 1 Mil. Pop.	284	285	156	154	90	83
< 250K Pop.	189	233	104	99	71	52
Nonmetro	242	221	130	125	88	99
Urbanized	186	153	101	77	64	59
Less Urbanized	154	174	81	97	60	79
Completely Rural	66	58	24	26	20	13

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.87D (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.25	0.26	0.14	0.14	0.09	0.09
GEOGRAPHIC DIVISION						
Northeast	0.54	0.53	0.30	0.28	0.20	0.20
New England	0.98	1.11	0.58	0.46	0.25	0.36
Middle Atlantic	0.65	0.60	0.35	0.34	0.26	0.24
Midwest	0.43	0.44	0.21	0.23	0.13	0.17
East North Central	0.50	0.52	0.24	0.29	0.15	0.23
West North Central	0.82	0.75	0.39	0.36	0.23	0.20
South	0.44	0.42	0.26	0.22	0.17	0.15
South Atlantic	0.67	0.59	0.39	0.32	0.26	0.23
East South Central	0.81	1.06	0.41	0.48	0.31	0.33
West South Central	0.75	0.67	0.45	0.38	0.27	0.23
West	0.65	0.69	0.33	0.41	0.22	0.23
Mountain	0.95	1.12	0.53	0.61	0.38	0.33
Pacific	0.83	0.86	0.41	0.52	0.27	0.30
COUNTY TYPE						
Large Metro	0.38	0.38	0.19	0.21	0.13	0.13
Small Metro	0.45	0.46	0.26	0.24	0.17	0.14
250K – 1 Mil. Pop.	0.56	0.54	0.33	0.31	0.20	0.18
< 250K Pop.	0.78	0.87	0.44	0.40	0.29	0.22
Nonmetro	0.55	0.56	0.32	0.32	0.23	0.26
Urbanized	0.83	0.89	0.53	0.49	0.36	0.38
Less Urbanized	0.75	0.79	0.45	0.50	0.34	0.42
Completely Rural	1.53	1.52	0.64	0.74	0.52	0.38

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.87P (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2010 Versus 2009

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	0.5991		0.6546		0.5504	
GEOGRAPHIC DIVISION						
Northeast	0.5015		0.5181		0.9716	
New England	0.7033		0.7068		0.1165	
Middle Atlantic	0.2981		0.5971		0.5161	
Midwest	0.5554		0.5901		0.2889	
East North Central	0.8256		0.8123		0.3223	
West North Central	0.1492		0.4791		0.6798	
South	0.5165		0.0330		0.1066	
South Atlantic	0.3791		0.2380		0.2804	
East South Central	0.1280		0.9452		0.9015	
West South Central	0.3448		0.0185		0.1426	
West	0.6013		0.1678		0.9676	
Mountain	0.4491		0.9518		0.2842	
Pacific	0.8360		0.1339		0.6286	
COUNTY TYPE						
Large Metro	0.8114		0.6489		0.8225	
Small Metro	0.5495		0.3448		0.4308	
250K – 1 Mil. Pop.	0.2231		0.3676		0.6903	
< 250K Pop.	0.5295		0.7691		0.0553	
Nonmetro	0.2306		0.4189		0.9435	
Urbanized	0.3165		0.2345		0.9672	
Less Urbanized	0.4368		0.8372		0.9347	
Completely Rural	0.9656		0.6271		0.5179	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.