

Table 1.88C Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2009 and 2010

Drug or Method of Administration	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
HALLUCINOGENS	665	689	43	44	133	134	644	654
PCP (Angel Dust, Phencyclidine)	297	321	13	11	35	34	292	317
LSD (Acid)	562	562	20	21	78	79	556	550
Peyote	279	331	13	8	28	29	275	328
Mescaline	361	382	6	8	29	32	358	377
Psilocybin (Mushrooms)	503	543	27	27	107	105	488	522
Ecstasy (MDMA)	370	429	34	35	103	114	351	396
INHALANTS	473	473	66	61	94	90	463	458
Amyl Nitrite, Poppers, Locker Room Odorizers, or Rush	305	314	28	25	38	42	301	313
Correction Fluid, Degreaser, or Cleaning Fluid	124	147	31	29	30	29	117	142
Gasoline or Lighter Fluid	188	186	40	34	35	38	182	179
Glue, Shoe Polish, or Toluene	176	233	37	39	33	37	168	227
Halothane, Ether, or Other Anesthetics	110	121	10	14	24	25	107	118
Lacquer Thinner or Other Paint Solvents	121	156	25	24	27	31	114	153
Lighter Gases (Butane, Propane)	74	80	21	17	19	22	70	74
Nitrous Oxide or Whippits	366	352	24	25	78	75	355	341
Spray Paints	108	150	37	31	28	37	98	144
Other Aerosol Sprays	132	130	32	28	38	40	124	121
NEEDLE USE^{1,2,3}	217	235	10	8	32	30	215	232
Heroin Needle Use ^{1,2}	137	158	6	6	22	24	136	156
Cocaine Needle Use ^{1,2}	163	208	5	*	19	20	162	206
Stimulant Needle Use ^{1,2,3}	154	172	6	4	21	18	152	170
Methamphetamine Needle Use ^{1,2,3}	144	164	5	4	19	17	143	161
HEROIN	220	248	9	9	37	41	217	243
Smoke Heroin ²	135	149	7	7	25	25	131	145
Sniff or Snort Heroin ²	174	218	6	5	31	33	170	215

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

³ Estimates of Stimulant Needle Use, Methamphetamine Needle Use, and the Needle Use summary variable include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.88D Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: Standard Errors of Percentages, 2009 and 2010

Drug or Method of Administration	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
HALLUCINOGENS	0.26	0.27	0.18	0.18	0.40	0.39	0.33	0.33
PCP (Angel Dust, Phencyclidine)	0.12	0.13	0.05	0.05	0.11	0.10	0.15	0.16
LSD (Acid)	0.22	0.22	0.08	0.09	0.23	0.23	0.29	0.28
Peyote	0.11	0.13	0.05	0.03	0.08	0.08	0.14	0.17
Mescaline	0.14	0.15	0.02	0.03	0.09	0.09	0.18	0.19
Psilocybin (Mushrooms)	0.20	0.21	0.11	0.11	0.32	0.31	0.25	0.27
Ecstasy (MDMA)	0.15	0.17	0.14	0.14	0.31	0.34	0.18	0.20
INHALANTS	0.19	0.19	0.27	0.25	0.28	0.26	0.24	0.23
Amyl Nitrite, Poppers, Locker Room Odorizers, or Rush	0.12	0.12	0.11	0.10	0.11	0.12	0.16	0.16
Correction Fluid, Degreaser, or Cleaning Fluid	0.05	0.06	0.13	0.12	0.09	0.09	0.06	0.07
Gasoline or Lighter Fluid	0.07	0.07	0.16	0.14	0.10	0.11	0.09	0.09
Glue, Shoe Polish, or Toluene	0.07	0.09	0.15	0.16	0.10	0.11	0.09	0.12
Halothane, Ether, or Other Anesthetics	0.04	0.05	0.04	0.06	0.07	0.07	0.06	0.06
Lacquer Thinner or Other Paint Solvents	0.05	0.06	0.10	0.10	0.08	0.09	0.06	0.08
Lighter Gases (Butane, Propane)	0.03	0.03	0.08	0.07	0.06	0.07	0.04	0.04
Nitrous Oxide or Whippits	0.15	0.14	0.10	0.10	0.23	0.22	0.18	0.17
Spray Paints	0.04	0.06	0.15	0.13	0.08	0.11	0.05	0.07
Other Aerosol Sprays	0.05	0.05	0.13	0.11	0.11	0.12	0.06	0.06
NEEDLE USE^{1,2,3}	0.09	0.09	0.04	0.03	0.10	0.09	0.11	0.12
Heroin Needle Use ^{1,2}	0.05	0.06	0.03	0.02	0.07	0.07	0.07	0.08
Cocaine Needle Use ^{1,2}	0.06	0.08	0.02	*	0.06	0.06	0.08	0.11
Stimulant Needle Use ^{1,2,3}	0.06	0.07	0.03	0.02	0.06	0.05	0.08	0.09
Methamphetamine Needle Use ^{1,2,3}	0.06	0.06	0.02	0.01	0.06	0.05	0.07	0.08
HEROIN	0.09	0.10	0.04	0.04	0.11	0.12	0.11	0.12
Smoke Heroin ²	0.05	0.06	0.03	0.03	0.08	0.07	0.07	0.07
Sniff or Snort Heroin ²	0.07	0.09	0.02	0.02	0.09	0.10	0.09	0.11

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

³ Estimates of Stimulant Needle Use, Methamphetamine Needle Use, and the Needle Use summary variable include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.88P Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime, by Age Group: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Drug or Method of Administration	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
HALLUCINOGENS	0.9606		0.9617		0.2886		0.9163	
PCP (Angel Dust, Phencyclidine)	0.9523		0.2169		0.4484		0.9516	
LSD (Acid)	0.5338		0.5906		0.1628		0.6482	
Peyote	0.2081		0.1849		0.6302		0.1768	
Mescaline	0.5087		0.1295		0.6686		0.5580	
Psilocybin (Mushrooms)	0.7880		0.3289		0.5406		0.8768	
Ecstasy (MDMA)	0.0045		0.2818		0.9335		0.0036	
INHALANTS	0.1639		0.0050		0.0365		0.5054	
Amyl Nitrite, Poppers, Locker Room Odorizers, or Rush	0.5178		0.3946		0.6424		0.5153	
Correction Fluid, Degreaser, or Cleaning Fluid	0.5078		0.2870		0.7393		0.3731	
Gasoline or Lighter Fluid	0.1331		0.0114		0.5280		0.3888	
Glue, Shoe Polish, or Toluene	0.1346		0.9201		0.8232		0.1237	
Halothane, Ether, or Other Anesthetics	0.4397		0.7355		0.4900		0.4909	
Lacquer Thinner or Other Paint Solvents	0.5912		0.0817		0.9539		0.3754	
Lighter Gases (Butane, Propane)	0.3488		0.0136		0.9200		0.7686	
Nitrous Oxide or Whippits	0.6002		0.9065		0.0334		0.8870	
Spray Paints	0.6401		0.0017		0.2720		0.2574	
Other Aerosol Sprays	0.1697		0.0810		0.2759		0.4975	
NEEDLE USE^{1,2,3}	0.5820		0.3512		0.1721		0.7339	
Heroin Needle Use ^{1,2}	0.2493		0.7832		0.8071		0.2627	
Cocaine Needle Use ^{1,2}	0.6244		*		0.9743		0.5824	
Stimulant Needle Use ^{1,2,3}	0.6006		0.2755		0.0514		0.7957	
Methamphetamine Needle Use ^{1,2,3}	0.8431		0.3054		0.1035		0.9804	
HEROIN	0.2269		0.4295		0.6426		0.2416	
Smoke Heroin ²	0.6491		0.4583		0.6954		0.6754	
Sniff or Snort Heroin ²	0.0575		0.1904		0.6103		0.0416	

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "noncore" module of the NSDUH questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NSDUH data are not edited for consistency across the "core" and "noncore" modules of the interview.

³ Estimates of Stimulant Needle Use, Methamphetamine Needle Use, and the Needle Use summary variable include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.89C Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2009 and 2010

Pain Reliever	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
Darvocet [®] , Darvon [®] , or Tylenol [®] with Codeine	437	480	45	45	93	91	419	467
Percocet [®] , Percodan [®] , or Tylox [®]	369	397	29	28	91	91	351	372
Vicodin [®] , Lortab [®] , or Lorcet [®]	446	488	46	43	127	122	413	456
Codeine	258	288	25	28	67	74	242	275
Demerol [®]	167	189	10	9	33	33	163	184
Dilaudid [®]	100	105	6	5	21	28	97	101
Fioricet [®]	64	58	5	5	14	13	62	56
Fiorinal [®]	69	78	3	5	8	6	68	77
Hydrocodone	299	296	27	32	88	89	285	270
Methadone	126	131	13	17	37	48	120	119
Morphine	134	176	17	18	43	55	122	163
OxyContin [®]	222	245	27	28	70	75	204	224
Phenaphen [®] with Codeine	77	86	9	11	20	21	74	83
Propoxyphene	45	31	6	8	16	12	41	27
SK-65 [®]	24	18	3	5	6	11	23	13
Stadol [®]	18	24	1	2	9	6	16	24
Talacen [®]	34	15	4	3	4	6	33	14
Talwin [®]	64	67	4	4	4	5	64	66
Talwin [®] NX	21	20	2	3	6	7	20	18
Tramadol	103	110	12	11	36	38	96	100
Ultram [®]	93	102	9	9	28	26	88	100
SELECTED GROUPS OF DRUGS¹								
Propoxyphene or Codeine Products ²	472	489	48	48	101	105	451	474
Oxycodone Products ³	392	412	36	35	99	102	366	382
Hydrocodone Products ⁴	471	499	47	46	131	123	438	468
Tramadol Products ⁵	118	134	15	13	42	40	110	124

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Pain Relievers module but fall into this category.

² Includes Darvocet[®], Darvon[®] or Tylenol[®] with Codeine, codeine, Phenaphen[®] with Codeine, propoxyphene, and SK-65[®].

³ Includes Percocet[®], Percodan[®] or Tylox[®], and OxyContin[®].

⁴ Includes Vicodin[®], Lortab[®], or Lorcet[®], and hydrocodone.

⁵ Includes tramadol and Ultram[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.89D Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: Standard Errors of Percentages, 2009 and 2010

Pain Reliever	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
Darvocet [®] , Darvon [®] , or Tylenol [®] with Codeine	0.17	0.19	0.18	0.19	0.28	0.27	0.22	0.24
Percocet [®] , Percodan [®] , or Tylox [®]	0.15	0.16	0.12	0.12	0.27	0.27	0.18	0.19
Vicodin [®] , Lortab [®] , or Lorcet [®]	0.18	0.19	0.18	0.18	0.38	0.36	0.21	0.23
Codeine	0.10	0.11	0.10	0.12	0.20	0.22	0.12	0.14
Demerol [®]	0.07	0.07	0.04	0.04	0.10	0.10	0.08	0.09
Dilaudid [®]	0.04	0.04	0.03	0.02	0.06	0.08	0.05	0.05
Fioricet [®]	0.03	0.02	0.02	0.02	0.04	0.04	0.03	0.03
Fiorinal [®]	0.03	0.03	0.01	0.02	0.02	0.02	0.04	0.04
Hydrocodone	0.12	0.12	0.11	0.13	0.26	0.26	0.15	0.14
Methadone	0.05	0.05	0.05	0.07	0.11	0.14	0.06	0.06
Morphine	0.05	0.07	0.07	0.07	0.13	0.16	0.06	0.08
OxyContin [®]	0.09	0.10	0.11	0.12	0.21	0.22	0.11	0.11
Phenaphen [®] with Codeine	0.03	0.03	0.04	0.04	0.06	0.06	0.04	0.04
Propoxyphene	0.02	0.01	0.02	0.03	0.05	0.04	0.02	0.01
SK-65 [®]	0.01	0.01	0.01	0.02	0.02	0.03	0.01	0.01
Stadol [®]	0.01	0.01	0.00	0.01	0.03	0.02	0.01	0.01
Talacen [®]	0.01	0.01	0.01	0.01	0.01	0.02	0.02	0.01
Talwin [®]	0.03	0.03	0.02	0.02	0.01	0.01	0.03	0.03
Talwin [®] NX	0.01	0.01	0.01	0.01	0.02	0.02	0.01	0.01
Tramadol	0.04	0.04	0.05	0.05	0.11	0.11	0.05	0.05
Ultram [®]	0.04	0.04	0.04	0.04	0.08	0.08	0.05	0.05
SELECTED GROUPS OF DRUGS¹								
Propoxyphene or Codeine Products ²	0.19	0.19	0.20	0.20	0.30	0.31	0.23	0.24
Oxycodone Products ³	0.16	0.16	0.15	0.15	0.29	0.30	0.19	0.20
Hydrocodone Products ⁴	0.19	0.20	0.19	0.19	0.39	0.36	0.23	0.24
Tramadol Products ⁵	0.05	0.05	0.06	0.05	0.13	0.12	0.06	0.06

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Pain Relievers module but fall into this category.

² Includes Darvocet[®], Darvon[®] or Tylenol[®] with Codeine, codeine, Phenaphen[®] with Codeine, propoxyphene, and SK-65[®].

³ Includes Percocet[®], Percodan[®] or Tylox[®], and OxyContin[®].

⁴ Includes Vicodin[®], Lortab[®], or Lorcet[®], and hydrocodone.

⁵ Includes tramadol and Ultram[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.89P Nonmedical Use of Specific Pain Relievers in Lifetime, by Age Group: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Pain Reliever	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
Darvocet [®] , Darvon [®] , or Tylenol [®] with Codeine	0.0651		0.2199		0.0144		0.1885	
Percocet [®] , Percodan [®] , or Tylox [®]	0.8721		0.2579		0.4440		0.6903	
Vicodin [®] , Lortab [®] , or Lorcet [®]	0.6988		0.0806		0.2101		0.3870	
Codeine	0.3099		0.3887		0.1916		0.4068	
Demerol [®]	0.5398		0.6152		0.6916		0.4804	
Dilaudid [®]	0.9751		0.4977		0.0108		0.5297	
Fioricet [®]	0.4878		0.9140		0.2789		0.6466	
Fiorinal [®]	0.4013		0.5826		0.3416		0.4317	
Hydrocodone	0.2179		0.0866		0.3517		0.4311	
Methadone	0.4233		0.2358		0.0513		0.9785	
Morphine	0.2468		0.3714		0.8948		0.2116	
OxyContin [®]	0.4505		0.8487		0.8880		0.4554	
Phenaphen [®] with Codeine	0.9388		0.6732		0.9564		0.9002	
Propoxyphene	0.0191		0.1963		0.0614		0.0394	
SK-65 [®]	0.8833		0.1659		0.3533		0.3591	
Stadol [®]	0.6784		0.5017		0.4100		0.4890	
Talacen [®]	0.2864		0.5454		0.1756		0.2009	
Talwin [®]	0.6496		0.9867		0.7849		0.6620	
Talwin [®] NX	0.5922		0.7422		0.5031		0.3979	
Tramadol	0.0942		0.8089		0.5720		0.1072	
Ultram [®]	0.3247		0.3680		0.0232		0.7372	
SELECTED GROUPS OF DRUGS¹								
Propoxyphene or Codeine Products ²	0.0264		0.4689		0.0242		0.0783	
Oxycodone Products ³	0.7724		0.3526		0.9000		0.7080	
Hydrocodone Products ⁴	0.5795		0.2192		0.4731		0.4025	
Tramadol Products ⁵	0.6921		0.4442		0.1585		0.3369	

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Pain Relievers module but fall into this category.

² Includes Darvocet[®], Darvon[®] or Tylenol[®] with Codeine, codeine, Phenaphen[®] with Codeine, propoxyphene, and SK-65[®].

³ Includes Percocet[®], Percodan[®] or Tylox[®], and OxyContin[®].

⁴ Includes Vicodin[®], Lortab[®], or Lorcet[®], and hydrocodone.

⁵ Includes tramadol and Ultram[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.90C Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2009 and 2010

Tranquilizer	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
Klonopin® or Clonazepam	220	194	16	19	60	59	206	187
Xanax®, Alprazolam, Ativan®, or Lorazepam	355	361	27	33	91	96	337	343
Valium® or Diazepam	384	411	17	21	71	75	375	403
Atarax®	58	33	6	6	8	10	57	30
BuSpar®	53	66	7	5	16	19	49	65
Equanil®	29	28	3	4	5	4	28	27
Flexeril®	140	158	8	10	35	35	134	151
Librium®	123	117	4	4	11	12	122	117
Limbitrol®	20	28	5	4	11	5	16	27
Meprobamate	41	19	5	4	7	8	41	17
Miltown®	36	23	2	4	3	4	36	22
Rohypnol®	44	60	7	4	13	11	43	59
Serax®	17	20	4	4	7	9	15	17
Soma®	155	182	14	16	49	56	147	168
Tranxene®	35	23	6	*	6	7	34	22
Vistaril®	45	60	4	6	10	11	44	58
SELECTED GROUPS OF DRUGS¹								
Benzodiazepines ²	476	481	31	35	101	103	455	466
Meprobamate Products ³	62	40	6	6	8	9	61	38
Muscle Relaxants ⁴	193	222	16	19	56	60	182	207

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Tranquilizers module but fall into this category.

² Includes Klonopin® or clonazepam, Xanax®, alprazolam, Ativan® or lorazepam, Valium® or diazepam, Librium®, Limbitrol®, Rohypnol®, Serax®, and Tranxene®.

³ Includes Equanil®, meprobamate, and Miltown®.

⁴ Includes Flexeril® and Soma®.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.90D Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: Standard Errors of Percentages, 2009 and 2010

Tranquilizer	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
Klonopin® or Clonazepam	0.09	0.08	0.06	0.08	0.18	0.17	0.11	0.10
Xanax®, Alprazolam, Ativan®, or Lorazepam	0.14	0.14	0.11	0.13	0.27	0.28	0.17	0.18
Valium® or Diazepam	0.15	0.16	0.07	0.08	0.21	0.22	0.19	0.21
Atarax®	0.02	0.01	0.02	0.03	0.03	0.03	0.03	0.02
BuSpar®	0.02	0.03	0.03	0.02	0.05	0.06	0.03	0.03
Equanil®	0.01	0.01	0.01	0.02	0.01	0.01	0.01	0.01
Flexeril®	0.06	0.06	0.03	0.04	0.10	0.10	0.07	0.08
Librium®	0.05	0.05	0.02	0.02	0.03	0.04	0.06	0.06
Limbitrol®	0.01	0.01	0.02	0.02	0.03	0.02	0.01	0.01
Meprobamate	0.02	0.01	0.02	0.02	0.02	0.02	0.02	0.01
Miltown®	0.01	0.01	0.01	0.02	0.01	0.01	0.02	0.01
Rohypnol®	0.02	0.02	0.03	0.02	0.04	0.03	0.02	0.03
Serax®	0.01	0.01	0.01	0.02	0.02	0.03	0.01	0.01
Soma®	0.06	0.07	0.06	0.07	0.15	0.16	0.08	0.09
Tranxene®	0.01	0.01	0.02	*	0.02	0.02	0.02	0.01
Vistaril®	0.02	0.02	0.02	0.02	0.03	0.03	0.02	0.03
SELECTED GROUPS OF DRUGS¹								
Benzodiazepines ²	0.19	0.19	0.13	0.14	0.30	0.30	0.23	0.24
Meprobamate Products ³	0.02	0.02	0.02	0.02	0.02	0.03	0.03	0.02
Muscle Relaxants ⁴	0.08	0.09	0.06	0.08	0.17	0.18	0.09	0.11

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Tranquilizers module but fall into this category.

² Includes Klonopin® or clonazepam, Xanax®, alprazolam, Ativan® or lorazepam, Valium® or diazepam, Librium®, Limbitrol®, Rohypnol®, Serax®, and Tranxene®.

³ Includes Equanil®, meprobamate, and Miltown®.

⁴ Includes Flexeril® and Soma®.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.90P Nonmedical Use of Specific Tranquilizers in Lifetime, by Age Group: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Tranquilizer	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
Klonopin® or Clonazepam	0.9176		0.1693		0.7414		0.8697	
Xanax®, Alprazolam, Ativan®, or Lorazepam	0.1623		0.1486		0.1575		0.3684	
Valium® or Diazepam	0.9518		0.7972		0.7275		0.9412	
Atarax®	0.0576		0.6827		0.6348		0.0348	
BuSpar®	0.6433		0.1333		0.8949		0.5634	
Equanil®	0.7034		0.5168		0.5625		0.7080	
Flexeril®	0.4649		0.3767		0.3384		0.3617	
Librium®	0.6274		0.8357		0.5332		0.5773	
Limbitrol®	0.7543		0.4009		0.1159		0.2621	
Meprobamate	0.3492		0.7028		0.6980		0.3124	
Miltown®	0.6893		0.1942		0.7893		0.5614	
Rohypnol®	0.6119		0.1803		0.6632		0.4483	
Serax®	0.6907		0.7148		0.7811		0.6101	
Soma®	0.2568		0.4745		0.7389		0.2220	
Tranxene®	0.0296		*		0.8760		0.0658	
Vistaril®	0.3178		0.5095		0.5685		0.4000	
SELECTED GROUPS OF DRUGS¹								
Benzodiazepines ²	0.8680		0.5933		0.4240		0.9251	
Meprobamate Products ³	0.3339		0.7532		0.6561		0.2747	
Muscle Relaxants ⁴	0.3535		0.3007		0.4133		0.2715	

*Low precision; no estimate reported.

¹ Includes other-specify drug responses that are not asked about explicitly in the Tranquilizers module but fall into this category.

² Includes Klonopin® or clonazepam, Xanax®, alprazolam, Ativan® or lorazepam, Valium® or diazepam, Librium®, Limbitrol®, Rohypnol®, Serax®, and Tranxene®.

³ Includes Equanil®, meprobamate, and Miltown®.

⁴ Includes Flexeril® and Soma®.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.91C Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2009 and 2010

Stimulant	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
Methamphetamine, Desoxyn [®] , or Methedrine ¹	403	451	19	18	62	62	393	438
Prescription Diet Pills ²	318	361	16	15	41	43	314	359
Ritalin [®] or Methylphenidate	220	249	20	22	72	71	205	231
Cylert [®]	38	26	5	3	11	11	36	23
Dexedrine [®]	203	201	6	7	20	18	202	200
Dextroamphetamine	74	130	7	7	22	23	71	129
Didrex [®]	52	30	4	3	7	7	51	29
Eskatrol [®]	43	27	4	5	4	8	43	25
Ionamin [®]	77	54	6	2	3	3	76	54
Mazanor [®]	11	10	2	2	3	7	10	7
Obedrin-LA [®]	*	*	3	*	2	2	*	*
Plegine [®]	7	*	3	2	4	4	*	*
Preludin [®]	88	98	4	3	5	9	88	98
Sanorex [®]	26	21	3	3	6	2	25	21
Tenuate [®]	40	22	3	4	6	3	39	22
SELECTED GROUPS OF DRUGS³								
Amphetamine, Dextroamphetamine, and Phentermine Products ⁴	357	391	19	20	57	65	349	381
Mazindol Products ⁵	28	23	4	3	6	7	27	22
Methylphenidate or Dexmethylphenidate Products ⁶	220	249	20	22	72	70	205	231

*Low precision; no estimate reported.

¹ Also known as crank, crystal, ice, or speed. Estimates of Use of Methamphetamine, Desoxyn[®], or Methedrine include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Respondents were asked about their use of prescription diet pills and were given the following as examples: amphetamines, Benzedrine[®], Biphedamine[®], Fastin[®], or phentermine. However, respondents were not given an exhaustive list of examples of prescription diet pills.

³ Includes other-specify drug responses that are not asked about explicitly in the Stimulants module but fall into this category.

⁴ Includes prescription diet pills, Dexedrine[®], dextroamphetamine, and Ionamin[®].

⁵ Includes Mazanor[®] and Sanorex[®].

⁶ Includes Ritalin[®], other products containing methylphenidate, and products containing dexmethylphenidate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.91D Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: Standard Errors of Percentages, 2009 and 2010

Stimulant	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
Methamphetamine, Desoxyn [®] , or Methedrine ¹	0.16	0.18	0.08	0.08	0.18	0.18	0.20	0.22
Prescription Diet Pills ²	0.13	0.14	0.06	0.06	0.12	0.12	0.16	0.18
Ritalin [®] or Methylphenidate	0.09	0.10	0.08	0.09	0.21	0.21	0.11	0.12
Cylert [®]	0.02	0.01	0.02	0.01	0.03	0.03	0.02	0.01
Dexedrine [®]	0.08	0.08	0.02	0.03	0.06	0.05	0.10	0.10
Dextroamphetamine	0.03	0.05	0.03	0.03	0.06	0.07	0.04	0.07
Didrex [®]	0.02	0.01	0.02	0.01	0.02	0.02	0.03	0.01
Eskatrol [®]	0.02	0.01	0.02	0.02	0.01	0.02	0.02	0.01
Ionamin [®]	0.03	0.02	0.02	0.01	0.01	0.01	0.04	0.03
Mazanor [®]	0.00	0.00	0.01	0.01	0.01	0.02	0.01	0.00
Obedrin-LA [®]	*	*	0.01	*	0.01	0.01	*	*
Plegine [®]	0.00	*	0.01	0.01	0.01	0.01	*	*
Preludin [®]	0.04	0.04	0.02	0.01	0.01	0.03	0.05	0.05
Sanorex [®]	0.01	0.01	0.01	0.01	0.02	0.01	0.01	0.01
Tenuate [®]	0.02	0.01	0.01	0.02	0.02	0.01	0.02	0.01
SELECTED GROUPS OF DRUGS³								
Amphetamine, Dextroamphetamine, and Phentermine Products ⁴	0.14	0.15	0.08	0.08	0.17	0.19	0.18	0.20
Mazindol Products ⁵	0.01	0.01	0.02	0.01	0.02	0.02	0.01	0.01
Methylphenidate or Dexmethylphenidate Products ⁶	0.09	0.10	0.08	0.09	0.22	0.21	0.11	0.12

*Low precision; no estimate reported.

¹ Also known as crank, crystal, ice, or speed. Estimates of Use of Methamphetamine, Desoxyn[®], or Methedrine include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Respondents were asked about their use of prescription diet pills and were given the following as examples: amphetamines, Benzedrine[®], Biphedamine[®], Fastin[®], or phentermine. However, respondents were not given an exhaustive list of examples of prescription diet pills.

³ Includes other-specify drug responses that are not asked about explicitly in the Stimulants module but fall into this category.

⁴ Includes prescription diet pills, Dexedrine[®], dextroamphetamine, and Ionamin[®].

⁵ Includes Mazanor[®] and Sanorex[®].

⁶ Includes Ritalin[®], other products containing methylphenidate, and products containing dexmethylphenidate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.91P Nonmedical Use of Specific Stimulants in Lifetime, by Age Group: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Stimulant	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
Methamphetamine, Desoxyn [®] , or Methedrine ¹	0.8892		0.3105		0.0366		0.6405	
Prescription Diet Pills ²	0.2315		0.1923		0.8725		0.2546	
Ritalin [®] or Methylphenidate	0.9516		0.5661		0.6910		0.9658	
Cylert [®]	0.0981		0.4308		0.6453		0.0638	
Dexedrine [®]	0.0630		0.4927		0.4217		0.0735	
Dextroamphetamine	0.4421		0.9637		0.8858		0.4518	
Didrex [®]	0.0651		0.6564		0.6584		0.0775	
Eskatrol [®]	0.2914		0.7093		0.1742		0.1946	
Ionamin [®]	0.1732		0.1209		0.5717		0.2148	
Mazanor [®]	0.7246		0.6733		0.4087		0.9825	
Obedrin-LA [®]	*		*		0.6913		*	
Plegine [®]	*		0.5834		0.8623		*	
Preludin [®]	0.7829		0.4725		0.3217		0.8251	
Sanorex [®]	0.5956		0.7972		0.0283		0.9500	
Tenuate [®]	0.2289		0.5820		0.2497		0.2647	
SELECTED GROUPS OF DRUGS³								
Amphetamine, Dextroamphetamine, and Phentermine Products ⁴	0.5199		0.2077		0.3021		0.4287	
Mazindol Products ⁵	0.7465		0.3968		0.5576		0.9590	
Methylphenidate or Dexmethylphenidate Products ⁶	0.9270		0.7469		0.6953		0.9742	

*Low precision; no estimate reported.

¹ Also known as crank, crystal, ice, or speed. Estimates of Use of Methamphetamine, Desoxyn[®], or Methedrine include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 National Findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Respondents were asked about their use of prescription diet pills and were given the following as examples: amphetamines, Benzedrine[®], Biphedamine[®], Fastin[®], or phentermine. However, respondents were not given an exhaustive list of examples of prescription diet pills.

³ Includes other-specify drug responses that are not asked about explicitly in the Stimulants module but fall into this category.

⁴ Includes prescription diet pills, Dexedrine[®], dextroamphetamine, and Ionamin[®].

⁵ Includes Mazanor[®] and Sanorex[®].

⁶ Includes Ritalin[®], other products containing methylphenidate, and products containing dexmethylphenidate.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.92C Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Standard Errors of Numbers in Thousands, 2009 and 2010

Sedative	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
Methaqualone, Sopor [®] , or Quaalude [®]	295	304	6	9	20	18	295	302
Barbiturates ¹	206	204	7	6	16	13	205	204
Restoril [®] or Temazepam	128	116	10	9	22	20	125	115
Amytal [®]	54	80	3	2	9	10	53	80
Butisol [®]	25	39	3	6	4	7	24	38
Chloral Hydrate	44	40	1	3	10	6	43	39
Dalmane [®]	66	52	6	4	7	8	65	51
Halcion [®]	82	115	5	7	9	11	81	114
Phenobarbital	127	122	5	6	15	12	127	121
Placidyl [®]	75	75	1	2	3	6	75	75
Tuinal [®]	119	88	4	1	5	6	119	88
SELECTED GROUPS OF DRUGS²								
Temazepam, Flurazepam, or Triazolam ³	151	152	13	12	23	21	148	151
Any Barbiturates ⁴	229	223	9	10	23	18	228	223

*Low precision; no estimate reported.

¹ Respondents were asked about their use of barbiturates and were given the following as examples: Nembutal[®], pentobarbital, Seconal[®], secobarbital, or butalbital. However, respondents were not given an exhaustive list of examples of barbiturates.

² Includes other-specify drug responses that are not asked about explicitly in the Sedatives module but fall into this category.

³ Includes Restoril[®], temazepam, Dalmane[®], or Halcion[®].

⁴ Includes barbiturates, Amytal[®], Butisol[®], phenobarbital, and Tuinal[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.92D Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: Standard Errors of Percentages, 2009 and 2010

Sedative	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
Methaqualone, Sopor [®] , or Quaalude [®]	0.12	0.12	0.02	0.04	0.06	0.05	0.15	0.15
Barbiturates ¹	0.08	0.08	0.03	0.02	0.05	0.04	0.11	0.10
Restoril [®] or Temazepam	0.05	0.05	0.04	0.04	0.06	0.06	0.06	0.06
Amytal [®]	0.02	0.03	0.01	0.01	0.03	0.03	0.03	0.04
Butisol [®]	0.01	0.02	0.01	0.03	0.01	0.02	0.01	0.02
Chloral Hydrate	0.02	0.02	0.00	0.01	0.03	0.02	0.02	0.02
Dalmane [®]	0.03	0.02	0.03	0.02	0.02	0.02	0.03	0.03
Halcion [®]	0.03	0.05	0.02	0.03	0.03	0.03	0.04	0.06
Phenobarbital	0.05	0.05	0.02	0.03	0.05	0.03	0.07	0.06
Placidyl [®]	0.03	0.03	0.00	0.01	0.01	0.02	0.04	0.04
Tuinal [®]	0.05	0.03	0.02	0.00	0.01	0.02	0.06	0.04
SELECTED GROUPS OF DRUGS²								
Temazepam, Flurazepam, or Triazolam ³	0.06	0.06	0.05	0.05	0.07	0.06	0.08	0.08
Any Barbiturates ⁴	0.09	0.09	0.04	0.04	0.07	0.05	0.12	0.11

*Low precision; no estimate reported.

¹ Respondents were asked about their use of barbiturates and were given the following as examples: Nembutal[®], pentobarbital, Seconal[®], secobarbital, or butalbital. However, respondents were not given an exhaustive list of examples of barbiturates.

² Includes other-specify drug responses that are not asked about explicitly in the Sedatives module but fall into this category.

³ Includes Restoril[®], temazepam, Dalmane[®], or Halcion[®].

⁴ Includes barbiturates, Amytal[®], Butisol[®], phenobarbital, and Tuinal[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 1.92P Nonmedical Use of Specific Sedatives in Lifetime, by Age Group: P Values from Tests of Differences of Percentages, 2010 Versus 2009

Sedative	Total (2009)	Total (2010)	Aged 12-17 (2009)	Aged 12-17 (2010)	Aged 18-25 (2009)	Aged 18-25 (2010)	Aged 26+ (2009)	Aged 26+ (2010)
Methaqualone, Sopor [®] , or Quaalude [®]	0.0941		0.1492		0.1806		0.1022	
Barbiturates ¹	0.1237		0.4108		0.6549		0.1376	
Restoril [®] or Temazepam	0.3232		0.8773		0.6370		0.3599	
Amytal [®]	0.8445		0.2554		0.6695		0.8560	
Butisol [®]	0.6906		0.1009		0.2811		0.9930	
Chloral Hydrate	0.7182		0.2300		0.4399		0.7683	
Dalmane [®]	0.4508		0.2805		0.9838		0.5013	
Halcion [®]	0.7689		0.3648		0.3403		0.8895	
Phenobarbital	0.4646		0.3064		0.4353		0.4842	
Placidyl [®]	0.3934		0.3706		0.7161		0.3683	
Tuinal [®]	0.1580		0.0785		0.7407		0.1657	
SELECTED GROUPS OF DRUGS²								
Temazepam, Flurazepam, or Triazolam ³	0.5604		0.9363		0.7845		0.5821	
Any Barbiturates ⁴	0.1933		0.8954		0.2873		0.2247	

*Low precision; no estimate reported.

¹ Respondents were asked about their use of barbiturates and were given the following as examples: Nembutal[®], pentobarbital, Seconal[®], secobarbital, or butalbital. However, respondents were not given an exhaustive list of examples of barbiturates.

² Includes other-specify drug responses that are not asked about explicitly in the Sedatives module but fall into this category.

³ Includes Restoril[®], temazepam, Dalmane[®], or Halcion[®].

⁴ Includes barbiturates, Amytal[®], Butisol[®], phenobarbital, and Tuinal[®].

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.