

Table 2.11A Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	174,119	174,194	83,336	83,200	69,713	69,559
12	232	204	129	93	36	40
13	481	474	296	306	138	146
14	868 ^b	709	608	525	296	255
15	1,324	1,238	968	876	543	512
16	1,673	1,525	1,206	1,151	748	664
17	2,047	1,913	1,553	1,437	1,099	980
18	2,806	2,705	2,367	2,244	1,795	1,679
19	3,001	2,852	2,422	2,306	1,810	1,770
20	2,865 ^a	3,154	2,183	2,383	1,736	1,931
21	2,964	2,931	2,234	2,212	1,797	1,787
22	3,035	2,934	2,288	2,204	1,828	1,761
23	2,984	2,927	2,110	2,027	1,721	1,647
24	3,046	3,025	2,138	2,071	1,742	1,700
25	2,658	2,849	1,826	1,978	1,551	1,630
26-29	12,776	12,799	8,178	8,216	6,929	6,994
30-34	14,795	15,046	8,608	8,407	7,401	7,072
35-39	14,567	13,668	7,415	7,043	6,228	5,932
40-44	15,862	15,387	7,526	7,531	6,529	6,499
45-49	17,307	17,330	7,714	7,901	6,747	6,978
50-54	16,697	16,863	7,448	7,353	6,773	6,335
55-59	14,244	14,629	5,148	5,835	4,617	5,151
60-64	11,487	12,009	3,741	3,751	3,332	3,367
65 or Older	26,400	27,024	5,230	5,351	4,316	4,729

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.11B Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	69.1	68.7	33.1	32.8	27.7	27.4
12	6.3	5.3	3.5	2.4	1.0	1.1
13	12.5	11.6	7.7	7.5	3.6	3.6
14	21.0 ^a	18.2	14.7	13.5	7.2	6.5
15	30.1	29.8	22.0	21.1	12.3	12.3
16	39.8 ^a	36.7	28.7	27.7	17.8	16.0
17	47.0	45.1	35.7	33.9	25.3	23.1
18	56.6	57.0	47.8	47.3	36.2	35.4
19	65.7	65.1	53.0	52.6	39.6	40.4
20	69.5	69.1	53.0	52.2	42.1	42.3
21	71.5	68.7	53.9	51.9	43.3	41.9
22	75.2 ^a	71.3	56.7	53.6	45.3	42.8
23	74.4	72.8	52.6	50.4	42.9	41.0
24	74.0	74.6	51.9	51.1	42.3	41.9
25	73.4	72.6	50.4	50.4	42.8	41.6
26-29	76.0	76.8	48.7	49.3	41.2	42.0
30-34	76.2	75.7	44.3	42.3	38.1	35.6
35-39	73.3	70.8	37.3	36.5	31.3	30.7
40-44	76.1	75.1	36.1	36.7	31.3	31.7
45-49	77.1	77.9	34.4	35.5	30.1	31.4
50-54	75.0	76.6	33.4	33.4	30.4	28.8
55-59	76.9	75.9	27.8	30.3	24.9	26.7
60-64	74.2	73.1	24.2	22.8	21.5	20.5
65 or Older	69.5	69.4	13.8	13.7	11.4	12.1

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.12A Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	162,681	162,762	69,165	68,458	58,661	58,256
12	181	150	98	65	18	25
13	341	369	208	243	86	116
14	694 ^b	573	480 ^a	398	230	204
15	1,074	1,002	741	687	411	389
16	1,405 ^a	1,227	941	882	585	513
17	1,763	1,627	1,221	1,164	860	784
18	2,489	2,329	1,990	1,822	1,453	1,331
19	2,668	2,517	2,062	1,929	1,554	1,482
20	2,571	2,792	1,897	1,995	1,503	1,559
21	2,737	2,664	1,951	1,864	1,549	1,527
22	2,836	2,717	1,962	1,878	1,569	1,512
23	2,759	2,672	1,847	1,726	1,503	1,383
24	2,841	2,803	1,868	1,816	1,539	1,490
25	2,496	2,661	1,608	1,645	1,359	1,381
26-29	11,915	12,065	7,102	6,968	6,114	6,127
30-34	13,909	14,194	7,206	7,099	6,201	6,140
35-39	13,475	12,885	5,970	5,654	5,108	4,762
40-44	14,725	14,284	5,842	6,114	5,225	5,441
45-49	16,154	16,271	6,605	6,384	5,946	5,821
50-54	15,955	16,066	6,058	6,137	5,738	5,556
55-59	13,516	13,909	4,460	4,887	3,999	4,386
60-64	11,101	11,381	3,054	3,071	2,744	2,768
65 or Older	25,078	25,605	3,992	4,031	3,365	3,558

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.12B Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	64.6	64.2	27.5	27.0	23.3	23.0
12	4.9	3.9	2.7	1.7	0.5	0.6
13	8.9	9.0	5.4	6.0	2.2	2.8
14	16.8	14.7	11.6	10.2	5.6	5.2
15	24.4	24.1	16.8	16.5	9.3	9.4
16	33.4 ^b	29.5	22.4	21.2	13.9	12.3
17	40.5	38.3	28.1	27.4	19.8	18.5
18	50.2	49.1	40.2	38.4	29.3	28.0
19	58.4	57.4	45.1	44.0	34.0	33.8
20	62.4	61.2	46.0	43.7	36.5	34.1
21	66.0	62.5	47.1	43.7	37.4	35.8
22	70.3 ^b	66.0	48.6	45.6	38.9	36.8
23	68.7	66.5	46.0	42.9	37.5	34.4
24	69.0	69.1	45.4	44.8	37.4	36.8
25	68.9	67.8	44.4	41.9	37.5	35.2
26-29	70.9	72.4	42.3	41.8	36.4	36.8
30-34	71.7	71.5	37.1	35.7	31.9	30.9
35-39	67.8	66.7	30.0	29.3	25.7	24.7
40-44	70.6	69.7	28.0	29.8	25.1	26.5
45-49	72.0	73.2	29.4	28.7	26.5	26.2
50-54	71.6	73.0	27.2	27.9	25.8	25.2
55-59	72.9	72.2	24.1	25.4	21.6	22.8
60-64	71.7	69.3	19.7	18.7	17.7	16.9
65 or Older	66.0	65.8	10.5	10.4	8.9	9.1

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.13A Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	44,804	44,916	12,063	12,334	8,559	8,855
12	53	53	26	26	8	9
13	124	123	71	82	32	39
14	234	191	145	149	75	55
15	347	344	245	247	102	108
16	493	465	347	316	148	152
17	628	589	430	382	211	198
18	868	857	582	627	325	373
19	917	887	522	531	298	302
20	870	971	492	493	296	332
21	882	899	416	428	250	231
22	988	902	410	416	260	265
23	858	896	371	365	222	233
24	822	842	299	354	188	225
25	830	875	279	341	214	224
26-29	3,577	3,765	1,146	1,183	692	867
30-34	4,790	4,620	1,372	1,228	1,096	897
35-39	5,087	5,089	1,162	1,315	953	1,160
40-44	5,272	5,145	1,281 ^b	903	1,132 ^b	767
45-49	4,602	4,653	691	915	557	738
50-54	3,821	3,650	616	488	590	389
55-59	2,662	2,610	303	496	243	451
60-64	2,029	2,006	293	342	234	235
65 or Older	4,050	4,484	561	707	432	605

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.13B Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	17.8	17.7	4.8	4.9	3.4	3.5
12	1.4	1.4	0.7	0.7	0.2	0.2
13	3.2	3.0	1.8	2.0	0.8	1.0
14	5.7	4.9	3.5	3.8	1.8	1.4
15	7.9	8.3	5.6	5.9	2.3	2.6
16	11.7	11.2	8.3	7.6	3.5	3.7
17	14.4	13.9	9.9	9.0	4.9	4.7
18	17.5	18.1	11.7	13.2	6.6	7.9
19	20.1	20.2	11.4	12.1	6.5	6.9
20	21.1	21.3	11.9	10.8	7.2	7.3
21	21.3	21.1	10.0	10.0	6.0	5.4
22	24.5	21.9	10.2	10.1	6.4	6.4
23	21.4	22.3	9.3	9.1	5.5	5.8
24	20.0	20.8	7.3	8.7	4.6	5.5
25	22.9	22.3	7.7	8.7	5.9	5.7
26-29	21.3	22.6	6.8	7.1	4.1	5.2
30-34	24.7	23.3	7.1	6.2	5.6	4.5
35-39	25.6	26.4	5.9	6.8	4.8	6.0
40-44	25.3	25.1	6.1 ^a	4.4	5.4 ^b	3.7
45-49	20.5	20.9	3.1	4.1	2.5	3.3
50-54	17.2	16.6	2.8	2.2	2.6	1.8
55-59	14.4	13.5	1.6	2.6	1.3	2.3
60-64	13.1	12.2	1.9	2.1	1.5	1.4
65 or Older	10.7	11.5	1.5	1.8	1.1	1.6

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.14A Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	90,115	88,334	26,402	25,883	13,269	13,177
12	67 ^a	34	36	17	10	9
13	205 ^a	149	115	91	44	33
14	352 ^a	272	234	195	81	59
15	625	541	478 ^b	373	185	158
16	795	737	564	529	285 ^b	205
17	1,019	938	771 ^a	662	374	318
18	1,660	1,586	1,285	1,261	657	641
19	1,730	1,663	1,259	1,121	554	559
20	1,719	1,772	1,051	1,053	564	553
21	1,787	1,792	1,042	999	506	481
22	1,853	1,764	907	916	441	426
23	1,833	1,806	847	804	438	400
24	1,820	1,748	805	748	355	378
25	1,599	1,719	630 ^a	750	322	376
26-29	7,773	7,785	2,917	2,843	1,203	1,278
30-34	8,630	8,600	2,647	2,392	1,474	1,205
35-39	7,850	7,425	2,029	2,103	1,038	1,065
40-44	7,635	8,333	1,877	2,121	984	1,077
45-49	8,710	8,240	1,834	1,970	929	1,156
50-54	8,571	7,829	1,977	1,692	1,067	889
55-59	7,235	7,013	1,305	1,447	729	761
60-64	5,392	5,527	928	817	588	530
65 or Older	11,257	11,059	865	979	440	619

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.14B Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	35.8	34.8	10.5	10.2	5.3	5.2
12	1.8 ^a	0.9	1.0 ^a	0.4	0.3	0.2
13	5.3 ^a	3.7	3.0	2.2	1.2	0.8
14	8.5	7.0	5.6	5.0	2.0	1.5
15	14.2	13.0	10.9 ^a	9.0	4.2	3.8
16	18.9	17.7	13.4	12.7	6.8 ^b	4.9
17	23.4	22.1	17.7	15.6	8.6	7.5
18	33.5	33.4	25.9	26.6	13.3	13.5
19	37.8	37.9	27.5	25.6	12.1	12.8
20	41.7	38.8	25.5	23.1	13.7	12.1
21	43.1	42.0	25.1	23.4	12.2	11.3
22	45.9	42.9	22.5	22.3	10.9	10.4
23	45.7	44.9	21.1	20.0	10.9	10.0
24	44.2	43.1	19.5	18.5	8.6	9.3
25	44.1	43.8	17.4	19.1	8.9	9.6
26-29	46.3	46.7	17.4	17.1	7.2	7.7
30-34	44.5	43.3	13.6	12.0	7.6	6.1
35-39	39.5	38.4	10.2	10.9	5.2	5.5
40-44	36.6 ^a	40.7	9.0	10.3	4.7	5.3
45-49	38.8	37.1	8.2	8.9	4.1	5.2
50-54	38.5	35.6	8.9	7.7	4.8	4.0
55-59	39.0	36.4	7.0	7.5	3.9	3.9
60-64	34.8	33.7	6.0	5.0	3.8	3.2
65 or Older	29.6	28.4	2.3	2.5	1.2	1.6

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.15A Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	208,545	209,295	168,141	168,396	130,621	131,342
12	342 ^a	270	200	168	88	60
13	708	709	496	511	172	188
14	1,290 ^b	1,105	961 ^a	832	367	351
15	1,961 ^a	1,784	1,594	1,470	745 ^a	643
16	2,359	2,170	1,900	1,813	942	876
17	2,723	2,536	2,315	2,154	1,308	1,190
18	3,601	3,430	3,202	3,036	2,101	1,974
19	3,650	3,422	3,278	3,074	2,342	2,167
20	3,468 ^a	3,796	3,121 ^a	3,428	2,337	2,549
21	3,722	3,857	3,573	3,679	2,964	3,052
22	3,732	3,708	3,533	3,470	2,890	2,837
23	3,617	3,692	3,305	3,449	2,798	2,852
24	3,798	3,707	3,504	3,379	2,884	2,786
25	3,222 ^b	3,594	2,941 ^b	3,301	2,451 ^a	2,745
26-29	15,148	15,171	13,658	13,438	11,156	10,883
30-34	17,611	17,932	15,297	15,537	12,126	12,699
35-39	17,813	17,148	14,938	14,660	11,813	11,684
40-44	19,014	18,512	15,822	15,796	12,742	12,521
45-49	20,388	20,454	16,692	16,391	13,226	13,098
50-54	20,082	19,812	15,544	15,745	12,223	12,677
55-59	16,818	17,466	12,941	12,830	10,306	10,160
60-64	13,793	14,474	9,946	10,561	7,787	8,469
65 or Older	29,685	30,547	19,380	19,673	14,852	14,879

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.15B Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2009 and 2010

Age Category	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	82.8	82.5	66.8	66.4	51.9	51.8
12	9.3 ^b	7.1	5.4	4.4	2.4	1.6
13	18.4	17.4	12.9	12.5	4.5	4.6
14	31.2 ^a	28.4	23.2	21.4	8.9	9.0
15	44.6	42.9	36.2	35.4	16.9	15.5
16	56.1 ^b	52.2	45.2	43.6	22.4	21.1
17	62.6	59.8	53.2	50.8	30.1	28.0
18	72.7	72.3	64.6	64.0	42.4	41.6
19	79.9	78.1	71.7	70.1	51.2	49.4
20	84.2	83.2	75.8	75.1	56.7	55.8
21	89.8	90.5	86.2	86.3	71.5	71.6
22	92.5 ^a	90.1	87.5 ^a	84.3	71.6	68.9
23	90.1	91.8	82.3 ^b	85.8	69.7	70.9
24	92.3	91.4	85.1	83.4	70.1	68.7
25	88.9 ^a	91.6	81.2	84.1	67.7	70.0
26-29	90.1	91.1	81.3	80.7	66.4	65.3
30-34	90.7	90.3	78.8	78.2	62.5	63.9
35-39	89.7	88.8	75.2	75.9	59.5	60.5
40-44	91.2	90.3	75.9	77.1	61.1	61.1
45-49	90.8	92.0	74.4	73.7	58.9	58.9
50-54	90.2	90.0	69.8	71.5	54.9	57.6
55-59	90.8	90.7	69.8	66.6	55.6	52.7
60-64	89.1	88.1	64.3	64.3	50.3	51.6
65 or Older	78.2	78.5	51.0	50.5	39.1	38.2

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.16A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Numbers in Thousands, 2009 and 2010

Age Category	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	130,621	131,342	59,561	58,550	17,129	16,931
12	88	60	36 ^a	14	3	2
13	172	188	86	67	12	8
14	367	351	179	161	19	25
15	745 ^a	643	419	381	98	71
16	942	876	552	516	120	114
17	1,308	1,190	904 ^a	772	265 ^b	189
18	2,101	1,974	1,505	1,350	400	391
19	2,342	2,167	1,631	1,505	579	535
20	2,337	2,549	1,601	1,702	562	623
21	2,964	3,052	1,999	2,043	691	755
22	2,890	2,837	2,039	1,870	718	669
23	2,798	2,852	1,845	1,836	623	584
24	2,884	2,786	1,848	1,804	579	531
25	2,451 ^a	2,745	1,530 ^a	1,720	460	551
26-29	11,156	10,883	6,517	6,435	1,809	1,902
30-34	12,126	12,699	6,639	6,878	1,849	1,847
35-39	11,813	11,684	5,429	5,516	1,464	1,505
40-44	12,742	12,521	5,623	5,426	1,415	1,527
45-49	13,226	13,098	5,443	5,548	1,512	1,609
50-54	12,223	12,677	4,533	4,537	1,418	1,317
55-59	10,306	10,160	3,549	3,258	1,094	1,001
60-64	7,787	8,469	1,937	2,268	588	537
65 or Older	14,852	14,879	3,717	2,942	852	636

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.16B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Percentages, 2009 and 2010

Age Category	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	51.9	51.8	23.7	23.1	6.8	6.7
12	2.4	1.6	1.0 ^a	0.4	0.1	0.0
13	4.5	4.6	2.2	1.6	0.3	0.2
14	8.9	9.0	4.3	4.1	0.5	0.6
15	16.9	15.5	9.5	9.2	2.2	1.7
16	22.4	21.1	13.1	12.4	2.8	2.7
17	30.1	28.0	20.8 ^a	18.2	6.1 ^a	4.5
18	42.4	41.6	30.4	28.5	8.1	8.2
19	51.2	49.4	35.7	34.3	12.7	12.2
20	56.7	55.8	38.9	37.3	13.6	13.7
21	71.5	71.6	48.2	47.9	16.7	17.7
22	71.6	68.9	50.5 ^b	45.5	17.8	16.3
23	69.7	70.9	46.0	45.7	15.5	14.5
24	70.1	68.7	44.9	44.5	14.1	13.1
25	67.7	70.0	42.2	43.8	12.7	14.0
26-29	66.4	65.3	38.8	38.6	10.8	11.4
30-34	62.5	63.9	34.2	34.6	9.5	9.3
35-39	59.5	60.5	27.3	28.6	7.4	7.8
40-44	61.1	61.1	27.0	26.5	6.8	7.5
45-49	58.9	58.9	24.2	25.0	6.7	7.2
50-54	54.9	57.6	20.4	20.6	6.4	6.0
55-59	55.6	52.7	19.2	16.9	5.9	5.2
60-64	50.3	51.6	12.5	13.8	3.8	3.3
65 or Older	39.1	38.2	9.8 ^a	7.6	2.2	1.6

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.