

**Table 2.1A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Numbers in Thousands, 2009 and 2010**

<b>Substance</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	174,119	174,194	83,336	83,200	69,713	69,559
Cigarettes	162,681	162,762	69,165	68,458	58,661	58,256
Smokeless Tobacco	44,804	44,916	12,063	12,334	8,559	8,855
Cigars	90,115	88,334	26,402	25,883	13,269	13,177
Pipe Tobacco <sup>2</sup>	35,097	34,229	--	--	2,087	2,150
<b>ALCOHOL</b>	208,545	209,295	168,141	168,396	130,621	131,342
Binge Alcohol Use <sup>3</sup>	--	--	--	--	59,561	58,550
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	17,129	16,931

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.1B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Percentages, 2009 and 2010**

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	69.1	68.7	33.1	32.8	27.7	27.4
Cigarettes	64.6	64.2	27.5	27.0	23.3	23.0
Smokeless Tobacco	17.8	17.7	4.8	4.9	3.4	3.5
Cigars	35.8	34.8	10.5	10.2	5.3	5.2
Pipe Tobacco <sup>2</sup>	13.9	13.5	--	--	0.8	0.8
<b>ALCOHOL</b>	82.8	82.5	66.8	66.4	51.9	51.8
Binge Alcohol Use <sup>3</sup>	--	--	--	--	23.7	23.1
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	6.8	6.7

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.2A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Numbers in Thousands, 2009 and 2010**

<b>Substance</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	713	678	425	400	174	187
Cigarettes	522	519	305	309	105	141
Smokeless Tobacco	177	176	97	108	40	48
Cigars	272 <sup>b</sup>	183	152	108	54	42
Pipe Tobacco <sup>2</sup>	62	46	--	--	18	17
<b>ALCOHOL</b>	1,050	979	695	679	260	249
Binge Alcohol Use <sup>3</sup>	--	--	--	--	122 <sup>a</sup>	80
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	15	9

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.2B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Percentages, 2009 and 2010**

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	9.5	8.6	5.7	5.1	2.3	2.4
Cigarettes	6.9	6.6	4.1	3.9	1.4	1.8
Smokeless Tobacco	2.4	2.2	1.3	1.4	0.5	0.6
Cigars	3.6 <sup>b</sup>	2.3	2.0 <sup>a</sup>	1.4	0.7	0.5
Pipe Tobacco <sup>2</sup>	0.8	0.6	--	--	0.2	0.2
<b>ALCOHOL</b>	14.0 <sup>a</sup>	12.4	9.3	8.6	3.5	3.1
Binge Alcohol Use <sup>3</sup>	--	--	--	--	1.6 <sup>a</sup>	1.0
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	0.2	0.1

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.3A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Numbers in Thousands, 2009 and 2010**

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	2,192 <sup>b</sup>	1,947	1,575 <sup>a</sup>	1,401	839	766
Cigarettes	1,768 <sup>b</sup>	1,575	1,221 <sup>a</sup>	1,084	641	593
Smokeless Tobacco	581	535	390	395	177	164
Cigars	978 <sup>b</sup>	814	712 <sup>b</sup>	568	266	218
Pipe Tobacco <sup>2</sup>	193	147	--	--	77 <sup>a</sup>	47
<b>ALCOHOL</b>	3,251 <sup>b</sup>	2,889	2,554 <sup>b</sup>	2,302	1,113 <sup>a</sup>	994
Binge Alcohol Use <sup>3</sup>	--	--	--	--	598	542
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	117	96

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.3B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Percentages, 2009 and 2010**

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	25.7	24.2	18.5	17.4	9.8	9.5
Cigarettes	20.7	19.6	14.3	13.5	7.5	7.4
Smokeless Tobacco	6.8	6.6	4.6	4.9	2.1	2.0
Cigars	11.4 <sup>a</sup>	10.1	8.3 <sup>a</sup>	7.1	3.1	2.7
Pipe Tobacco <sup>2</sup>	2.3	1.8	--	--	0.9	0.6
<b>ALCOHOL</b>	38.1 <sup>a</sup>	35.9	29.9	28.6	13.0	12.4
Binge Alcohol Use <sup>3</sup>	--	--	--	--	7.0	6.7
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	1.4	1.2

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.4A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Numbers in Thousands, 2009 and 2010**

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	3,720 <sup>a</sup>	3,438	2,759	2,588	1,848 <sup>a</sup>	1,644
Cigarettes	3,168 <sup>b</sup>	2,854	2,162	2,046	1,445 <sup>a</sup>	1,297
Smokeless Tobacco	1,121	1,053	778	698	359	350
Cigars	1,814	1,676	1,335 <sup>a</sup>	1,190	659 <sup>b</sup>	523
Pipe Tobacco <sup>2</sup>	385	342	--	--	136 <sup>a</sup>	90
<b>ALCOHOL</b>	5,082 <sup>b</sup>	4,706	4,215	3,967	2,250 <sup>a</sup>	2,065
Binge Alcohol Use <sup>3</sup>	--	--	--	--	1,456 <sup>a</sup>	1,288
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	385 <sup>a</sup>	303

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.4B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Percentages, 2009 and 2010**

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	43.5 <sup>a</sup>	40.9	32.2	30.8	21.6 <sup>a</sup>	19.6
Cigarettes	37.0 <sup>b</sup>	34.0	25.3	24.4	16.9	15.4
Smokeless Tobacco	13.1	12.5	9.1	8.3	4.2	4.2
Cigars	21.2	19.9	15.6 <sup>a</sup>	14.2	7.7 <sup>b</sup>	6.2
Pipe Tobacco <sup>2</sup>	4.5	4.1	--	--	1.6 <sup>a</sup>	1.1
<b>ALCOHOL</b>	59.4 <sup>b</sup>	56.0	49.3 <sup>a</sup>	47.2	26.3	24.6
Binge Alcohol Use <sup>3</sup>	--	--	--	--	17.0 <sup>a</sup>	15.3
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	4.5 <sup>a</sup>	3.6

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.5A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Numbers in Thousands, 2009 and 2010**

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	8,672	8,710	6,972	6,932	5,341	5,380
Cigarettes	7,728	7,638	5,949	5,746	4,510	4,373
Smokeless Tobacco	2,656	2,716	1,596	1,652	920	1,007
Cigars	5,109	5,021	3,594	3,435	1,774	1,754
Pipe Tobacco <sup>2</sup>	1,001 <sup>a</sup>	1,200	--	--	249 <sup>a</sup>	337
<b>ALCOHOL</b>	10,719	10,648	9,601	9,538	6,780	6,690
Binge Alcohol Use <sup>3</sup>	--	--	--	--	4,736	4,557
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	1,541	1,549

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.5B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Percentages, 2009 and 2010**

<b>Substance</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	63.6	63.6	51.1	50.6	39.1	39.3
Cigarettes	56.6	55.8	43.6	42.0	33.1	31.9
Smokeless Tobacco	19.5	19.8	11.7	12.1	6.7	7.4
Cigars	37.4	36.7	26.3	25.1	13.0	12.8
Pipe Tobacco <sup>2</sup>	7.3 <sup>b</sup>	8.8	--	--	1.8 <sup>a</sup>	2.5
<b>ALCOHOL</b>	78.6	77.8	70.4	69.6	49.7	48.9
Binge Alcohol Use <sup>3</sup>	--	--	--	--	34.7	33.3
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	11.3	11.3

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.6A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Numbers in Thousands, 2009 and 2010**

<b>Substance</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	14,687	14,665	10,597	10,491	8,639	8,525
Cigarettes	13,669	13,517	9,237	8,928	7,521	7,294
Smokeless Tobacco	4,380	4,414	1,775	1,903	1,134	1,176
Cigars	8,890	8,829	4,231	4,217	2,063	2,062
Pipe Tobacco <sup>2</sup>	1,877	2,047	--	--	335	278
<b>ALCOHOL</b>	18,091	18,559	16,855	17,278	13,988	14,273
Binge Alcohol Use <sup>3</sup>	--	--	--	--	9,262	9,274
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	3,071	3,090

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.6B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Percentages, 2009 and 2010**

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	73.7 <sup>a</sup>	72.0	53.2 <sup>a</sup>	51.5	43.3	41.8
Cigarettes	68.6 <sup>b</sup>	66.3	46.3 <sup>b</sup>	43.8	37.7 <sup>a</sup>	35.8
Smokeless Tobacco	22.0	21.7	8.9	9.3	5.7	5.8
Cigars	44.6	43.3	21.2	20.7	10.3	10.1
Pipe Tobacco <sup>2</sup>	9.4	10.0	--	--	1.7	1.4
<b>ALCOHOL</b>	90.8	91.1	84.6	84.8	70.2	70.0
Binge Alcohol Use <sup>3</sup>	--	--	--	--	46.5	45.5
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	15.4	15.2

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.7A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Numbers in Thousands, 2009 and 2010**

<b>Substance</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	27,571	27,845	16,785	16,624	14,331	14,066
Cigarettes	25,823	26,258	14,308	14,067	12,315	12,267
Smokeless Tobacco	8,366	8,385	2,518	2,412	1,788	1,764
Cigars	16,403	16,385	5,564	5,235	2,677	2,482
Pipe Tobacco <sup>2</sup>	3,168	3,228	--	--	360	288
<b>ALCOHOL</b>	32,760	33,103	28,955	28,974	23,282	23,581
Binge Alcohol Use <sup>3</sup>	--	--	--	--	13,156	13,313
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	3,658	3,750

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.7B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Percentages, 2009 and 2010**

<b>Substance</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	76.1	76.2	46.3	45.5	39.6	38.5
Cigarettes	71.3	71.9	39.5	38.5	34.0	33.6
Smokeless Tobacco	23.1	23.0	7.0	6.6	4.9	4.8
Cigars	45.3	44.9	15.4	14.3	7.4	6.8
Pipe Tobacco <sup>2</sup>	8.7	8.8	--	--	1.0	0.8
<b>ALCOHOL</b>	90.5	90.6	80.0	79.3	64.3	64.6
Binge Alcohol Use <sup>3</sup>	--	--	--	--	36.3	36.5
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	10.1	10.3

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.8A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Numbers in Thousands, 2009 and 2010**

<b>Substance</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	116,564	116,909	44,221	44,764	38,542	38,991
Cigarettes	110,003	110,401	35,981	36,278	32,125	32,292
Smokeless Tobacco	27,523	27,637	4,907	5,166	4,141	4,346
Cigars	56,649	55,426	10,814	11,130	5,776	6,097
Pipe Tobacco <sup>2</sup>	28,410	27,218	--	--	912	1,092
<b>ALCOHOL</b>	137,592	138,412	105,265	105,657	82,949	83,489
Binge Alcohol Use <sup>3</sup>	--	--	--	--	30,232	29,496
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	8,342	8,134

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.8B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Percentages, 2009 and 2010**

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	74.1	73.7	28.1	28.2	24.5	24.6
Cigarettes	69.9	69.6	22.9	22.9	20.4	20.4
Smokeless Tobacco	17.5	17.4	3.1	3.3	2.6	2.7
Cigars	36.0	34.9	6.9	7.0	3.7	3.8
Pipe Tobacco <sup>2</sup>	18.0	17.2	--	--	0.6	0.7
<b>ALCOHOL</b>	87.4	87.2	66.9	66.6	52.7	52.6
Binge Alcohol Use <sup>3</sup>	--	--	--	--	19.2	18.6
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	5.3	5.1

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.9A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Numbers in Thousands, 2009 and 2010**

<b>Substance</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOBACCO PRODUCTS<sup>1</sup></b>	93,468	94,076	49,045	49,791	40,909	41,587
Cigarettes	84,701	85,394	36,763	36,968	30,937	31,361
Smokeless Tobacco	37,948	38,501	11,112	11,330	8,151	8,346
Cigars	65,916	64,963	20,515	20,048	10,679	10,507
Pipe Tobacco <sup>2</sup>	30,432	30,105	--	--	1,772	1,735
<b>ALCOHOL</b>	104,908	105,512	86,393	85,698	70,455	70,834
Binge Alcohol Use <sup>3</sup>	--	--	--	--	38,654	38,171
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	12,604	12,490

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.9B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males Aged 12 or Older: Percentages, 2009 and 2010**

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	76.4	76.2	40.1	40.3	33.5	33.7
Cigarettes	69.3	69.2	30.1	30.0	25.3	25.4
Smokeless Tobacco	31.0	31.2	9.1	9.2	6.7	6.8
Cigars	53.9	52.6	16.8	16.2	8.7	8.5
Pipe Tobacco <sup>2</sup>	24.9	24.4	--	--	1.4	1.4
<b>ALCOHOL</b>	85.8	85.5	70.6	69.4	57.6	57.4
Binge Alcohol Use <sup>3</sup>	--	--	--	--	31.6	30.9
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	10.3	10.1

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.10A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Numbers in Thousands, 2009 and 2010**

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	80,651	80,118	34,291	33,409	28,804	27,972
Cigarettes	77,980	77,368	32,402	31,490	27,724	26,895
Smokeless Tobacco	6,856	6,415	950	1,004	408	509
Cigars	24,199	23,371	5,887	5,835	2,590	2,670
Pipe Tobacco <sup>2</sup>	4,665	4,124	--	--	315	415
<b>ALCOHOL</b>	103,637	103,783	81,748	82,697	60,166	60,508
Binge Alcohol Use <sup>3</sup>	--	--	--	--	20,908	20,378
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	4,525	4,441

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.10B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females Aged 12 or Older: Percentages, 2009 and 2010**

Substance	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
<b>TOBACCO PRODUCTS<sup>1</sup></b>	62.3	61.5	26.5	25.7	22.2	21.5
Cigarettes	60.2	59.4	25.0	24.2	21.4	20.7
Smokeless Tobacco	5.3	4.9	0.7	0.8	0.3	0.4
Cigars	18.7	18.0	4.5	4.5	2.0	2.1
Pipe Tobacco <sup>2</sup>	3.6	3.2	--	--	0.2	0.3
<b>ALCOHOL</b>	80.0	79.7	63.1	63.5	46.5	46.5
Binge Alcohol Use <sup>3</sup>	--	--	--	--	16.1	15.7
Heavy Alcohol Use <sup>3</sup>	--	--	--	--	3.5	3.4

\*Low precision; no estimate reported.

-- Not available.

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

<sup>2</sup> Information about past year use of pipe tobacco was not collected.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.