

**Table 2.57A (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	44,962	45,036	12,142	12,383	8,615	8,879
<b>GEOGRAPHIC DIVISION</b>						
Northeast	6,362	6,574	1,528	1,539	1,028	1,013
New England	1,675	1,864	375	386	240	224
Middle Atlantic	4,687	4,710	1,153	1,153	788	788
Midwest	11,460	11,092	3,211	3,090	2,206	2,136
East North Central	7,205	7,038	1,923	1,946	1,331	1,345
West North Central	4,255	4,054	1,288	1,144	875	791
South	17,960	17,632	5,073	5,252	3,753	3,953
South Atlantic	8,588	8,208	2,208	2,258	1,513	1,667
East South Central	3,534	3,822	1,046	1,203	793	970
West South Central	5,839	5,602	1,818	1,791	1,447	1,316
West	9,180	9,738	2,330	2,502	1,628	1,777
Mountain	3,607	3,730	981	1,219	742	843
Pacific	5,574	6,007	1,349	1,283	886	934
<b>COUNTY TYPE</b>						
Large Metro	18,944	19,605	4,201	4,523	2,841	3,075
Small Metro	15,457	15,326	4,367	4,464	3,089	3,145
250K – 1 Mil. Pop.	9,467	9,543	2,623	2,832	1,833	1,980
< 250K Pop.	5,990	5,783	1,744	1,633	1,256	1,165
Nonmetro	10,561	10,105	3,575	3,396	2,685	2,658
Urbanized	4,165	4,093	1,274	1,301	841	980
Less Urbanized	5,167	4,949	1,874	1,591	1,542	1,260
Completely Rural	1,229	1,064	427	503	302	418

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.57B (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	17.9	17.8	4.8	4.9	3.4	3.5
<b>GEOGRAPHIC DIVISION</b>						
Northeast	13.7	14.1	3.3	3.3	2.2	2.2
New England	13.8	15.2	3.1	3.2	2.0	1.8
Middle Atlantic	13.7	13.7	3.4	3.4	2.3	2.3
Midwest	20.8	20.0	5.8	5.6	4.0	3.9
East North Central	18.7	18.3	5.0	5.1	3.5	3.5
West North Central	25.5	24.1	7.7	6.8	5.2	4.7
South	19.5	19.0	5.5	5.6	4.1	4.3
South Atlantic	17.7	16.8	4.5	4.6	3.1	3.4
East South Central	23.6	25.3	7.0	8.0	5.3	6.4
West South Central	20.5	19.4	6.4	6.2	5.1	4.6
West	15.8	16.6	4.0	4.3	2.8	3.0
Mountain	20.2	20.7	5.5	6.8	4.2	4.7
Pacific	13.8	14.7	3.3	3.2	2.2	2.3
<b>COUNTY TYPE</b>						
Large Metro	14.2	14.4	3.1	3.3	2.1	2.3
Small Metro	20.1	19.8	5.7	5.8	4.0	4.1
250K – 1 Mil. Pop.	18.7	18.5	5.2	5.5	3.6	3.8
< 250K Pop.	22.9	22.3	6.7	6.3	4.8	4.5
Nonmetro	25.6	25.0	8.7	8.4	6.5	6.6
Urbanized	23.1	24.0	7.1	7.6	4.7	5.7
Less Urbanized	27.2	25.2	9.9 <sup>a</sup>	8.1	8.1 <sup>a</sup>	6.4
Completely Rural	29.4	28.6	10.2	13.5	7.2	11.2

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.58A (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	1,889	1,783	1,272	1,215	583	570
<b>GEOGRAPHIC DIVISION</b>						
Northeast	243	263	196	206	78	91
New England	58	76	46	64	15	25
Middle Atlantic	185	187	150	142	63	65
Midwest	448	421	318	296	148	128
East North Central	281	259	197	176	86	74
West North Central	168	162	122	120	61	54
South	852	789	523	495	257	262
South Atlantic	380	360	243	225	121	117
East South Central	218	214	139	133	70	72
West South Central	254	214	141	137	66	73
West	346	309	234	218	101	89
Mountain	139	142	86	103	41	41
Pacific	207	167	148	115	59	48
<b>COUNTY TYPE</b>						
Large Metro	697	641	446	462	179	192
Small Metro	630	617	448	415	209	203
250K – 1 Mil. Pop.	392	373	284	255	132	131
< 250K Pop.	238	244	164	160	77	71
Nonmetro	562	525	378	338	196	176
Urbanized	229	177	165 <sup>a</sup>	118	75	52
Less Urbanized	268	295	167	178	89	98
Completely Rural	65	52	47	41	31	26

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.58B (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	7.7	7.3	5.2	5.0	2.4	2.3
<b>GEOGRAPHIC DIVISION</b>						
Northeast	5.6	6.2	4.6	4.9	1.8	2.1
New England	5.2	6.9	4.2	5.8	1.4	2.3
Middle Atlantic	5.8	6.0	4.7	4.5	2.0	2.1
Midwest	8.3	7.9	5.9	5.6	2.7	2.4
East North Central	7.4	6.9	5.2	4.7	2.3	2.0
West North Central	10.3	10.2	7.5	7.5	3.8	3.4
South	9.5	8.8	5.8	5.5	2.9	2.9
South Atlantic	8.4	8.0	5.3	5.0	2.7	2.6
East South Central	15.0	14.8	9.6	9.2	4.8	5.0
West South Central	8.4	7.1	4.7	4.5	2.2	2.4
West	5.9	5.3	4.0	3.7	1.7	1.5
Mountain	7.7	7.9	4.8	5.7	2.3	2.3
Pacific	5.1	4.1	3.6	2.9	1.4	1.2
<b>COUNTY TYPE</b>						
Large Metro	5.3	4.9	3.4	3.5	1.4	1.5
Small Metro	8.2	8.2	5.8	5.5	2.7	2.7
250K – 1 Mil. Pop.	7.7	7.4	5.5	5.1	2.6	2.6
< 250K Pop.	9.3	9.8	6.4	6.5	3.0	2.9
Nonmetro	14.4	13.9	9.7	8.9	5.0	4.6
Urbanized	13.3	11.6	9.6	7.7	4.4	3.4
Less Urbanized	14.9	15.2	9.3	9.2	5.0	5.1
Completely Rural	16.9	16.5	12.0	12.9	8.1	8.1

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.59A (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	7,081	7,160	3,386	3,569	2,056	2,182
<b>GEOGRAPHIC DIVISION</b>						
Northeast	1,154	1,106	496	539	294	327
New England	298	312	126	160	67 <sup>a</sup>	102
Middle Atlantic	856	794	370	380	227	226
Midwest	1,877	1,962	960	993	589	599
East North Central	1,215	1,250	619	638	369	375
West North Central	662	713	340	355	221	224
South	2,562	2,601	1,256	1,316	796	830
South Atlantic	1,213	1,209	588	593	378	369
East South Central	484	523	274	255	180	191
West South Central	865	868	394	468	237	270
West	1,488	1,492	675	720	377	426
Mountain	595	548	270	297	170	166
Pacific	893	943	405	423	207	260
<b>COUNTY TYPE</b>						
Large Metro	2,803	3,033	1,223	1,395	694	760
Small Metro	2,604	2,596	1,271	1,318	784	819
250K – 1 Mil. Pop.	1,518	1,505	721	773	425	472
< 250K Pop.	1,085	1,091	550	545	359	347
Nonmetro	1,674	1,531	892	856	578	603
Urbanized	773	661	395	326	258	199
Less Urbanized	774	744	423	456	271	353
Completely Rural	126	126	74	73	49	51

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.59B (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	21.1	21.0	10.1	10.5	6.1	6.4
<b>GEOGRAPHIC DIVISION</b>						
Northeast	18.9	18.3	8.1	8.9	4.8	5.4
New England	18.7	19.2	7.9	9.8	4.2 <sup>a</sup>	6.3
Middle Atlantic	18.9	18.0	8.2	8.6	5.0	5.1
Midwest	25.6	26.2	13.1	13.3	8.0	8.0
East North Central	23.9	24.3	12.2	12.4	7.3	7.3
West North Central	29.3	30.2	15.0	15.1	9.8	9.5
South	21.3	20.9	10.4	10.6	6.6	6.7
South Atlantic	19.8	19.1	9.6	9.4	6.2	5.8
East South Central	25.5	26.3	14.4	12.9	9.5	9.6
West South Central	21.6	21.1	9.8	11.4	5.9	6.6
West	18.4	18.4	8.3	8.9	4.7	5.2
Mountain	24.6	22.0	11.2	11.9	7.0	6.6
Pacific	15.7	16.8	7.1	7.5	3.7	4.6
<b>COUNTY TYPE</b>						
Large Metro	15.7	16.7	6.9	7.7	3.9	4.2
Small Metro	24.8	23.3	12.1	11.8	7.5	7.4
250K – 1 Mil. Pop.	22.7	21.4	10.8	11.0	6.4	6.7
< 250K Pop.	28.5	26.6	14.5	13.3	9.4	8.5
Nonmetro	31.9	31.9	17.0	17.9	11.0	12.6
Urbanized	30.8	31.3	15.7	15.5	10.3	9.4
Less Urbanized	32.2	31.9	17.6	19.6	11.3 <sup>a</sup>	15.1
Completely Rural	38.9	36.1	22.7	21.0	15.1	14.6

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.60A (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	35,992	36,092	7,484	7,600	5,976	6,127
<b>GEOGRAPHIC DIVISION</b>						
Northeast	4,965	5,205	836	794	656	595
New England	1,319	1,476	203	163	158	97
Middle Atlantic	3,646	3,729	633	631	498	498
Midwest	9,134	8,709	1,933	1,801	1,469	1,409
East North Central	5,709	5,529	1,107	1,132	876	896
West North Central	3,425	3,179	826	669	594	513
South	14,547	14,242	3,294	3,441	2,700	2,861
South Atlantic	6,995	6,638	1,378	1,441	1,014	1,181
East South Central	2,831	3,085	633	814	543	707
West South Central	4,720	4,519	1,283	1,186	1,144	973
West	7,346	7,937	1,421	1,564	1,150	1,262
Mountain	2,873	3,040	625	819	531	637
Pacific	4,473	4,897	796	746	619	625
<b>COUNTY TYPE</b>						
Large Metro	15,445	15,930	2,532	2,666	1,969	2,124
Small Metro	12,223	12,113	2,648	2,732	2,096	2,124
250K – 1 Mil. Pop.	7,557	7,665	1,617	1,804	1,276	1,377
< 250K Pop.	4,666	4,448	1,030	927	820	746
Nonmetro	8,324	8,049	2,304	2,202	1,911	1,880
Urbanized	3,163	3,255	714	857	508	729
Less Urbanized	4,124	3,909	1,283	957	1,181 <sup>a</sup>	809
Completely Rural	1,037	885	307	389	221	342

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.60B (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	18.6	18.5	3.9	3.9	3.1	3.1
<b>GEOGRAPHIC DIVISION</b>						
Northeast	13.8	14.4	2.3	2.2	1.8	1.6
New England	13.9	15.5	2.1	1.7	1.7	1.0
Middle Atlantic	13.8	14.0	2.4	2.4	1.9	1.9
Midwest	21.5	20.5	4.6	4.2	3.5	3.3
East North Central	19.3	18.6	3.7	3.8	3.0	3.0
West North Central	26.7	24.7	6.4	5.2	4.6	4.0
South	20.5	19.9	4.6	4.8	3.8	4.0
South Atlantic	18.4	17.4	3.6	3.8	2.7	3.1
East South Central	24.3	26.5	5.4	7.0	4.7	6.1
West South Central	22.0	20.8	6.0	5.5	5.3	4.5
West	16.6	17.7	3.2	3.5	2.6	2.8
Mountain	21.1	22.1	4.6	6.0	3.9	4.6
Pacific	14.6	15.7	2.6	2.4	2.0	2.0
<b>COUNTY TYPE</b>						
Large Metro	15.0	15.2	2.5	2.6	1.9	2.0
Small Metro	20.9	20.6	4.5	4.6	3.6	3.6
250K – 1 Mil. Pop.	19.5	19.4	4.2	4.6	3.3	3.5
< 250K Pop.	23.6	22.9	5.2	4.8	4.2	3.9
Nonmetro	25.9	25.3	7.2	6.9	6.0	5.9
Urbanized	22.9	24.2	5.2	6.4	3.7 <sup>a</sup>	5.4
Less Urbanized	27.8	25.4	8.7 <sup>a</sup>	6.2	8.0 <sup>b</sup>	5.3
Completely Rural	29.9	28.9	8.8	12.7	6.4	11.2

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.61A (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	43,073	43,253	10,870	11,169	8,032	8,309
<b>GEOGRAPHIC DIVISION</b>						
Northeast	6,119	6,311	1,332	1,333	950	922
New England	1,617	1,788	329	322	225	199
Middle Atlantic	4,502	4,523	1,003	1,011	725	723
Midwest	11,011	10,671	2,893	2,794	2,059	2,008
East North Central	6,924	6,779	1,726	1,770	1,244	1,271
West North Central	4,087	3,892	1,167	1,024	814	737
South	17,108	16,842	4,550	4,757	3,496	3,691
South Atlantic	8,208	7,847	1,966	2,034	1,392	1,550
East South Central	3,315	3,608	907	1,070	723	898
West South Central	5,585	5,387	1,677	1,654	1,381	1,243
West	8,834	9,429	2,096	2,284	1,527	1,688
Mountain	3,468	3,588	895	1,116	700	802
Pacific	5,366	5,840	1,201	1,168	827	885
<b>COUNTY TYPE</b>						
Large Metro	18,248	18,964	3,755	4,061	2,663	2,884
Small Metro	14,827	14,709	3,919	4,049	2,880	2,943
250K – 1 Mil. Pop.	9,075	9,170	2,339	2,577	1,701	1,849
< 250K Pop.	5,751	5,538	1,580	1,472	1,179	1,093
Nonmetro	9,998	9,580	3,196	3,058	2,489	2,483
Urbanized	3,936	3,916	1,109	1,183	766	928
Less Urbanized	4,899	4,653	1,706	1,413	1,452	1,162
Completely Rural	1,163	1,011	381	462	270	393

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.61B (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	19.0	18.9	4.8	4.9	3.5	3.6
<b>GEOGRAPHIC DIVISION</b>						
Northeast	14.5	14.9	3.2	3.2	2.3	2.2
New England	14.6	16.0	3.0	2.9	2.0	1.8
Middle Atlantic	14.5	14.5	3.2	3.2	2.3	2.3
Midwest	22.1	21.3	5.8	5.6	4.1	4.0
East North Central	20.0	19.5	5.0	5.1	3.6	3.7
West North Central	27.1	25.6	7.7	6.7	5.4	4.8
South	20.6	20.0	5.5	5.7	4.2	4.4
South Atlantic	18.6	17.6	4.5	4.6	3.2	3.5
East South Central	24.5	26.5	6.7	7.8	5.3	6.6
West South Central	21.9	20.8	6.6	6.4	5.4	4.8
West	16.9	17.8	4.0	4.3	2.9	3.2
Mountain	21.7	22.1	5.6	6.9	4.4	4.9
Pacific	14.8	15.9	3.3	3.2	2.3	2.4
<b>COUNTY TYPE</b>						
Large Metro	15.1	15.5	3.1	3.3	2.2	2.4
Small Metro	21.5	21.0	5.7	5.8	4.2	4.2
250K – 1 Mil. Pop.	19.9	19.7	5.1	5.5	3.7	4.0
< 250K Pop.	24.4	23.6	6.7	6.3	5.0	4.7
Nonmetro	26.8	26.1	8.6	8.3	6.7	6.8
Urbanized	24.1	25.2	6.8	7.6	4.7	6.0
Less Urbanized	28.4	26.3	9.9 <sup>a</sup>	8.0	8.4 <sup>a</sup>	6.6
Completely Rural	30.6	29.7	10.0	13.6	7.1	11.5

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.