

**Table 2.62A (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:  
Numbers in Thousands, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	90,460	88,760	26,592	26,061	13,345	13,273
<b>GEOGRAPHIC DIVISION</b>						
Northeast	16,338	16,127	5,141	4,925	2,460	2,524
New England	4,813	4,749	1,352	1,432	683	696
Middle Atlantic	11,525	11,378	3,789	3,492	1,777	1,828
Midwest	21,563	21,519	6,558	6,308	2,966	3,292
East North Central	14,629	14,632	4,441	4,487	2,081	2,345
West North Central	6,933	6,887	2,118	1,821	885	947
South	30,912	30,898	8,742	9,268	4,884	5,060
South Atlantic	16,115	16,398	4,429	4,769	2,459	2,558
East South Central	5,282	5,198	1,305 <sup>a</sup>	1,612	811	950
West South Central	9,514	9,302	3,009	2,887	1,615	1,552
West	21,647 <sup>a</sup>	20,216	6,151	5,560	3,035 <sup>b</sup>	2,398
Mountain	7,143	6,578	1,985	1,823	908	771
Pacific	14,504	13,638	4,165	3,737	2,127 <sup>a</sup>	1,627
<b>COUNTY TYPE</b>						
Large Metro	47,146	46,129	14,309	13,835	7,270	6,940
Small Metro	28,139	28,533	8,332	8,427	4,119	4,319
250K – 1 Mil. Pop.	18,179	18,770	5,471	5,471	2,584	2,818
< 250K Pop.	9,960	9,764	2,860	2,956	1,535	1,501
Nonmetro	15,175	14,097	3,951	3,798	1,956	2,014
Urbanized	6,675	6,276	1,782	1,689	895	793
Less Urbanized	6,760	6,639	1,853	1,877	924	1,071
Completely Rural	1,740 <sup>a</sup>	1,182	316	233	137	150

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.62B (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	35.9	35.0	10.6	10.3	5.3	5.2
<b>GEOGRAPHIC DIVISION</b>						
Northeast	35.2	34.7	11.1	10.6	5.3	5.4
New England	39.5	38.8	11.1	11.7	5.6	5.7
Middle Atlantic	33.7	33.2	11.1	10.2	5.2	5.3
Midwest	39.1	38.9	11.9	11.4	5.4	5.9
East North Central	38.0	38.0	11.5	11.6	5.4	6.1
West North Central	41.5	41.0	12.7	10.8	5.3	5.6
South	33.6	33.2	9.5	10.0	5.3	5.4
South Atlantic	33.2	33.5	9.1	9.7	5.1	5.2
East South Central	35.2	34.5	8.7	10.7	5.4	6.3
West South Central	33.4	32.2	10.6	10.0	5.7	5.4
West	37.2 <sup>a</sup>	34.4	10.6	9.5	5.2 <sup>b</sup>	4.1
Mountain	40.1	36.5	11.1	10.1	5.1	4.3
Pacific	35.9	33.5	10.3	9.2	5.3 <sup>a</sup>	4.0
<b>COUNTY TYPE</b>						
Large Metro	35.2	34.0	10.7	10.2	5.4	5.1
Small Metro	36.7	36.8	10.9	10.9	5.4	5.6
250K – 1 Mil. Pop.	35.9	36.5	10.8	10.6	5.1	5.5
< 250K Pop.	38.1	37.6	11.0	11.4	5.9	5.8
Nonmetro	36.8	34.9	9.6	9.4	4.7	5.0
Urbanized	37.0	36.7	9.9	9.9	5.0	4.6
Less Urbanized	35.5	33.8	9.7	9.6	4.9	5.4
Completely Rural	41.6 <sup>b</sup>	31.7	7.6	6.2	3.3	4.0

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.63A (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	3,088 <sup>b</sup>	2,693	2,218 <sup>b</sup>	1,882	990 <sup>b</sup>	788
<b>GEOGRAPHIC DIVISION</b>						
Northeast	532	481	397	356	160	138
New England	163	160	130	114	55	42
Middle Atlantic	369	321	267	242	105	96
Midwest	746 <sup>b</sup>	619	546 <sup>b</sup>	431	272 <sup>b</sup>	196
East North Central	527 <sup>b</sup>	415	384 <sup>b</sup>	301	189 <sup>b</sup>	135
West North Central	219	204	162	130	83	61
South	1,107 <sup>a</sup>	981	750	680	348	286
South Atlantic	554	475	365	326	175	153
East South Central	197	175	126	121	60	58
West South Central	356	331	260	233	113 <sup>a</sup>	75
West	703	612	525 <sup>a</sup>	415	210	168
Mountain	229	201	161	136	72	63
Pacific	474	410	364	279	137	105
<b>COUNTY TYPE</b>						
Large Metro	1,574 <sup>b</sup>	1,318	1,122 <sup>a</sup>	951	489 <sup>b</sup>	384
Small Metro	964	950	714	631	314	274
250K – 1 Mil. Pop.	638	658	473	452	205	195
< 250K Pop.	326	292	241 <sup>a</sup>	179	109	78
Nonmetro	551 <sup>b</sup>	426	381 <sup>a</sup>	300	186 <sup>a</sup>	131
Urbanized	238 <sup>b</sup>	161	171 <sup>a</sup>	114	88 <sup>b</sup>	40
Less Urbanized	263	232	185	160	86	75
Completely Rural	50	32	25	26	13	16

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.63B (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	12.5 <sup>b</sup>	11.1	9.0 <sup>b</sup>	7.7	4.0 <sup>b</sup>	3.2
<b>GEOGRAPHIC DIVISION</b>						
Northeast	12.3	11.4	9.2	8.4	3.7	3.3
New England	14.6	14.6	11.7	10.4	5.0	3.8
Middle Atlantic	11.6	10.2	8.3	7.7	3.3	3.1
Midwest	13.8 <sup>b</sup>	11.6	10.1 <sup>b</sup>	8.1	5.0 <sup>b</sup>	3.7
East North Central	13.9 <sup>b</sup>	11.1	10.1 <sup>b</sup>	8.0	5.0 <sup>b</sup>	3.6
West North Central	13.5	12.8	10.0	8.1	5.1	3.8
South	12.3 <sup>a</sup>	11.0	8.3	7.6	3.9	3.2
South Atlantic	12.2	10.6	8.0	7.2	3.9	3.4
East South Central	13.5	12.1	8.6	8.4	4.1	4.0
West South Central	11.8	11.0	8.6	7.7	3.7 <sup>a</sup>	2.5
West	12.0	10.5	8.9 <sup>a</sup>	7.1	3.6	2.9
Mountain	12.7	11.3	8.9	7.6	4.0	3.5
Pacific	11.6	10.2	8.9	6.9	3.4	2.6
<b>COUNTY TYPE</b>						
Large Metro	12.1 <sup>b</sup>	10.1	8.6 <sup>b</sup>	7.3	3.8 <sup>b</sup>	2.9
Small Metro	12.6	12.6	9.3	8.4	4.1	3.6
250K – 1 Mil. Pop.	12.5	13.1	9.2	9.0	4.0	3.9
< 250K Pop.	12.7	11.7	9.4 <sup>a</sup>	7.2	4.3	3.2
Nonmetro	14.1 <sup>b</sup>	11.2	9.8 <sup>a</sup>	7.9	4.8 <sup>a</sup>	3.5
Urbanized	13.9 <sup>a</sup>	10.6	10.0 <sup>a</sup>	7.5	5.1 <sup>b</sup>	2.6
Less Urbanized	14.6	12.0	10.3	8.3	4.8	3.9
Completely Rural	12.9	10.2	6.6	8.1	3.3	5.0

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.64A (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	14,132	13,988	7,904	7,735	3,871	3,858
<b>GEOGRAPHIC DIVISION</b>						
Northeast	2,574	2,551	1,416	1,414	660	680
New England	742	758	405	407	198	190
Middle Atlantic	1,831	1,793	1,011	1,006	462	490
Midwest	3,419	3,483	1,917	1,933	968	1,032
East North Central	2,353	2,339	1,329	1,314	678	714
West North Central	1,066	1,144	588	619	290	318
South	4,764	4,722	2,743	2,669	1,407	1,430
South Atlantic	2,458	2,351	1,388	1,330	696	696
East South Central	727	790	439	480	259	274
West South Central	1,578	1,581	915	859	452	460
West	3,375	3,232	1,829	1,719	836	716
Mountain	1,064	1,103	564	603	264	270
Pacific	2,311	2,128	1,265	1,116	573 <sup>a</sup>	446
<b>COUNTY TYPE</b>						
Large Metro	7,169	6,945	3,988	3,768	1,975	1,858
Small Metro	4,676	4,885	2,589	2,747	1,234	1,353
250K – 1 Mil. Pop.	2,867	3,054	1,558	1,713	739	842
< 250K Pop.	1,808	1,831	1,031	1,033	496	511
Nonmetro	2,287	2,158	1,327	1,220	662	647
Urbanized	1,170	1,037	670	578	329	298
Less Urbanized	977	975	578	573	294	309
Completely Rural	140	146	79	69	38	40

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.64B (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	42.1	41.1	23.5	22.7	11.5	11.3
<b>GEOGRAPHIC DIVISION</b>						
Northeast	42.0	42.2	23.1	23.4	10.8	11.2
New England	46.5	46.7	25.4	25.1	12.4	11.7
Middle Atlantic	40.5	40.6	22.3	22.8	10.2	11.1
Midwest	46.6	46.5	26.1	25.8	13.2	13.8
East North Central	46.4	45.5	26.2	25.6	13.4	13.9
West North Central	47.2	48.6	26.0	26.3	12.8	13.5
South	39.6	38.0	22.8	21.5	11.7	11.5
South Atlantic	40.2	37.2	22.7	21.1	11.4	11.0
East South Central	38.2	39.8	23.1	24.1	13.6	13.8
West South Central	39.4	38.4	22.8	20.9	11.3	11.2
West	41.7	39.8	22.6	21.2	10.3	8.8
Mountain	44.0	44.2	23.3	24.1	10.9	10.8
Pacific	40.7	37.9	22.3	19.9	10.1	7.9
<b>COUNTY TYPE</b>						
Large Metro	40.2 <sup>a</sup>	38.2	22.3 <sup>a</sup>	20.8	11.1	10.2
Small Metro	44.6	43.9	24.7	24.7	11.8	12.2
250K – 1 Mil. Pop.	42.9	43.5	23.3	24.4	11.0	12.0
< 250K Pop.	47.5	44.7	27.1	25.2	13.0	12.5
Nonmetro	43.6	45.0	25.3	25.5	12.6	13.5
Urbanized	46.5	49.1	26.6	27.4	13.1	14.1
Less Urbanized	40.7	41.8	24.1	24.6	12.3	13.2
Completely Rural	43.2	41.9	24.4	19.7	11.6	11.5

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.65A (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:  
Numbers in Thousands, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	73,240	72,079	16,470	16,444	8,484	8,627
<b>GEOGRAPHIC DIVISION</b>						
Northeast	13,233	13,094	3,328	3,155	1,639	1,706
New England	3,908	3,831	816	911	430	464
Middle Atlantic	9,325	9,264	2,512	2,244	1,210	1,242
Midwest	17,397	17,417	4,095	3,944	1,726	2,064
East North Central	11,749	11,878	2,728	2,872	1,213	1,495
West North Central	5,649	5,538	1,367	1,072	512	569
South	25,041	25,195	5,250	5,919	3,130	3,344
South Atlantic	13,103	13,572	2,675	3,113	1,587	1,709
East South Central	4,358	4,233	740 <sup>a</sup>	1,011	492	617
West South Central	7,581	7,390	1,834	1,795	1,050	1,017
West	17,569	16,373	3,797	3,426	1,989 <sup>a</sup>	1,514
Mountain	5,849	5,274	1,261	1,084	572	438
Pacific	11,720	11,099	2,537	2,342	1,417	1,076
<b>COUNTY TYPE</b>						
Large Metro	38,403	37,866	9,199	9,115	4,805	4,699
Small Metro	22,500	22,699	5,028	5,050	2,571	2,692
250K – 1 Mil. Pop.	14,673	15,058	3,440	3,306	1,641	1,781
< 250K Pop.	7,826	7,641	1,588	1,744	930	911
Nonmetro	12,337	11,513	2,243	2,278	1,108	1,237
Urbanized	5,267	5,077	941	997	478	456
Less Urbanized	5,521	5,432	1,091	1,143	544	687
Completely Rural	1,550 <sup>b</sup>	1,004	211	138	87	94

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.65B (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	37.8	36.9	8.5	8.4	4.4	4.4
<b>GEOGRAPHIC DIVISION</b>						
Northeast	36.8	36.1	9.3	8.7	4.6	4.7
New England	41.3	40.2	8.6	9.6	4.5	4.9
Middle Atlantic	35.2	34.7	9.5	8.4	4.6	4.6
Midwest	41.0	41.0	9.7	9.3	4.1	4.9
East North Central	39.7	40.1	9.2	9.7	4.1	5.0
West North Central	44.1	43.1	10.7	8.3	4.0	4.4
South	35.3	35.2	7.4	8.3	4.4	4.7
South Atlantic	34.6	35.5	7.1	8.2	4.2	4.5
East South Central	37.5	36.3	6.4 <sup>a</sup>	8.7	4.2	5.3
West South Central	35.3	34.0	8.5	8.3	4.9	4.7
West	39.7 <sup>a</sup>	36.5	8.6	7.6	4.5 <sup>a</sup>	3.4
Mountain	43.0	38.4	9.3	7.9	4.2	3.2
Pacific	38.3	35.7	8.3	7.5	4.6	3.5
<b>COUNTY TYPE</b>						
Large Metro	37.3	36.2	8.9	8.7	4.7	4.5
Small Metro	38.4	38.6	8.6	8.6	4.4	4.6
250K – 1 Mil. Pop.	37.8	38.2	8.9	8.4	4.2	4.5
< 250K Pop.	39.6	39.4	8.0	9.0	4.7	4.7
Nonmetro	38.4	36.1	7.0	7.1	3.5	3.9
Urbanized	38.1	37.8	6.8	7.4	3.5	3.4
Less Urbanized	37.2	35.3	7.4	7.4	3.7	4.5
Completely Rural	44.6 <sup>b</sup>	32.8	6.1	4.5	2.5	3.1

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.66A (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:  
Numbers in Thousands, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	87,372	86,067	24,374	24,179	12,355	12,485
<b>GEOGRAPHIC DIVISION</b>						
Northeast	15,806	15,645	4,744	4,568	2,299	2,386
New England	4,650	4,589	1,222	1,318	628	654
Middle Atlantic	11,156	11,057	3,522	3,250	1,671	1,731
Midwest	20,817	20,900	6,012	5,877	2,694	3,096
East North Central	14,102	14,217	4,057	4,186	1,891	2,209
West North Central	6,715	6,683	1,955	1,692	803	887
South	29,805	29,917	7,992	8,588	4,537	4,774
South Atlantic	15,561	15,923	4,064	4,443	2,283	2,405
East South Central	5,085	5,023	1,179 <sup>a</sup>	1,491	751	892
West South Central	9,159	8,971	2,749	2,654	1,502	1,477
West	20,944 <sup>a</sup>	19,604	5,626	5,145	2,825 <sup>a</sup>	2,230
Mountain	6,914	6,377	1,825	1,687	835	708
Pacific	14,030	13,228	3,801	3,458	1,990 <sup>a</sup>	1,522
<b>COUNTY TYPE</b>						
Large Metro	45,572	44,811	13,187	12,884	6,781	6,557
Small Metro	27,175	27,584	7,617	7,797	3,805	4,045
250K – 1 Mil. Pop.	17,540	18,112	4,998	5,019	2,379	2,623
< 250K Pop.	9,635	9,472	2,619	2,777	1,425	1,422
Nonmetro	14,625	13,672	3,570	3,499	1,770	1,883
Urbanized	6,437	6,114	1,610	1,575	807	754
Less Urbanized	6,497	6,407	1,669	1,717	838	996
Completely Rural	1,690 <sup>a</sup>	1,150	291	207	124	134

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.66B (REVISED) Cigar Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2009 and 2010**

<b>Geographic Characteristic</b>	<b>Lifetime (2009)</b>	<b>Lifetime (2010)</b>	<b>Past Year (2009)</b>	<b>Past Year (2010)</b>	<b>Past Month (2009)</b>	<b>Past Month (2010)</b>
<b>TOTAL</b>	38.5	37.5	10.7	10.5	5.4	5.4
<b>GEOGRAPHIC DIVISION</b>						
Northeast	37.6	37.0	11.3	10.8	5.5	5.6
New England	42.0	41.1	11.0	11.8	5.7	5.9
Middle Atlantic	36.0	35.5	11.4	10.4	5.4	5.6
Midwest	41.8	41.8	12.1	11.8	5.4	6.2
East North Central	40.7	40.9	11.7	12.0	5.5	6.4
West North Central	44.5	43.9	13.0	11.1	5.3	5.8
South	35.9	35.6	9.6	10.2	5.5	5.7
South Atlantic	35.3	35.8	9.2	10.0	5.2	5.4
East South Central	37.6	36.8	8.7 <sup>a</sup>	10.9	5.6	6.5
West South Central	35.9	34.7	10.8	10.3	5.9	5.7
West	40.0 <sup>a</sup>	37.0	10.8	9.7	5.4 <sup>b</sup>	4.2
Mountain	43.2	39.2	11.4	10.4	5.2	4.4
Pacific	38.6	36.0	10.5	9.4	5.5 <sup>a</sup>	4.1
<b>COUNTY TYPE</b>						
Large Metro	37.7	36.5	10.9	10.5	5.6	5.3
Small Metro	39.3	39.4	11.0	11.1	5.5	5.8
250K – 1 Mil. Pop.	38.5	39.0	11.0	10.8	5.2	5.6
< 250K Pop.	40.9	40.3	11.1	11.8	6.1	6.1
Nonmetro	39.2	37.3	9.6	9.5	4.7	5.1
Urbanized	39.4	39.3	9.9	10.1	4.9	4.8
Less Urbanized	37.7	36.2	9.7	9.7	4.9	5.6
Completely Rural	44.5 <sup>b</sup>	33.8	7.7	6.1	3.3	3.9

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

<sup>a</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.