

Table 2.77A Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Numbers in Thousands, 2009 and 2010

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	20,102 ^a	19,222	17,066	16,486	10,403	9,998
GENDER						
Male	10,538	10,059	8,926	8,660	5,595	5,552
Female	9,564	9,164	8,140	7,826	4,808 ^a	4,446
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	16,171 ^b	15,257	13,865 ^a	13,180	8,575	8,127
White	12,342 ^b	11,563	10,856 ^b	10,168	6,869	6,475
Black or African American	2,636	2,543	2,058	2,061	1,170	1,151
American Indian or Alaska Native	124	113	91	87	49	47
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	612	582	501	488	248	239
Two or More Races	394	412	305	339	201	185
Hispanic or Latino	3,931	3,965	3,201	3,306	1,828	1,872
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	6,340	6,009	5,565	5,285	3,640	3,533
Female, White, Not Hispanic	6,001 ^a	5,554	5,291 ^a	4,883	3,229 ^a	2,942
Male, Black, Not Hispanic	1,371	1,291	1,072	1,039	634	616
Female, Black, Not Hispanic	1,265	1,252	986	1,022	537	535
Male, Hispanic	2,194	2,160	1,795	1,829	1,015	1,104
Female, Hispanic	1,737	1,805	1,406	1,476	813	767

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.77B Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Percentages, 2009 and 2010

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	52.5 ^b	50.5	44.6 ^a	43.3	27.2	26.3
GENDER						
Male	53.6 ^b	51.3	45.4	44.1	28.5	28.3
Female	51.4 ^a	49.8	43.8	42.5	25.8 ^a	24.1
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	52.2 ^b	50.2	44.8 ^a	43.4	27.7	26.8
White	54.6 ^b	52.3	48.0 ^b	46.0	30.4	29.3
Black or African American	46.0	45.2	35.9	36.6	20.4	20.4
American Indian or Alaska Native	55.3	54.4	40.6	41.9	22.0	22.9
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	39.6	37.6	32.4	31.5	16.1	15.4
Two or More Races	53.7	53.8	41.7	44.2	27.5	24.2
Hispanic or Latino	54.0	51.6	44.0	43.1	25.1	24.4
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	54.6 ^a	52.6	47.9	46.3	31.3	30.9
Female, White, Not Hispanic	54.6 ^b	52.0	48.1 ^a	45.7	29.4	27.5
Male, Black, Not Hispanic	47.2	45.8	36.9	36.9	21.8	21.9
Female, Black, Not Hispanic	44.7	44.5	34.8	36.3	19.0	19.0
Male, Hispanic	58.1 ^a	53.4	47.6	45.2	26.9	27.3
Female, Hispanic	49.5	49.7	40.1	40.7	23.2	21.1

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.78A Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Numbers in Thousands, 2009 and 2010

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	188,443	190,073	151,075	151,910	120,218	121,344
GENDER						
Male	94,370	95,454	77,467	77,039	64,860	65,282
Female	94,073	94,619	73,608	74,871	55,357	56,062
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	165,457	166,464	132,986	133,596	107,049	107,855
White	135,798	135,994	110,005	110,196	89,422	89,968
Black or African American	19,831	20,371	15,249	15,399	11,701	11,796
American Indian or Alaska Native	849	*	553	530	405	393
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	6,355	6,658	5,093	5,427	3,898	4,157
Two or More Races	2,026	2,107	1,566	1,656	1,205	1,251
Hispanic or Latino	22,986	23,608	18,089	18,313	13,169	13,489
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	67,235	67,426	55,044	54,806	47,174	47,654
Female, White, Not Hispanic	68,563	68,568	54,961	55,390	42,248	42,315
Male, Black, Not Hispanic	9,325	9,366	7,464	7,324	6,141	6,042
Female, Black, Not Hispanic	10,506	11,004	7,785	8,075	5,560	5,754
Male, Hispanic	12,851	13,200	10,696	10,567	8,170	8,187
Female, Hispanic	10,134	10,409	7,393	7,746	4,999	5,302

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.78B Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Percentages, 2009 and 2010

Demographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	88.2	88.2	70.7	70.5	56.3	56.3
GENDER						
Male	91.9	92.0	75.5	74.2	63.2	62.9
Female	84.8	84.7	66.4	67.0	49.9	50.2
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	89.5	89.3	71.9	71.6	57.9	57.8
White	92.3	91.9	74.7	74.5	60.8	60.8
Black or African American	81.5	82.8	62.7	62.6	48.1	47.9
American Indian or Alaska Native	84.9	*	55.3	53.1	40.5	39.4
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	67.0	67.2	53.7	54.8	41.1	42.0
Two or More Races	91.2	87.4	70.5	68.7	54.3	51.9
Hispanic or Latino	80.2	81.2	63.1	63.0	45.9	46.4
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	94.8	94.4	77.6	76.8	66.5	66.7
Female, White, Not Hispanic	89.9	89.6	72.1	72.4	55.4	55.3
Male, Black, Not Hispanic	86.8	85.5	69.5	66.9	57.2	55.2
Female, Black, Not Hispanic	77.3	80.6	57.3	59.2	40.9	42.1
Male, Hispanic	87.4	88.6	72.7	71.0	55.5	55.0
Female, Hispanic	72.6	73.3	53.0	54.5	35.8	37.3

*Low precision; no estimate reported.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.79A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Numbers in Thousands, 2009 and 2010

Demographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	10,403	9,998	6,912 ^a	6,467	2,058	1,958
GENDER						
Male	5,595	5,552	4,025	3,889	1,371	1,318
Female	4,808 ^a	4,446	2,887 ^a	2,579	687	640
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	8,575	8,127	5,663 ^a	5,236	1,773	1,649
White	6,869	6,475	4,745 ^a	4,388	1,593	1,480
Black or African American	1,170	1,151	581	559	100	96
American Indian or Alaska Native	49	47	38	38	12	13
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	3
Asian	248	239	141	120	18	18
Two or More Races	201	185	134	116	45	39
Hispanic or Latino	1,828	1,872	1,249	1,231	285	308
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	3,640	3,533	2,729	2,571	1,062	974
Female, White, Not Hispanic	3,229 ^a	2,942	2,016	1,817	531	506
Male, Black, Not Hispanic	634	616	324	331	71	67
Female, Black, Not Hispanic	537	535	257	229	30	29
Male, Hispanic	1,015	1,104	779	800	189	226
Female, Hispanic	813	767	470	431	96	82

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.79B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Demographic Characteristics: Percentages, 2009 and 2010

Demographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	27.2	26.3	18.1 ^a	17.0	5.4	5.1
GENDER						
Male	28.5	28.3	20.5	19.8	7.0	6.7
Female	25.8 ^a	24.1	15.5 ^a	14.0	3.7	3.5
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	27.7	26.8	18.3	17.2	5.7	5.4
White	30.4	29.3	21.0	19.8	7.0	6.7
Black or African American	20.4	20.4	10.1	9.9	1.7	1.7
American Indian or Alaska Native	22.0	22.9	16.8	18.4	5.2	6.4
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	2.8
Asian	16.1	15.4	9.1	7.8	1.2	1.2
Two or More Races	27.5	24.2	18.3	15.1	6.1	5.1
Hispanic or Latino	25.1	24.4	17.1	16.0	3.9	4.0
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	31.3	30.9	23.5	22.5	9.1	8.5
Female, White, Not Hispanic	29.4	27.5	18.3	17.0	4.8	4.7
Male, Black, Not Hispanic	21.8	21.9	11.2	11.7	2.4	2.4
Female, Black, Not Hispanic	19.0	19.0	9.1	8.1	1.0	1.0
Male, Hispanic	26.9	27.3	20.6	19.8	5.0	5.6
Female, Hispanic	23.2	21.1	13.4	11.9	2.7	2.3

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.80A Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Numbers in Thousands, 2009 and 2010

Demographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	120,218	121,344	52,650	52,082	15,071	14,973
GENDER						
Male	64,860	65,282	34,629	34,283	11,233	11,173
Female	55,357	56,062	18,021	17,799	3,838	3,801
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	107,049	107,855	44,901	44,068	13,479	13,404
White	89,422	89,968	37,416	36,355	11,814	11,599
Black or African American	11,701	11,796	5,376	5,415	1,262	1,262
American Indian or Alaska Native	405	393	234	260	90	69
Native Hawaiian or Other Pacific Islander	*	*	*	*	25	*
Asian	3,898	4,157	1,085	1,297	144	260
Two or More Races	1,205	1,251	578	568	143	145
Hispanic or Latino	13,169	13,489	7,749	8,015	1,592	1,569
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	47,174	47,654	24,232	23,973	8,635	8,730
Female, White, Not Hispanic	42,248	42,315	13,183	12,382	3,179	2,868
Male, Black, Not Hispanic	6,141	6,042	3,419	3,260	931	850
Female, Black, Not Hispanic	5,560	5,754	1,957	2,155	331	411
Male, Hispanic	8,170	8,187	5,580	5,561	1,351	1,172
Female, Hispanic	4,999	5,302	2,169	2,453	241	397

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.80B Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Demographic Characteristics: Percentages, 2009 and 2010

Demographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	56.3	56.3	24.7	24.2	7.1	6.9
GENDER						
Male	63.2	62.9	33.7	33.0	10.9	10.8
Female	49.9	50.2	16.2	15.9	3.5	3.4
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	57.9	57.8	24.3	23.6	7.3	7.2
White	60.8	60.8	25.4	24.6	8.0	7.8
Black or African American	48.1	47.9	22.1	22.0	5.2	5.1
American Indian or Alaska Native	40.5	39.4	23.4	26.0	9.0	7.0
Native Hawaiian or Other Pacific Islander	*	*	*	*	3.6	*
Asian	41.1	42.0	11.4	13.1	1.5	2.6
Two or More Races	54.3	51.9	26.0	23.6	6.5	6.0
Hispanic or Latino	45.9	46.4	27.0	27.5	5.6	5.4
GENDER/RACE/HISPANIC ORIGIN						
Male, White, Not Hispanic	66.5	66.7	34.2	33.6	12.2	12.2
Female, White, Not Hispanic	55.4	55.3	17.3	16.2	4.2	3.7
Male, Black, Not Hispanic	57.2	55.2	31.8	29.8	8.7	7.8
Female, Black, Not Hispanic	40.9	42.1	14.4	15.8	2.4	3.0
Male, Hispanic	55.5	55.0	37.9	37.3	9.2	7.9
Female, Hispanic	35.8	37.3	15.5	17.3	1.7	2.8

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly the Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.81A (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	20,145 ^a	19,196	17,081	16,434	10,392	9,934
GEOGRAPHIC DIVISION						
Northeast	3,743	3,542	3,313	3,140	2,158	1,987
New England	942	912	857	798	561	518
Middle Atlantic	2,802	2,630	2,455	2,342	1,597	1,469
Midwest	4,540 ^a	4,242	3,840	3,640	2,336	2,204
East North Central	3,212 ^a	2,948	2,730	2,536	1,662	1,505
West North Central	1,327	1,294	1,110	1,104	674	698
South	7,117	7,026	5,915	5,936	3,560	3,458
South Atlantic	3,567	3,604	2,989	3,047	1,791	1,794
East South Central	1,186	1,125	963	925	541	503
West South Central	2,364	2,297	1,963	1,963	1,227	1,161
West	4,745	4,386	4,013	3,718	2,339	2,285
Mountain	1,403	1,368	1,172	1,131	694	687
Pacific	3,341	3,018	2,841	2,588	1,644	1,598
COUNTY TYPE						
Large Metro	10,472	9,936	8,896	8,617	5,355	5,212
Small Metro	6,362	6,231	5,417	5,275	3,418	3,228
250K – 1 Mil. Pop.	4,048	4,001	3,444	3,381	2,110	2,051
< 250K Pop.	2,315	2,230	1,973	1,894	1,308	1,177
Nonmetro	3,311	3,029	2,767	2,542	1,620	1,494
Urbanized	1,532 ^b	1,219	1,297 ^b	1,033	751	639
Less Urbanized	1,502	1,574	1,233	1,317	722	737
Completely Rural	277	236	237	192	147	119

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.81B (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	52.7 ^b	50.5	44.7 ^a	43.3	27.2	26.2
GEOGRAPHIC DIVISION						
Northeast	55.2	53.5	48.8	47.4	31.8	30.0
New England	56.1	53.9	51.1	47.2	33.4	30.6
Middle Atlantic	54.9	53.3	48.1	47.5	31.3	29.8
Midwest	53.6 ^b	50.7	45.3	43.5	27.6	26.4
East North Central	54.2 ^b	50.4	46.0 ^a	43.3	28.0 ^a	25.7
West North Central	52.2	51.6	43.7	44.1	26.5	27.9
South	51.2	49.9	42.6	42.2	25.6	24.6
South Atlantic	50.9	50.7	42.6	42.9	25.6	25.2
East South Central	51.3	49.4	41.6	40.6	23.4	22.1
West South Central	51.7	49.0	42.9	41.9	26.9	24.8
West	52.2 ^a	49.1	44.2	41.7	25.7	25.6
Mountain	50.7	50.0	42.4	41.4	25.1	25.1
Pacific	52.8 ^a	48.7	44.9	41.8	26.0	25.8
COUNTY TYPE						
Large Metro	52.3 ^b	49.4	44.4	42.8	26.7	25.9
Small Metro	52.9	52.1	45.1	44.1	28.4	27.0
250K – 1 Mil. Pop.	51.6	51.4	43.9	43.5	26.9	26.4
< 250K Pop.	55.4	53.3	47.2	45.3	31.3	28.2
Nonmetro	53.5	51.3	44.7	43.1	26.2	25.3
Urbanized	55.3 ^a	50.7	46.8	43.0	27.1	26.6
Less Urbanized	52.5	52.0	43.1	43.5	25.2	24.3
Completely Rural	50.0	50.1	42.6	40.8	26.4	25.2

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

**Table 2.82A (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics:
Numbers in Thousands, 2009 and 2010**

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	188,388	190,068	151,055	151,947	120,250	121,440
GEOGRAPHIC DIVISION						
Northeast	35,715	36,082	29,857	29,870	24,246	24,957
New England	9,792	9,783	8,309	8,308	6,947	7,224
Middle Atlantic	25,924	26,300	21,548	21,562	17,300	17,733
Midwest	42,407	43,046	34,527	35,110	27,870	28,084
East North Central	29,444	29,831	23,711	24,433	19,099	19,684
West North Central	12,964	13,215	10,817	10,676	8,771	8,399
South	67,058	67,806	52,157	52,086	40,173	40,711
South Atlantic	36,262	36,254	28,701	28,638	22,094	22,749
East South Central	10,592	10,699	7,350	7,432	5,489	5,661
West South Central	20,204	20,853	16,106	16,016	12,589	12,301
West	43,208	43,133	34,512	34,881	27,961	27,688
Mountain	13,195	13,055	10,380	10,273	8,482	8,254
Pacific	30,013	30,078	24,133	24,607	19,479	19,434
COUNTY TYPE						
Large Metro	100,661	102,835	83,742	84,634	67,302	68,245
Small Metro	57,333	57,350	44,867	45,298	35,615	36,026
250K – 1 Mil. Pop.	38,039	38,300	29,632	30,573	23,649	24,318
< 250K Pop.	19,294	19,050	15,235	14,725	11,966	11,708
Nonmetro	30,395	29,882	22,446	22,015	17,333	17,170
Urbanized	13,536	13,051	10,469	9,717	8,095	7,832
Less Urbanized	13,728	14,123	9,969	10,388	7,640	7,867
Completely Rural	3,130	2,708	2,009	1,910	1,599	1,471

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.82B (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Lifetime (2009)	Lifetime (2010)	Past Year (2009)	Past Year (2010)	Past Month (2009)	Past Month (2010)
TOTAL	88.2	88.1	70.7	70.5	56.3	56.3
GEOGRAPHIC DIVISION						
Northeast	90.2	90.4	75.4	74.8	61.2	62.5
New England	93.2	92.7	79.1	78.7	66.2	68.4
Middle Atlantic	89.1	89.6	74.1	73.4	59.5	60.4
Midwest	90.8	91.6	73.9	74.7	59.7	59.8
East North Central	90.5	91.3	72.9	74.8	58.7	60.2
West North Central	91.6	92.4	76.4	74.6	61.9	58.7
South	85.8	85.9	66.7	66.0	51.4	51.6
South Atlantic	87.2	86.5	69.1	68.4	53.2	54.3
East South Central	83.6	83.6	58.0	58.1	43.3	44.2
West South Central	84.5	86.2	67.3	66.2	52.6	50.8
West	88.0	86.5	70.3	70.0	56.9	55.5
Mountain	87.7	85.3	69.0	67.1	56.4	53.9
Pacific	88.1	87.1	70.8	71.2	57.2	56.3
COUNTY TYPE						
Large Metro	88.5	89.0	73.6	73.2	59.2	59.0
Small Metro	88.6	87.6	69.3	69.2	55.0	55.0
250K – 1 Mil. Pop.	88.9	87.6	69.2	69.9	55.3	55.6
< 250K Pop.	88.0	87.4	69.5	67.6	54.6	53.7
Nonmetro	86.7	86.5	64.0	63.7	49.4	49.7
Urbanized	88.6	88.9	68.5	66.2	53.0	53.4
Less Urbanized	84.9	85.0	61.7	62.5	47.3	47.3
Completely Rural	86.2	83.3	55.3	58.7	44.0	45.2

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.83A (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	10,392	9,934	6,945 ^a	6,423	2,063	1,940
GEOGRAPHIC DIVISION						
Northeast	2,158	1,987	1,405	1,282	402	397
New England	561	518	392	340	137	103
Middle Atlantic	1,597	1,469	1,014	943	265	294
Midwest	2,336	2,204	1,654 ^a	1,482	510	453
East North Central	1,662	1,505	1,194 ^b	995	369	300
West North Central	674	698	460	487	142	153
South	3,560	3,458	2,311	2,160	709	687
South Atlantic	1,791	1,794	1,143	1,071	363	343
East South Central	541	503	367	340	110	107
West South Central	1,227	1,161	802	749	236	237
West	2,339	2,285	1,575	1,499	441	403
Mountain	694	687	462	453	132	118
Pacific	1,644	1,598	1,113	1,045	309	285
COUNTY TYPE						
Large Metro	5,355	5,212	3,475	3,292	1,004	954
Small Metro	3,418	3,228	2,314	2,106	715	641
250K – 1 Mil. Pop.	2,110	2,051	1,399	1,292	390	379
< 250K Pop.	1,308	1,177	915	814	325	261
Nonmetro	1,620	1,494	1,156	1,025	344	345
Urbanized	751	639	536	433	173	152
Less Urbanized	722	737	510	508	140	173
Completely Rural	147	119	109	84	32	21

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.83B (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 20, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	27.2	26.2	18.2 ^a	16.9	5.4	5.1
GEOGRAPHIC DIVISION						
Northeast	31.8	30.0	20.7	19.4	5.9	6.0
New England	33.4	30.6	23.3	20.1	8.2	6.1
Middle Atlantic	31.3	29.8	19.8	19.1	5.2	6.0
Midwest	27.6	26.4	19.5 ^a	17.7	6.0	5.4
East North Central	28.0 ^a	25.7	20.1 ^b	17.0	6.2	5.1
West North Central	26.5	27.9	18.1	19.4	5.6	6.1
South	25.6	24.6	16.6	15.4	5.1	4.9
South Atlantic	25.6	25.2	16.3	15.1	5.2	4.8
East South Central	23.4	22.1	15.9	14.9	4.8	4.7
West South Central	26.9	24.8	17.5	16.0	5.2	5.1
West	25.7	25.6	17.3	16.8	4.9	4.5
Mountain	25.1	25.1	16.7	16.6	4.8	4.3
Pacific	26.0	25.8	17.6	16.9	4.9	4.6
COUNTY TYPE						
Large Metro	26.7	25.9	17.3	16.4	5.0	4.7
Small Metro	28.4	27.0	19.3	17.6	5.9	5.4
250K – 1 Mil. Pop.	26.9	26.4	17.9	16.6	5.0	4.9
< 250K Pop.	31.3	28.2	21.9	19.5	7.8	6.2
Nonmetro	26.2	25.3	18.7	17.4	5.6	5.9
Urbanized	27.1	26.6	19.4	18.0	6.2	6.3
Less Urbanized	25.2	24.3	17.8	16.8	4.9	5.7
Completely Rural	26.4	25.2	19.7	17.9	5.7	4.4

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.84A (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Numbers in Thousands, 2009 and 2010

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	120,250	121,440	52,744	52,112	15,111	14,959
GEOGRAPHIC DIVISION						
Northeast	24,246	24,957	10,067	10,252	2,991	2,716
New England	6,947	7,224	2,695	2,762	835	808
Middle Atlantic	17,300	17,733	7,372	7,490	2,156	1,909
Midwest	27,870	28,084	13,011	12,558	3,860	3,627
East North Central	19,099	19,684	8,905	8,782	2,646	2,517
West North Central	8,771	8,399	4,106	3,776	1,214	1,110
South	40,173	40,711	17,922	17,089	5,230	5,106
South Atlantic	22,094	22,749	9,169	8,842	2,599	2,744
East South Central	5,489	5,661	2,655	2,533	902	760
West South Central	12,589	12,301	6,099	5,714	1,729	1,602
West	27,961	27,688	11,745	12,213	3,031	3,509
Mountain	8,482	8,254	3,676	3,867	929	1,074
Pacific	19,479	19,434	8,070	8,346	2,102	2,436
COUNTY TYPE						
Large Metro	67,302	68,245	28,730	28,668	7,639	7,693
Small Metro	35,615	36,026	15,739	15,506	4,955	4,864
250K – 1 Mil. Pop.	23,649	24,318	9,981	10,177	3,200	3,148
< 250K Pop.	11,966	11,708	5,758	5,329	1,755	1,716
Nonmetro	17,333	17,170	8,275	7,939	2,517	2,401
Urbanized	8,095	7,832	3,760	3,566	1,151	1,056
Less Urbanized	7,640	7,867	3,840	3,725	1,140	1,094
Completely Rural	1,599	1,471	676	647	226	251

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.

Table 2.84B (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 21 or Older, by Geographic Characteristics: Percentages, 2009 and 2010

Geographic Characteristic	Alcohol Use (2009)	Alcohol Use (2010)	Binge Alcohol Use (2009)	Binge Alcohol Use (2010)	Heavy Alcohol Use (2009)	Heavy Alcohol Use (2010)
TOTAL	56.3	56.3	24.7	24.2	7.1	6.9
GEOGRAPHIC DIVISION						
Northeast	61.2	62.5	25.4	25.7	7.6	6.8
New England	66.2	68.4	25.7	26.2	7.9	7.7
Middle Atlantic	59.5	60.4	25.3	25.5	7.4	6.5
Midwest	59.7	59.8	27.9	26.7	8.3	7.7
East North Central	58.7	60.2	27.4	26.9	8.1	7.7
West North Central	61.9	58.7	29.0	26.4	8.6	7.8
South	51.4	51.6	22.9	21.7	6.7	6.5
South Atlantic	53.2	54.3	22.1	21.1	6.3	6.5
East South Central	43.3	44.2	20.9	19.8	7.1	5.9
West South Central	52.6	50.8	25.5	23.6	7.2	6.6
West	56.9	55.5	23.9	24.5	6.2	7.0
Mountain	56.4	53.9	24.4	25.3	6.2	7.0
Pacific	57.2	56.3	23.7	24.2	6.2	7.0
COUNTY TYPE						
Large Metro	59.2	59.0	25.3	24.8	6.7	6.7
Small Metro	55.0	55.0	24.3	23.7	7.7	7.4
250K – 1 Mil. Pop.	55.3	55.6	23.3	23.3	7.5	7.2
< 250K Pop.	54.6	53.7	26.3	24.5	8.0	7.9
Nonmetro	49.4	49.7	23.6	23.0	7.2	6.9
Urbanized	53.0	53.4	24.6	24.3	7.5	7.2
Less Urbanized	47.3	47.3	23.8	22.4	7.1	6.6
Completely Rural	44.0	45.2	18.6	19.9	6.2	7.7

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2010 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

^a Difference between estimate and 2010 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2010 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality (formerly Office of Applied Studies), National Survey on Drug Use and Health, 2009 and 2010.